

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PCT/03 / 0 0 8 5 4

27-05-2003

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0201662-4
Patent application number

(86) Ingivningsdatum 2002-05-31
Date of filing

REC'D 20 JUN 2003

WIPO PCT

Stockholm, 2003-03-11

För Patent- och registreringsverket
For the Patent- and Registration Office

Hjördis Segerlund
Hjördis Segerlund

Avgift
Fee 170:-

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

PHARMACEUTICAL COMBINATION

Field of the Invention

- 5 This invention relates to a new combination of pharmaceutically-active compounds. In particular the invention relates to a combination of thrombin inhibitor of a particular class or a pharmaceutically-acceptable derivative thereof and certain antiarrhythmic oxabispindines or pharmaceutically acceptable salts thereof.

10 Background to the Invention

- Atrial fibrillation (AF) is characterised by grossly disorganised atrial electrical activity that is irregular in respect of both rate and rhythm. Patients with AF have no visually discernible timing pattern in atrial electrical activity when measured by surface ECG, or in
15 electrogram sequences recorded by catheter electrodes.

- During AF, the regular pumping action of the atria is replaced by irregular, disorganised and quivering spasms of atrial tissue. These spasms may be experienced as irregular heartbeat, palpitations, discomfort, dizziness and/or angina pectoris. Further, the
20 inefficient pumping action of the heart tends to lead to significant morbidity related to reduced blood flow. More seriously, the reduced cardiac output can lead to blood pooling in the left atria and the formation of blood clots. Blood clots, mostly originating in the left atrium, can dislodge and travel through the bloodstream to organs, e.g. the brain, spleen, kidneys etc. If the clot travels to the brain, this may result in cerebral stroke and even
25 death.

In the US alone, AF affects an estimated two million people, with approximately 160,000 new cases being diagnosed each year. It has been estimated that AF is responsible for over 70,000 strokes each year in the US, and that the cost of treating these patients is more than

- R^{5a} to R^{5d} independently represent, at each occurrence, H, C_{1-6} alkyl (which latter group is optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro, aryl and Het²), aryl or Het³, or R^{5d} , together with R^8 , represents C_{3-6} alkylene (which alkylene group is optionally interrupted by an O atom and/or is optionally substituted by one or more C_{1-3} alkyl groups);
- R^6 represents H, C_{1-6} alkyl (optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, $-C(O)R^{10a}$, $-C(O)OR^{10b}$ or $-C(O)N(H)R^{10c}$;
- R^{10a} , R^{10b} and R^{10c} independently represent C_{1-6} alkyl (optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, or R^{10a} represents H;
- R^7 represents C_{1-12} alkyl (optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro, aryl, C_{1-6} alkoxy and Het⁴);
- R^8 represents H, C_{1-12} alkyl, C_{1-6} alkoxy (which latter two groups are optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl and C_{1-4} alkoxy), -D-aryl, -D-aryloxy, -D-Het⁵, -D-N(H)C(O) R^{11a} , -D-S(O)₂ R^{12a} , -D-C(O) R^{11b} , -D-C(O)OR^{12b}, -D-C(O)N(R^{11c}) R^{11d} , or R^8 , together with R^{5d} , represents C_{3-6} alkylene (which alkylene group is optionally interrupted by an O atom and/or is optionally substituted by one or more C_{1-3} alkyl groups);
- R^{11a} to R^{11d} independently represent H, C_{1-6} alkyl (optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, or R^{11c} and R^{11d} together represent C_{3-6} alkylene;
- R^9 , R^{12a} and R^{12b} independently represent C_{1-6} alkyl (optionally substituted and/or terminated by one or more substituents selected from -OH, halo, cyano, nitro and aryl) or aryl;
- D represents a direct bond or C_{1-6} alkylene;
- X represents O or S;

R^2 represents H, halo, C_{1-6} alkyl, $-OR^{13}$, $-E-N(R^{14})R^{15}$ or, together with R^3 , represents $=O$;

R^3 represents H, C_{1-6} alkyl or, together with R^2 , represents $=O$;

R^{13} represents H, C_{1-6} alkyl, $-E$ -aryl, $-E$ -Het⁶, $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or

$-C(O)N(R^{17a})R^{17b}$;

R^{14} represents H, C_{1-6} alkyl, $-E$ -aryl, $-E$ -Het⁶, $-C(O)R^{16a}$, $-C(O)OR^{16b}$,

$-S(O)_2R^{16c}$, $-[C(O)]_pN(R^{17a})R^{17b}$ or $-C(NH)NH_2$;

R^{15} represents H, C_{1-6} alkyl, $-E$ -aryl or $-C(O)R^{16d}$;

R^{16a} to R^{16d} independently represent, at each occurrence when used herein, C_{1-6} alkyl

(optionally substituted and/or terminated by one or more substituents selected from halo,

aryl and Het⁷), aryl, Het⁸, or R^{16a} and R^{16d} independently represent H;

R^{17a} and R^{17b} independently represent, at each occurrence when used herein, H or C_{1-6} alkyl

(optionally substituted and/or terminated by one or more substituents selected from halo,

aryl and Het⁹), aryl, Het¹⁰, or together represent C_{3-6} alkylene, optionally interrupted by an

O atom;

E represents, at each occurrence when used herein, a direct bond or

C_{1-4} alkylene;

p represents 1 or 2;

Het¹ to Het¹⁰ independently represent five- to twelve-membered heterocyclic groups

containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, which

groups are optionally substituted by one or more substituents selected from $-OH$, oxo, halo,

cyano, nitro,

C_{1-6} alkyl, C_{1-6} alkoxy, aryl, aryloxy, $-N(R^{18a})R^{18b}$, $-C(O)R^{18c}$, $-C(O)OR^{18d}$, -

$C(O)N(R^{18e})R^{18f}$, $-N(R^{18g})C(O)R^{18h}$ and $-N(R^{18i})S(O)_2R^{18j}$;

R^{18a} to R^{18j} independently represent C_{1-6} alkyl, aryl or R^{18a} to R^{18i} independently represent

H;

A represents a direct bond, $-J-$, $-J-N(R^{19})-$ or $-J-O-$ (in which latter two groups, $N(R^{19})-$ or

O- is attached to the carbon atom bearing R^2 and R^3);

B represents $-Z-$, $-Z-N(R^{20})-$, $-N(R^{20})-Z-$, $-Z-S(O)_n-$, $-Z-O-$ (in which latter two groups, Z is attached to the carbon atom bearing R^2 and R^3),

$-N(R^{20})C(O)O-Z-$, (in which latter group, $-N(R^{20})$ is attached to the carbon atom bearing R^2 and R^3) or $-C(O)N(R^{20})-$ (in which latter group,

5 $-C(O)$ is attached to the carbon atom bearing R^2 and R^3);

J represents C_{1-6} alkylene optionally substituted by one or more substituents selected from $-OH$, halo and amino;

Z represents a direct bond or C_{1-4} alkylene;

n represents 0, 1 or 2;

10 R^{19} and R^{20} independently represent H or C_{1-6} alkyl;

G represents CH or N;

R^4 represents one or more optional substituents selected from $-OH$, cyano, halo, nitro, C_{1-6} alkyl (optionally terminated by $-N(H)C(O)OR^{21a}$),

C_{1-6} alkoxy, $-N(R^{22a})R^{22b}$, $-C(O)R^{22c}$, $-C(O)OR^{22d}$, $-C(O)N(R^{22e})R^{22f}$, $-N(R^{22g})C(O)R^{22h}$, $-N(R^{22i})C(O)N(R^{22j})R^{22k}$, $-N(R^{22m})S(O)_2R^{21b}$, $-S(O)_2R^{21c}$, and/or $-OS(O)_2R^{21d}$;

R^{21a} to R^{21d} independently represent C_{1-6} alkyl;

20 R^{22a} and R^{22b} independently represent H, C_{1-6} alkyl or together represent C_{3-6} alkylene, resulting in a four- to seven-membered nitrogen-containing ring;

R^{22c} to R^{22m} independently represent H or C_{1-6} alkyl; and

R^{41} to R^{46} independently represent H or C_{1-3} alkyl;

25 wherein each aryl and aryloxy group, unless otherwise specified, is optionally substituted;

provided that

(a) the compound is not:

30 3,7-dibenzoyl-9-oxa-3,7-diazabicyclo[3.3.1]nonane;

(b) when A represents $-J-N(R^{19})-$ or $-J-O-$, then:

(i) J does not represent C_1 alkylene; and

(ii) B does not represent $-N(R^{20})-$, $-N(R^{20})-Z-$ (in which latter group $N(R^{20})$ is attached to the carbon atom bearing R^2 and R^3),

5 $-S(O)_n-$, $-O-$ or $-N(R^{20})C(O)O-Z-$ when R^2 and R^3 do not together represent $=O$; and

(c) when R^2 represents $-OR^{13}$ or $-N(R^{14})(R^{15})$, then:

(i) A does not represent $-J-N(R^{19})-$ or $-J-O-$; and

(ii) B does not represent $-N(R^{20})-$, $-N(R^{20})-Z-$ (in which latter group $N(R^{20})$ is attached to the carbon atom bearing R^2 and R^3),

10 $-S(O)_n-$, $-O-$ or $-N(R^{20})C(O)O-Z-$;

or a pharmaceutically acceptable derivative thereof.

This definition will hereinafter be referred to as a compound as defined in claim 1 of WO
15 01/28992. The definition of "a pharmaceutically acceptable derivative thereof" is that used
in WO 01/28992 which is now repeated. Pharmaceutically acceptable derivatives include
salts and solvates. Salts which may be mentioned include acid addition salts. Specific
salts that may be mentioned include arylsulfonate salts, such as toluenesulfonate and,
especially, benzenesulfonate salts. Solvates that may be mentioned include hydrates, such
20 as monohydrates of the compounds of the invention.

Pharmaceutically acceptable derivatives also include, at the oxabispidine or (when G
represents N) pyridyl nitrogens, C_{1-4} alkyl quaternary ammonium salts and N-oxides,
provided that when a N-oxide is present:

25 no Het (Het^1 , Het^2 , Het^3 , Het^4 , Het^5 , Het^6 , Het^7 , Het^8 , Het^9 and Het^{10}) group contains an
unoxidised S-atom; and/or

n does not represent 0 when B represents $-Z-S(O)_n-$.

The compounds of the invention may exhibit tautomerism. All tautomeric forms and
30 mixtures thereof are included within the scope of the invention.

Claim 34 of WO 01/28992 provides a list of compounds as follows

A compound which is:

- 4-{2-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzonitrile;
- 5 7-[4-(4-cyanophenyl)-4-(3,4-dimethoxyphenoxy)butyl]-*N*-ethyl-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxamide;
- 4-({3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile;
- 4-{3-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-
10 hydroxypropoxy}benzonitrile;
- 4-(2-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;
- 4-(((2*S*)-2-amino-3-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)oxy)benzonitrile;
- 15 *tert*-butyl 2-{7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate;
- tert*-butyl 2-{7-[4-(4-cyanophenyl)butyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate;
- tert*-butyl 2-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate;
- 20 4-(2-{7-[4-(4-pyridinyl)butyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-ethoxy)benzonitrile
- tert*-butyl 2-{7-[4-(4-pyridinyl)butyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate;
- 4-{3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile;
- 25 4-{3-[7-(3,4-dimethoxyphenethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile;
- 4-{2-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-ethoxy}benzonitrile;
- 30 4-({3-[7-(butylsulfonyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-

amino)benzonitrile;

4-({3-[7-(3,4-dimethoxyphenethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl} amino)benzonitrile;

4-[4-[7-(butylsulfonyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile;

4-{1-(3,4-dimethoxyphenoxy)-4-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]butyl}benzonitrile;

4-[4-[7-(3,4-dimethoxyphenethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-1-(3,4-dimethoxyphenoxy)butyl]benzonitrile;

2-(4-acetyl-1-piperazinyl)ethyl 7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate;

7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-*N*-ethyl-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxamide;

4-{3-[7-(butylsulfonyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile;

2-(4-acetyl-1-piperazinyl)ethyl 7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate;

7-[2-(4-cyanophenoxy)ethyl]-*N*-ethyl-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxamide;

4-{2-[7-(butylsulfonyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile;

4-{2-[7-(3,4-dimethoxyphenethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile;

2-(4-acetyl-1-piperazinyl)ethyl 7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate;

7-[3-(4-cyanoanilino)propyl]-*N*-ethyl-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxamide;

2-(4-acetyl-1-piperazinyl)ethyl 7-[4-(4-cyanophenyl)-4-(3,4-dimethoxyphenoxy)butyl]-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate;

4-{3-[7-(cyclopropylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-

hydroxypropoxy}benzonitrile;

4-(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;

4-(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;

2-{7-[3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-*N*-isopropylacetamide;

4-(3-{7-[3-(ethylsulfonyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;

4-(2-hydroxy-3-{7-[2-(2-methoxyethoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile;

4-(2-hydroxy-3-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile;

4-({3-[7-(cyclopropylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile;

4-[(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-[(3-{7-[2-(4-methyl-1,3-thiazol-5-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-[(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

2-{7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-*N*-isopropylacetamide;

4-[(3-{7-[3-(ethylsulfonyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-[(3-{7-[2-(2-methoxyethoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-({3-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile;

4-[(3-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-

non-3-yl}propyl)amino]benzonitrile;

4-{2-[7-(cyclopropylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-ethoxy}benzonitrile;

4-(2-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-(2-{7-[2-(4-methyl-1,3-thiazol-5-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-(2-{7-[3-(4-acetyl-1-piperazinyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-*N*-isopropylacetamide;

4-(2-{7-[3-(ethylsulfonyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-ethoxy)benzonitrile;

4-(2-{7-[2-(2-methoxyethoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-(2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy)benzonitrile;

4-[(3-{7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-(cyclopropylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-[2-(4-methyl-1,3-thiazol-5-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-[3-(4-acetyl-1-piperazinyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

2-(7-{3-[(4-cyanophenyl)sulfonyl]propyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)-*N*-isopropylacetamide;

4-[(3-{7-[3-(ethylsulfonyl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-

propyl)sulfonyl]benzonitrile;

4-[(3-{7-[2-(2-methoxyethoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;

4-[(3-{7-[2-(4-fluorophenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-(2-{7-[2-(4-fluorophenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-{2-[7-(tetrahydro-2H-pyran-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}benzonitrile;

4-(3-{7-[2-(4-fluorophenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropoxy)benzonitrile;

4-(2-hydroxy-3-[7-(tetrahydro-2H-pyran-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy)benzonitrile;

4-[(3-{7-(2-fluoro-3,3-dimethylbutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-[(3-{7-(2-hydroxy-3,3-dimethylbutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-[(3-{7-(3,3-dimethylbutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-[(3-{7-(2-oxopropyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)amino]benzonitrile;

4-(2-{7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-

yl)ethyl)benzonitrile;

4-{4-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-
butyl}benzonitrile;

4-{2-[7-(2-phenoxyethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}-
5 benzonitrile;

2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-
N,N-diethylacetamide;

4-[(3-{7-[4-(4-fluorophenyl)-4-oxobutyl]-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl}propyl)amino]benzonitrile;

10 4-[(7-{3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)-
methyl]benzonitrile;

4-{2-[7-(2,4-difluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-
ethoxy}benzonitrile;

4-[(3-{7-[4-(difluoromethoxy)benzyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-
15 yl}propyl)amino]benzonitrile;

4-[(3-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-
propyl)amino]benzonitrile;

4-[(3-{7-[3-(4-bromophenyl)-3-oxopropyl]-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl}propyl)amino]benzonitrile;

20 4-{2-[7-(2,2-difluoroethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}-
benzonitrile;

4-[(3-{7-(2-phenoxyethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-
amino]benzonitrile;

4-(2-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)-
25 ethoxy)benzonitrile;

4-[(2*S*)-3-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-
diazabicyclo[3.3.1]non-3-yl}-2-hydroxypropyl)oxy]benzonitrile;

4-[(2*S*)-2-hydroxy-3-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}propyl)oxy]benzonitrile;

30 4-{2-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-

ethoxy}isophthalonitrile;

4-(2-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl}ethoxy)isophthalonitrile;

4-(2-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-
ethoxy)isophthalonitrile;

tert-butyl 2-{7-[2-(2,4-dicyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethylcarbamate;

4-({(2*S*)-2-amino-3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-
yl]propyl}oxy)benzonitrile;

4-(((2*S*)-2-amino-3-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diaza-
bicyclo[3.3.1]non-3-yl}propyl)oxy)benzonitrile;

4-{3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-
propoxy}benzonitrile;

4-(3-{7-[2-(4-fluorophenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-
3-yl}propoxy)benzonitrile;

4-(3-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-
propoxy)benzonitrile;

4-(4-{7-[2-(1*H*-pyrrol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-
butyl)benzonitrile;

4-{[(2*S*)-3-(7-{2-[4-(*tert*-butoxy)phenoxy]ethyl}-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl)-2-hydroxypropyl]oxy}benzonitrile;

4-(((2*S*)-3-{7-[2-(3,5-dimethyl-1*H*-pyrazol-1-yl)ethyl]-9-oxa-3,7-diaza-
bicyclo[3.3.1]non-3-yl}-2-hydroxypropyl)oxy)benzonitrile;

4-{3-[7-(imidazo[1,2-*a*]pyridin-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl]propoxy}benzonitrile;

4-{3-[7-(2-phenoxyethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}-
benzonitrile;

4-(3-{7-[2-(3,5-dimethyl-1*H*-pyrazol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}propoxy)benzonitrile;

4-({3-[7-(imidazo[1,2-*a*]pyridin-2-ylmethyl)-9-oxa-3,7-diazabicyclo-

- [3.3.1]non-3-yl}propyl}amino)benzonitrile;
4-({3-[7-(2,4-difluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-
propyl}amino)benzonitrile;
4-({3-[7-(2-[4-(*tert*-butoxy)phenoxy]ethyl)-9-oxa-3,7-diazabicyclo[3.3.1]-
5 non-3-yl}propyl}amino)benzonitrile;
4-[2-[7-(imidazo[1,2-a]pyridin-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-
yl]ethoxy)benzonitrile;
tert-butyl 2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl}ethylcarbamate;
10 4-({3-[7-(2-[4-(*tert*-butoxy)phenoxy]ethyl)-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl}propyl)sulfonyl}benzonitrile;
4-[(3-[7-[2-(3,5-dimethyl-1*H*-pyrazol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}propyl)sulfonyl]benzonitrile;
4-({3-[7-(2,4-difluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-
15 propyl)sulfonyl}benzonitrile;
4-[2-[7-(imidazo[1,2-a]pyridin-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl]ethoxy]isophthalonitrile;
4-[2-(7-{2-[4-(*tert*-butoxy)phenoxy]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]-
non-3-yl)ethoxy]isophthalonitrile;
20 4-(2-[7-[2-(3,5-dimethyl-1*H*-pyrazol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl]ethoxy)isophthalonitrile;
4-(4-[7-[2-(1*H*-imidazol-4-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-
yl}butyl)benzonitrile;
4-[4-[7-(imidazo[1,2-a]pyridin-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]-
25 non-3-yl]butyl}benzonitrile;
4-[4-[7-(2-phenoxyethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]butyl]-
benzonitrile;
4-(4-[7-[2-(3,5-dimethyl-1*H*-pyrazol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl]butyl)benzonitrile;

4-[3-(7-{2-oxo-2-[4-(1-pyrrolidinyl)phenyl]ethyl}-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl)propoxy]benzonitrile;

4-(3-{7-[2-(4-hydroxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl}propoxy)benzonitrile;

5 4-(3-{7-[2-(4-methylphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl}propoxy)benzonitrile;

4-(3-{7-[2-(4-methoxyphenyl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl}propoxy)benzonitrile;

10 4-(3-{7-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-oxoethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile;

4-(2-{7-[2-(2,6-dimethylphenoxy)-1-methylethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethoxy)benzonitrile;

4-(3-{7-[2-oxo-2-(3-oxo-3,4-dihydro-2H-1,4-benzoxazin-6-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propoxy)benzonitrile;

15 *tert*-butyl 2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl}ethylcarbamate;

N-(*tert*-butyl)-*N'*-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)urea;

20 *tert*-butyl 2-((7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl)methyl)-1-pyrrolidinecarboxylate;

4-[[3-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propyl]amino]-benzonitrile;

4-[[3-{7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl]amino]benzonitrile;

tert-butyl 2-{7-[2-(4-nitrophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-

25 non-3-yl}ethylcarbamate (*m/z* = 437);

tert-butyl 2-[7-(2-{4-[(methylsulfonyl)amino]phenoxy}ethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethylcarbamate;

tert-butyl 2-{7-[2-(4-aminophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl}ethylcarbamate;


4-((3-[7-(phenylsulfonyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl)-amino)benzonitrile; or

4-((3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl)amino)benzamide.

- 5 This list of compounds and including pharmaceutically acceptable derivatives of the compounds as defined in WO 01/28992 will hereinafter be referred to as a compound of Claim 34 of WO 01/28992.

PCT/SE02/00724 discloses modified release formulations of the following compounds
10 which are described in WO 01/28992:


(a) 4-((3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl)amino)benzonitrile:



15

which compound is referred to hereinafter as Compound A. Compound A is specifically disclosed in WO 01/28992 both in the form of the free base and in the form of a benzenesulphonate salt;

(b) *tert*-butyl 2-{7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethylcarbamate:




5

in the form of the free base, which compound is referred to hereinafter as Compound B;

(c) *tert*-butyl 2-{7-[4-(4-cyanophenyl)butyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethylcarbamate:


10



in the form of the free base, which compound is referred to hereinafter as Compound C;
and

15

(d) *tert*-butyl 2-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate:




in the form of the free base, which compound is referred to hereinafter as Compound D.

Current drug therapies for AF include antiarrhythmic drugs, administered with a view to re-establishing and maintaining a normal heartbeat or to controlling heart rate, and anticoagulant and/or thrombolytic drugs, administered with a view to preventing thromboembolism and/or cerebral stroke.

Coagulation is the result of a complex series of enzymatic reactions. One of the ultimate steps in this series of reactions is the conversion of the proenzyme prothrombin to the active enzyme thrombin.

Thrombin is known to play a central role in coagulation. It activates platelets, leading to platelet aggregation, converts fibrinogen into fibrin monomers, which polymerise spontaneously into fibrin polymers, and activates factor XIII, which in turn crosslinks the polymers to form insoluble fibrin. Furthermore, thrombin activates factor V and factor VIII leading to a "positive feedback" generation of thrombin from prothrombin.

International patent application PCT/SE01/02657 discloses thrombin-inhibiting compounds of formula



wherein


R^a represents $-OH$ or $-CH_2OH$;

R^1 represents at least one optional halo substituent;


R^2 represents one or two C_{1-3} alkoxy substituents, the alkyl parts of which substituents are themselves substituted with one or more fluoro substituents (i.e. R^2 represents one or two fluoroalkoxy(C_{1-3}) groups);

Y represents $-CH_2-$ or $-(CH_2)_2-$; and

R^3 represents a structural fragment of formula I(i) or I(ii):



I(i)



I(ii)

wherein

R^4 represents H or one or more fluoro substituents; and

one or two of X_1 , X_2 , X_3 and X_4 represent $-N-$ and the others represent $-CH-$,

or a pharmaceutically-acceptable derivative thereof in claim 1. Such compounds are hereinafter referred to as a compound of claim 1 in PCT/SE01/02657.

Claim 20 of PCT/SE01/02657 discloses the following compounds:

- Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Cl)(5-OCHF₂)-(S)CH(CH₂OH)C(O)-Aze-Pab;
 Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab;
 5 Ph(3-OCHF₂)-(R)CH(OH)-CO-Aze-Pab;
 Ph(3-OCF₃)-(R)CH(OH)-CO-Aze-Pab;
 Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab;
 10 Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Br)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab;
 Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab;
 15 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Pro-Pab;
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((2-amidino)-5-pyridinyl);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-amidino)-2-pyrimidinyl);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(3-F);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF);
 20 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF).
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OEt);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OnPr);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OiPr);
 25 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OcBu);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(COOcPentyl);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Z);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 30 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂-3-(5-Me-isoxazole));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂-3-pyridine);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OiBu);

- Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OEt);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(ONb);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OcHexyl);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OcBu);
 5 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂CH₂OPh(3-CF₃));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(ONb(4-Cl));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(ONb(3-MeO));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(ONb(2-Br));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(ONb(4-Me));
 10 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(O-4-heptyl);
 Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 15 Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl, 5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH);
 20 Ph(3-Cl, 5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OH);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Pro-Pab(OMe);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((2-methoxy-amidino)-5-pyridinyl);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-methoxy-amidino)-2-pyrimidinyl);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(OMe); or
 25 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(OMe).

Such compounds and pharmaceutically-acceptable derivatives of these compounds are hereinafter referred to as a compound of claim 20 in PCT/SE01/02657.

The following compounds represent sub-set 1 of the compounds of claim 20 of
 30 PCT/SE01/02657:

- Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF)(OMe);

Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe);

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab;

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab(OH);

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF);

5 Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-(S)Aze-Pab; or,

Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-(S)Aze-Pab(OH).

The following compounds represent sub-set 2 of the compounds of claim 20 of PCT/SE01/02657:

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF)(OMe) ;

10 Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe) ; or

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab.

The following compound represent sub-set 3 of the compounds of claim 20 of PCT/SE01/02657:

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF)(OMe).

15


Combinations of a compound from any one of sub-sets 1, 2 and 3 and a compound A, B, C or D are particular combinations of the present invention.

The term "pharmaceutically-acceptable derivatives" in PCT/SE01/02657 includes

20 pharmaceutically-acceptable salts (e.g. acid addition salts).


In PCT/SE01/02657 pharmaceutically acceptable derivatives of compounds of formula I also include "protected" derivatives, and/or compounds that act as prodrugs, of compounds of formula I.

25 Compounds that may act as prodrugs of compounds of formula I that may be mentioned include compounds of formula Ia,




Ia

wherein R^{3a} represents a structural fragment of formula I(iii) or I(iv):



I(iii)



I(iv)

5 wherein R^5 represents OR^6 or $C(O)OR^7$;

R^6 represents H, C_{1-10} alkyl, C_{1-3} alkylaryl or C_{1-3} alkyloxyaryl (the alkyl parts of which latter two groups are optionally interrupted by one or more oxygen atoms, and the aryl parts of which latter two groups are optionally substituted by one or more substituents selected from halo, phenyl, methyl or methoxy, which latter three groups are also

10 optionally substituted by one or more halo substituents); R^7 represents C_{1-10} alkyl (which latter group is optionally interrupted by one or more oxygen atoms), or C_{1-3} alkylaryl or C_{1-3} alkyloxyaryl (the alkyl parts of which latter two groups are optionally interrupted by one or more oxygen atoms, and the aryl parts of which latter two groups are optionally substituted by one or more substituents selected from halo, phenyl, methyl or methoxy, which latter three groups are also optionally substituted by one or more halo substituents); and

15 R^a , R^1 , R^2 , Y, R^4 , X_1 , X_2 , X_3 and X_4 are as hereinbefore defined, and pharmaceutically-acceptable derivatives thereof.

20 The term "pharmaceutically-acceptable derivatives" of compounds of formula Ia includes pharmaceutically-acceptable salts (e.g. acid addition salts).

The wavy lines on the bonds in the fragments of formulae I(iii) and I(iv) signify the bond positions of the fragments.

25 In particular, compounds of the invention are potent inhibitors of thrombin either as such and/or (e.g. in the case of prodrugs), are metabolised following administration to form

potent inhibitors of thrombin, for example as may be demonstrated in the tests described below.

By "prodrug of a thrombin inhibitor", we include compounds that form a thrombin inhibitor, in an experimentally-detectable amount, and within a predetermined time (e.g. about 1 hour), following oral or parenteral administration (see, for example, Test E below) or, alternatively, following incubation in the presence of liver microsomes (see, for example, Test G below).

10 However, it is estimated that only 40% of patients with AF who should benefit from anticoagulant therapy do so, owing to the risks associated with existing treatments. This also includes patients whose anticoagulant therapy is in combination with cardioversion (electrical or chemical). In particular, of the currently-available oral anticoagulants, warfarin (a vitamin K antagonist) carries the risk of bleeding, and the need for frequent
15 laboratory control. Vitamin K antagonists also demonstrate a notable risk of interaction with other drugs and certain foods, e.g. those that are rich in Vitamin K, and their use requires monitoring of the patient's blood coagulation status. Medication containing acetylsalicylic acid (an antiplatelet agent) also carries the risk of bleeding. Blood
20 coagulation is the key process involved in both haemostasis (i.e. the prevention of blood loss from a damaged vessel) and thrombosis (i.e. the formation of a blood clot in a blood vessel, sometimes leading to vessel obstruction).

There remains a need for a combination of an antiarrhythmic drug and an anti-coagulant drug that has fewer side-effects than existing therapies and will encourage the use of such a
25 combination in a higher percentage of AF patients, thus reducing morbidity and mortality in this patient group.

None of the above-mentioned documents disclose or suggest the administration of a compound of claim 1 in PCT/SE01/02657 in conjunction with a compound as defined in
30 claim 1 of WO 01/28992. Surprisingly, the administration of just such a combination gives rise to unexpected, beneficial effects.

Disclosure of the Invention

According to a first aspect of the invention there is provided a combination product
5 comprising :

- (1) a compound of claim 1 in PCT/SE01/02657;
- and
- (2) a compound as defined in claim 1 of WO 01/28992 .


10 According to a second aspect of the invention there is provided a combination product
comprising :

- (1) a compound of claim 20 in PCT/SE01/02657;
- and
- (2) a compound of Claim 34 of WO 01/28992.

15


According to a third aspect of the invention there is provided a combination product
comprising :

- (1) a compound of claim 20 in PCT/SE01/02657;
- and
- 20 (2) (a) 4-({3-[7-(3,3-dimethyl-2-oxobutyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}amino)benzonitrile:




which compound is referred to hereinafter as Compound A, or a pharmaceutically-acceptable salt thereof; or

(b) *tert*-butyl 2-{7-[3-(4-cyanoanilino)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethylcarbamate:



in the form of the free base, which compound is referred to hereinafter as Compound B or a pharmaceutically-acceptable salt thereof; or


(c) *tert*-butyl 2-{7-[4-(4-cyanophenyl)butyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethylcarbamate:



in the form of the free base, which compound is referred to hereinafter as Compound C or a pharmaceutically-acceptable salt thereof; or

(d) *tert*-butyl 2-{7-[(2*S*)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate:

5



in the form of the free base, which compound is referred to hereinafter as Compound D or a pharmaceutically-acceptable salt thereof;

10 wherein each of components (1) and (2) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

The combination product according to the invention provides for the administration of a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657

15 in conjunction with (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), and may thus be presented either as separate formulations, wherein at least one of those formulations comprises a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 and at least one

20 comprises (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or

Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), or may be presented (i.e. formulated) as a combined preparation (i.e. presented as a single

formulation including a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof).

5

Thus, there is further provided:

(1) a pharmaceutical formulation including a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 or a pharmaceutically-acceptable derivative thereof, and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier (which formulation is hereinafter referred to as a "combined preparation"); and

15

(2) a kit of parts comprising components:

(a) a pharmaceutical formulation including a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 or a pharmaceutically-acceptable derivative thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; and (b) a pharmaceutical formulation including (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof) in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier, which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

25

According to a further aspect of the invention, there is provided a method of making a kit of parts as defined above, which method comprises bringing a component (a), as defined above, into association with a component (b), as defined above, thus rendering the two components suitable for administration in conjunction with each other.

30

By bringing the two components "into association with" each other, we include that components (a) and (b) of the kit of parts may be:

- 5 (i) provided as separate formulations (i.e. independently of one another), which are subsequently brought together for use in conjunction with each other in combination therapy; or
- (ii) packaged and presented together as separate components of a "combination pack" for use in conjunction with each other in combination therapy.

10 Thus, there is further provided a kit of parts comprising:

- (I) one of components (a) and (b) as defined herein; together with
- (II) instructions to use that component in conjunction with the other of the two components.

- 15 The kits of parts described herein may comprise more than one formulation including an appropriate quantity/dose of a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 or derivative thereof, and/or more than one formulation including an appropriate quantity/dose of (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or
- 20 (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), in order to provide for repeat dosing. If more than one formulation (comprising either active compound) is present, such formulations may be the same, or may be different in terms of the dose of a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative) or (1) a compound as defined in claim 1 of WO 01/28992
- 25 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), chemical composition and/or physical form.

A further aspect of the invention provides a method of treatment of a condition where anticoagulant therapy is indicated, which comprises administration of a pharmaceutical

30 formulation including a compound of claim 1 in PCT/SE01/02657 or a compound of claim

20 in PCT/SE01/02657 (or a pharmaceutically-acceptable derivative thereof), and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

5

A further aspect of the invention provides a method of treatment of a condition where anticoagulant therapy is indicated (by which we mean where anticoagulation is required), which comprises administration of:

(a) a pharmaceutical formulation including a compound of claim 1 in PCT/SE01/02657 or
10 a compound of claim 20 in PCT/SE01/02657 or a pharmaceutically-acceptable derivative thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; in conjunction with

(b) a pharmaceutical formulation including (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C
15 or D (or pharmaceutically-acceptable salts thereof), in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier,
to a patient suffering from, or susceptible to, such a condition.

For the avoidance of doubt, as used herein, the term "treatment" includes therapeutic
20 and/or prophylactic treatment.

With respect to the kits of parts as described herein, by "administration in conjunction with", we include that respective formulations comprising a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof)
25 and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), are administered, sequentially, separately and/or simultaneously, over the course of treatment of the relevant condition, which condition may be acute or chronic.

Thus, in respect of the combination product according to the invention, the term "administration in conjunction with" includes that the two components of the combination product (a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof)) are administered (optionally repeatedly), either (in the case of a combined preparation) together, or (in the case of a kit of parts) sufficiently closely in time, to enable a beneficial effect for the patient, that is greater, over the course of the treatment of the relevant condition, than if either a formulation comprising a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657, or a formulation comprising (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), are administered (optionally repeatedly) alone, in the absence of the other component, over the same course of treatment. Determination of whether a combination provides a greater beneficial effect in respect of, and over the course of treatment of, a particular condition, will depend upon the condition to be treated or prevented, but may be achieved routinely by the skilled person.

Further, in the context of a kit of parts according to the invention, the term "in conjunction with" includes that one or other of the two formulations may be administered (optionally repeatedly) prior to, after, and/or at the same time as, administration with the other component. When used in this context, the terms "administered simultaneously" and "administered at the same time as" include that individual doses of a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), are administered within 48 hours (e.g. 24 hours) of each other.

Suitable daily doses of the compounds of a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) in therapeutic treatment of humans are about 0.001-100 mg/kg body weight at peroral administration and 0.001-50 mg/kg body weight at parenteral administration.

5

Suitable doses of (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), in the therapeutic and/or prophylactic treatment of mammalian, especially human, patients may be determined routinely by the medical practitioner or other skilled person, and include the respective doses discussed in WO 01/28992 which is hereby incorporated by reference.

10

In the case of antiarrhythmic oxabispidines typical daily doses of (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), are in the range 10 to 2000 mg, e.g. 25, such as 30, to 1200 mg of free base (i.e., in the case of a salt, excluding any weight resulting from the presence of a counter ion), irrespective of the number of compositions (e.g. tablets) that are administered during the course of that day. Preferred daily doses are in the range 50 to 1000 mg, such as 100 to 500 mg, for example 150mg, 200mg, 250 mg, 300mg, 350mg, 400mg or 450mg. Typical doses in individual compositions of the invention (e.g. tablets) are thus in the range 15 to 500 mg, for example 40 to 400 mg eg for example 150mg, 200mg, 250 mg, 300mg, 350mg or 400mg.

15

20

Specifically claimed herein are specific fixed dose combinations where any dose stated for a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) is combined with any dose stated for the antiarrhythmic oxabispidine, including the doses stated as limits for the ranges described.

25

In any event, the physician, or the skilled person, will be able to determine the actual dosage which will be most suitable for an individual patient, which is likely to vary with

30

the condition that is to be treated, as well as the age, weight, sex and response of the particular patient to be treated. The above-mentioned dosages are exemplary of the average case; there can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

5

When separate formulations are administered, the sequence in which the formulations comprising a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof), and the antiarrhythmic oxabispidine (or derivative thereof), may be administered (i.e. whether, and at what point, sequential, 10 separate and/or simultaneous administration takes place) may be determined by the physician or skilled person. For example, the sequence may depend upon many factors that will be evident to the skilled person, such as whether, at any time during the course or period of treatment, one or other of the formulations cannot be administered to the patient for practical reasons (e.g. the patient is unconscious and thus unable to take an oral 15 formulation comprising either a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) or the antiarrhythmic oxabispidine).

20

The method described herein may have the advantage that, in the treatment of conditions where anticoagulant therapy is indicated, it may be more convenient for the physician and/or patient than, be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, produce fewer side effects than, or that it may have other useful pharmacological properties over, similar methods known in the prior art for the treatment of such conditions.

25

A compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) and derivatives thereof, may be administered for systemic delivery using appropriate means of administration that are known to the skilled person.

30

Thus, in accordance with the invention, a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) and derivatives thereof, may be administered orally, intravenously, subcutaneously, buccally, rectally, dermally, nasally, tracheally, bronchially, topically, by any other parenteral route, or *via* inhalation, in the form of a pharmaceutical preparation comprising the active ingredient in a pharmaceutically-acceptable dosage form. Depending on the disorder, and the patient, to be treated, as well as the route of administration, the compositions may be administered at varying doses.

Preferred modes of delivery are systemic. For a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof), preferred modes of administration are oral, parenteral, more preferably intravenous, and especially subcutaneous. Preferred modes of administration are oral.

In the therapeutic treatment of mammals, and especially humans, a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) may be administered alone, but will generally be administered as a pharmaceutical formulation in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier, which may be selected with due regard to the intended route of administration and standard pharmaceutical practice.

The preparation of suitable formulations may be achieved non-inventively by the skilled person using routine techniques.

The combinations of the present invention are useful in both the prophylaxis and the treatment of cardiac arrhythmias, in particular atrial and ventricular arrhythmias (such as atrial fibrillation (e.g. atrial flutter)) and NVAf.

The combinations of the invention are thus indicated in the treatment or prophylaxis of cardiac diseases, or in indications related to cardiac diseases, in which arrhythmias are

believed to play a major role, including ischemic heart disorders, sudden heart attack, myocardial infarction, heart failure, cardiac surgery and thromboembolic events.

The term "ischemic disorders" will be understood by those skilled in the art to include any
5 condition, the results of which include a restriction in blood flow in a part of the body. In
this context, the term will also be understood to include thrombosis and hypercoagulability
in blood and/or organs, tissues, etc.

The term "thrombosis" will be understood by those skilled in the art to include the
10 formation, development or presence of a thrombus in animals including man, and which
may result in embolism and/or ischemia. The term may thus include conditions such as
atrophic thrombosis, arterial thrombosis, cardiac thrombosis, coronary thrombosis,
creeping thrombosis, infective thrombosis, mesenteric thrombosis, placental thrombosis,
propagating thrombosis, traumatic thrombosis and venous thrombosis.

15 The term "hypercoagulability" includes any state in which the blood is more readily
coagulated than usual.

The term "NVAf" may be understood by those skilled in the art to mean grossly
20 disorganised atrial electrical activity, which is irregular in respect of both rate and rhythm,
leading to a hypercoagulable state and an increased risk of thrombosis originating from the
left heart chambers, and particularly the left atrium. The term may thus also be understood
to include AF (chronic, persistent, permanent and/or intermittent (paroxysmal)) in the
absence of heart valvular disease (mostly rheumatic heart valvular disease e.g. mitral
25 stenosis), or prosthesis, and to exclude patients with rheumatic mitral stenosis.

Particular disease states that may be mentioned include the prevention/treatment of
ischemic heart disease, myocardial infarction, systemic embolic events in e.g. the kidneys,
spleen etc, and, more particularly, of cerebral ischemia, including cerebral thrombosis,
30 cerebral embolism and/or cerebral ischemia associated with non-cerebral thrombosis or

embolism (in other words, the treatment/prophylaxis of thrombotic, or ischemic, stroke and of transient ischemic attack (TIA)) in patients with, or at risk of, NVAF. The skilled person will appreciate that patients with NVAF who are at risk of stroke include elderly patients generally (e.g. those with an age of greater than 75 years); patients with
5 complicating health factors, such as hypertension, left ventricular dysfunction (e.g. left ventricular ejection fraction (LVEF) of less than 40%), symptomatic congestive heart failure, diabetes mellitus (especially in those patients of 65 years of age or greater) and/or coronary heart or artery disease (especially in those patients of 65 years of age or greater); and/or patients with a history of stroke, TIA and/or systemic embolism, all of which factors
10 may predispose such patients to stroke and/or thromboembolic events.

According to a further aspect of the invention, there is provided a method of treatment of an arrhythmia which method comprises administration of a combination of the invention to a person suffering from, or susceptible to, such a condition.

15

According to a further aspect of the invention, there is provided a method of treatment of atrial fibrillation which method comprises administration of a combination of the invention to a person suffering from, or susceptible to, such a condition.

20 According to a further aspect of the invention, there is provided a method of treatment of atrial flutter which method comprises administration of a combination of the invention to a person suffering from, or susceptible to, such a condition.

25 For the avoidance of doubt, by "treatment" we include the therapeutic treatment, as well as the prophylaxis, of a condition.

It is expected that the combinations of the present invention may provide one or more of the following advantages. Synergy between the components in terms of:

- response rate
- 30 - patient survival rate

- time to disease progression
- dose/response effects leading to lower doses with same efficacy.

Alternatively, it is expected that the combinations of the present invention may provide one
5 or more of the following advantages:

- lower toxicity/reduced side effects with similar/improved efficacy;
- improved physical properties, e.g. storage stability, flow properties etc.;
- ease of formulation for example, reduced drug/drug incompatibility problems;
- reduced drug/ drug interaction problems on administration, for example possible changes
10 in metabolism of one drug caused by the effect of the other drug;
- improved patient compliance;
- improved quality of life;
- covenient dosing regimes;
- or
- 15 lack of diminishing effects of one drug caused by the presence of the other drug.

It is expected that the combination of the present invention will lead to a reduced incidence
of strokes in patients susceptible to strokes by the treatment and prevention of atrial
fibrillation.

20

Improved patient compliance may be demonstrated by methods known to those skilled in
the art, for example by supplying patients with blister packs containing the combination of
the present invention wherein the date and time of the removal of a drug from the blister
pack is recorded.

25

In a further aspect the present invention provides a process for the preparation of a
combination product as described earlier comprising formulating (1) a compound of claim
1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative
thereof) with a pharmaceutically acceptable diluent or carrier; and then formulating (1) a
30 compound as defined in claim 1 of a compound of claim 1 in PCT/SE01/02657 or a

compound of claim 20 in PCT/SE01/02657 (or derivative thereof) or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof) in a dose as previously described herein with a pharmaceutically acceptable diluent or carrier ; and then combining these formulations to provide a combination product as previously described herein.

Compounds of WO 01/28992 may be prepared as described therein.

Compounds of PCT/SE01/02657 may be prepared as described below and analogous methods thereof.

General Experimental Details

TLC was performed on silica gel. Chiral HPLC analysis was performed using a 46 mm X 250 mm Chiralcel OD column with a 5 cm guard column. The column temperature was maintained at 35°C. A flow rate of 1.0 mL/min was used. A Gilson 115 UV detector at 228 nm was used. The mobile phase consisted of hexanes, ethanol and trifluoroacetic acid and the appropriate ratios are listed for each compound. Typically, the product was dissolved in a minimal amount of ethanol and this was diluted with the mobile phase.

LC-MS/MS was performed using a HP-1100 instrument equipped with a CTC-PAL injector and a 5 µm, 4x100 mm ThermoQuest, Hypersil BDS-C18 column. An API-3000 (Sciex) MS detector was used. The flow rate was 1.2 mL/min and the mobile phase (gradient) consisted of 10-90% acetonitrile with 90-10% of 4 mM aq. ammonium acetate, both containing 0.2% formic acid.

¹H NMR spectra were recorded using tetramethylsilane as the internal standard. ¹³C NMR spectra were recorded using the listed deuterated solvents as the internal standard.

Example 1

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OcBu)

(i) 3-Chloro-5-methoxybenzaldehyde

3,5-Dichloroanisole (74.0 g, 419 mmol) in THF (200 mL) was added dropwise to magnesium metal (14.2 g, 585 mmol, pre-washed with 0.5 N HCl) in THF (100 mL) at 25°C.

After the addition, 1,2-dibromoethane (3.9 g, 20.8 mmol) was added dropwise. The resultant dark brown mixture was heated at reflux for 3 h. The mixture was cooled to 0°C, and *N,N*-dimethylformamide (60 mL) was added in one portion. The mixture was partitioned with diethyl ether (3 x 400 mL) and 6N HCl (500 mL). The combined organic extracts were washed with brine (300 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to give an oil. Flash chromatography (2x) on silica gel eluting with Hex:EtOAc (4:1) afforded the sub-title compound (38.9 g, 54%) as a yellow oil.

¹H NMR (300 MHz, CDCl₃) δ 9.90 (s, 1H), 7.53 (s, 1H), 7.38 (s, 1H), 7.15 (s, 1H), 3.87 (s, 3H).

(ii) 3-Chloro-5-hydroxybenzaldehyde

A solution of 3-chloro-5-methoxybenzaldehyde (22.8 g, 134 mmol; see step (i) above) in CH₂Cl₂ (250 mL) was cooled to 0°C. Boron tribromide (15.8 mL, 167 mmol) was added dropwise over 15 min. After stirring, the reaction mixture for 2 h, H₂O (50 mL) was added slowly. The solution was then extracted with Et₂O (2 x 100 mL). The organic layers were combined, dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (4:1) afforded the sub-title compound (5.2 g, 25%).

¹H NMR (300 MHz, CDCl₃) δ 9.85 (s, 1H), 7.35 (s, 1H), 7.20 (s, 1H), 7.10 (s, 1H), 3.68 (s, 1H)

(iii) 3-Chloro-5-difluoromethoxybenzaldehyde

A solution of 3-chloro-5-hydroxybenzaldehyde (7.5g, 48 mmol; see step (ii) above) in 2-propanol (250 mL) and 30% KOH (100 mL) was heated to reflux. While stirring, CHClF₂ was bubbled into the reaction mixture for 2 h. The reaction mixture was cooled, acidified with 1N HCl and extracted with EtOAc (2 x 100 mL). The organics were washed with brine (100 mL), dried (Na₂SO₄), filtered and

concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (4:1) afforded the sub-title compound (4.6 g, 46%).

¹H NMR (300 MHz, CDCl₃) δ 9.95 (s, 1H), 7.72 (s, 1H), 7.52 (s, 1H), 7.40 (s, 1H), 6.60 (t, *J*_{H-F} = 71.1 Hz, 1H)

(iv) Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OTMS)CN

A solution of 3-chloro-5-difluoromethoxybenzaldehyde (4.6 g, 22.3 mmol; see step (iii) above) in CH₂Cl₂ (200 mL) was cooled to 0°C. ZnI₂ (1.8 g, 5.6 mmol) and trimethylsilyl cyanide (2.8 g, 27.9 mmol) were added and the reaction mixture was allowed to warm to room temperature and stirred for 15 h. The mixture was partially concentrated *in vacuo* yielding the sub-title compound as a liquid, which was used directly in step (v) below without further purification or characterization.

(v) Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OH)C(NH)OEt

Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OTMS)CN (6.82 g, assume 22.3 mmol; see step (iv) above) was added dropwise to HCl/EtOH (500 mL). The reaction mixture was stirred 15 h, then partially concentrated *in vacuo* yielding the sub-title compound as a liquid, which was used in step (vi) without further purification or characterization.

(vi) Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OH)C(O)OEt

Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OH)C(NH)OEt (6.24 g, assume 22.3 mmol; see step (v) above) was dissolved in THF (250 mL), 0.5M H₂SO₄ (400 mL) was added and the reaction was stirred at 40°C for 65 h, cooled and then partially concentrated *in vacuo* to remove most of the THF. The reaction mixture was then extracted with Et₂O (3 x 100 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound as a solid, which was used in step (vii) without further purification or characterization.

(vii) Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OH)C(O)OH

A solution of Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OH)C(O)OEt (6.25 g, assume 22.3 mmol; see step (vi) above) in 2-propanol (175 mL) and 20% KOH (350 mL) was stirred at room

temperature 15 h. The reaction was then partially concentrated *in vacuo* to remove most of the 2-propanol. The remaining mixture was acidified with 1M H₂SO₄, extracted with Et₂O (3 x 100 mL), dried (Na₂SO₄) and concentrated *in vacuo* to give a solid. Flash chromatography on silica gel eluting with CHCl₃:MeOH:concentrated NH₄OH (6:3:1) afforded the ammonium salt of the sub-title compound. The ammonium salt was then dissolved in a mixture of EtOAc (75 mL) and H₂O (75 mL) and acidified with 2N HCl. The organic layer was separated and washed with brine (50 mL), dried (Na₂SO₄) and concentrated *in vacuo* to afford the sub-title compound (3.2 g, 57% from steps (iv) to (vii)).

¹H NMR (300 MHz, CD₃OD) δ 7.38 (s, 1H), 7.22 (s, 1H), 7.15 (s, 1H), 6.89 (t, *J*_{H-F} = 71.1 Hz, 1H), 5.16 (s, 1H)

(viii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (a) and Ph(3-Cl)(5-OCHF₂)-(S)CH(OAc)C(O)OH (b)

A mixture of Ph(3-Cl)(5-OCHF₂)-(R,S)CH(OH)C(O)OH (3.2 g, 12.7 mmol; see step (vii) above) and Lipase PS "Amano" (-2.0 g) in vinyl acetate (125 mL) and MTBE (125 mL) was heated at reflux for 48 h. The reaction mixture was cooled, filtered through Celite® and the filter cake washed with EtOAc. The filtrate was concentrated *in vacuo* and subjected to flash chromatography on silica gel eluting with CHCl₃:MeOH:concentrated NH₄OH (6:3:1) yielding the ammonium salts of the sub-title compounds (a) and (b). Compound (a) as a salt was dissolved in H₂O, acidified with 2N HCl and extracted with EtOAc. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (a) (1.2 g, 37%).

For sub-title compound (a)

¹H NMR (300 MHz, CD₃OD) δ 7.38 (s, 1H), 7.22 (s, 1H), 7.15 (s, 1H), 6.89 (t, *J*_{H-F} = 71.1 Hz, 1H), 5.17 (s, 1H)

(ix) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

To a solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (1.1 g, 4.4 mmol; see step (viii) above) and H-Aze-Pab(Teoc) (see international patent application WO 00/42059, 2.6 g, 5.7 mmol) in DMF (50 mL) at 0°C was added PyBOP (2.8 g, 5.3 mmol) and collidine (1.3 g,

10.6 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature for an additional 15 h. The reaction mixture was concentrated *in vacuo* and flash chromatographed on silica gel (3 x), eluting first with CHCl₃:EtOH (9:1), then with EtOAc:EtOH (20:1) and finally eluting with CH₂Cl₂:CH₃OH (95:5) to afford the sub-title
5 compound (1.0 g, 37%) as a white solid.

¹H NMR (300 MHz, CD₃OD, mixture of rotamers) δ 7.79-7.85 (d, *J* = 8.7 Hz, 2H), 7.15-7.48 (m, 5H), 6.89 and 6.91 (t, *J*_{H-F} = 71.1 Hz, 1H), 5.12 and 5.20 (s, 1H), 4.75-4.85 (m, 1H), 3.97-4.55 (m, 6H), 2.10-2.75 (m, 2H), 1.05-1.15 (m, 2H), 0.09 (s, 9H)
10 MS (m/z) 611 (M + 1)⁺

(x) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OcBu, Teoc)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.051 g, 0.08 mmol; see step (ix) above), was dissolved in 3 mL of acetonitrile and 0.062 g
15 (0.5 mmol) of *O*-cyclobutylhydroxylamine hydrochloride was added. The mixture was heated at 70°C for 4.5 h. The solvent was evaporated and the residue was partitioned between water and ethyl acetate. The aqueous phase was extracted two more times with ethyl acetate and the combined organic phase was washed with water, brine, dried (Na₂SO₄), filtered and evaporated. Yield: 0.054 g (95%).

20 ¹H-NMR (400 MHz; CD₃OD): δ 8.66-8.50 (m, 1H), 7.45 (d, 2H), 7.29 (m, 3H), 7.15 (m, 2H), 6.88 (t, 1H major rotamer), 6.85 (t, 1H minor rotamer), 5.18 (s, 1H major rotamer), 5.12 (s, 1H minor rotamer), 5.16 (m, 1H minor rotamer), 4.78 (m, 1H major rotamer), 4.70 (m, 1H), 4.50-4.30 (m, 3H), 4.19-3.93 (m, 3H), 2.71-2.44 (m, 1H), 2.34-2.11 (m, 5H),
25 1.78 (m, 1H), 1.62 (m, 1H), 0.96 (m, 2H), 0.01 (s, 9H)

(xi) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OcBu)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OcBu, Teoc) (0.054 g, 0.08 mmol; see step (x) above), was dissolved in 0.5 mL of CH₂Cl₂ and
30 3 mL of TFA. The reaction was allowed to proceed for 60 minutes. TFA was evaporated and the residue was purified using preparative HPLC. The fractions of interest were pooled and freeze-dried (2x), yielding 23 mg (54%) of the title compound.

MS (m/z) 536 (M - 1)⁻; 538 (M + 1)⁺

¹H-NMR (400 MHz; CD₃OD): δ 7.56 (d, 2H), 7.33 (m, 3H), 7.15 (m, 2H), 6.89 (t, 1H major rotamer), 6.86 (t, 1H minor rotamer), 5.18 (s, 1H major rotamer; and m, 1H minor rotamer), 5.11 (s, 1H minor rotamer), 4.77 (m, 1H major rotamer), 4.58 (m, 1H), 4.42 (m, 2H), 4.34 (m, 1H major rotamer), 4.15 (m, 1H major rotamer), 4.06 (m, 1H minor rotamer), 3.97 (m, 1H minor rotamer), 2.66 (m, 1H minor rotamer), 2.52 (m, 1H major rotamer), 2.33-2.25 (m, 3H), 2.01-2.20 (m, 2H), 1.75 (m, 1H), 1.59 (m, 1H)

¹³C-NMR (100 MHz; CD₃OD) (carbonyl and/or amidine carbons, rotamers) δ 172.4, 172.3, 171.9, 171.4, 152.3

Example 2

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH)

(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH, Teoc)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.148 g, 0.24 mmol; see Example 1(ix) above), was dissolved in 9 mL of acetonitrile and 0.101 g (1.45 mmol) of hydroxylamine hydrochloride was added. The mixture was heated at 70°C for 2.5 h, filtered through Celite® and evaporated. The crude product (0.145 g; 75% pure) was used directly in the next step without further purification.

(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH, Teoc) (0.145 g, 0.23 mmol; see step (i) above), was dissolved in 0.5 mL of CH₂Cl₂ and 9 mL of TFA. The reaction was allowed to proceed for 60 minutes. TFA was evaporated and the residue was purified using preparative HPLC. The fractions of interest were pooled and freeze-dried (2x), yielding 72 mg (yield over two steps 62%) of the title compound.

MS (m/z) 482 (M - 1)⁻; 484 (M + 1)⁺

¹H-NMR (400 MHz; CD₃OD): δ 7.58 (d, 2H), 7.33 (m, 3H), 7.15 (m, 2H), 6.89 (t, 1H major rotamer), 6.86 (t, 1H minor rotamer), 5.18 (s, 1H major rotamer; and m, 1H minor rotamer), 5.12 (s, 1H minor rotamer), 4.77 (m, 1H major rotamer), 4.42 (m, 2H), 4.34 (m, 1H major rotamer), 4.14 (m, 1H major rotamer), 4.06 (m, 1H minor rotamer), 3.95 (m, 1H minor rotamer), 2.66 (m, 1H minor rotamer), 2.50 (m, 1H major rotamer), 2.27 (m, 1H major rotamer), 2.14 (m, 1H minor rotamer)

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 172.4, 172.3, 172.0, 171.4 152.3, 152.1

10 **Example 3**

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.045 g, 0.074 mmol; see Example 1(ix) above), was dissolved in 3 mL of TFA and allowed to react for 1 h. TFA was evaporated and the residue was freeze dried from water/acetonitrile to yield 0.043 g (100%) of the sub-title compound as its TFA salt.

¹H-NMR (400 MHz; CD₃OD) rotamers: δ 7.8-7.75 (m, 2H), 7.55-7.5 (m, 2H), 7.35 (m, 1H, major rotamer), 7.31 (m, 1H, minor rotamer), 7.19 (m, 1H, major rotamer), 7.15 (m, 1H), 7.12 (m, 1H, minor rotamer), 6.89 (t, 1H, major rotamer), 6.87 (t, 1H, minor rotamer), 5.22 (m, 1H, minor rotamer), 5.20 (s, 1H, major rotamer), 5.13 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.6-4.4 (m, 2H), 4.37 (m, 1H, major rotamer), 4.19 (m, 1H, major rotamer), 4.07 (m, 1H, minor rotamer), 3.98 (m, 1H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.55 (m, 1H, major rotamer), 2.29 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer)

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 172.6, 172.5, 172.0, 171.7, 167.0

MS (m/z) 465 (M - 1)⁻, 467 (M + 1)⁺

Example 4

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(COOCpenty)

To a solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA (74 mg, 0.13 mmol; see Example 3 above) and cyclopentylchloroformate (44 mg, 0.30 mmol) in methylene chloride (5 mL) was added aq. NaOH (0.5 mL, 2M, 1 mmol). The mixture was stirred at room temperature and the reaction was monitored with HPLC. After 2.5 hours, water was added and the liquid phases were separated. The aqueous phase was extracted twice with methylene chloride. The combined organic phases were dried (MgSO₄) and purified on silica gel (first methylene chloride, then EtOAc). After removal of the solvents *in vacuo*, the solid residue was dissolved in water/acetonitrile and freeze-dried to afford the title compound as a white solid. Yield: 33mg (44%)

MS (m/z) 579 (M + 1)⁺

¹H NMR (400MHz; CD₃OD): δ 7.79(d, 2H), 7.43-7.30(m, 5H), 7.20-7.11(m, 2H), 6.90(t, 1H, major rotamer), 6.87(t, 1H, minor rotamer), 5.19(dd, 1H, minor rotamer), 5.18(s, 1H, major rotamer), 5.13(m, 1H), 5.11(s, 1H, minor rotamer), 4.78(dd, 1H, major rotamer), 4.45(m, 2H), 4.35(m, 1H, major rotamer), 4.16(s, 1H, major rotamer), 4.06(s, 1H, minor rotamer), 3.97(s, 1H, minor rotamer), 2.68(m, 1H, minor rotamer), 2.52(s, 1H, major rotamer), 2.28(s, 1H, major rotamer), 2.16(s, 1H, minor rotamer), 1.90(m, 2H), 1.77(m, 4H), 1.61(m, 2H)

¹³C NMR (carbonyl and/or amidine protons; 100 MHz): δ 173.6, 173.1, 172.6, 170.3, 165.6

Example 5

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Z)

The title compound was prepared according to the procedure described in Example 4 above starting from Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA (73 mg, 0.13 mmol; see Example 3 above) and benzylchloroformate (35 mg, 0.21 mmol). Additional purification by reverse-phase HPLC (0.1M ammonium acetate/MeCN 40/60) was necessary. The appropriate fractions were concentrated *in vacuo* and extracted with EtOAc. Yield: 24mg (32%).

MS (m/z) 602 (M + 1)⁺

¹H NMR (400MHz; CD₃OD): Δ 7.80(d, 2H), 7.43-7.25(m, 8H), 7.20-7.10(m, 2H), 6.90(t, 1H, major rotamer), 6.88(t, 1H, minor rotamer), 5.18(dd, 1H, minor rotamer), 5.18(s, 2H), 5.17(s, 1H, rotamer), 5.11(s, 1H, rotamer), 4.78(dd, 1H, major rotamer), 4.45(m, 2H), 4.34(m, 1H, major rotamer), 4.15(s, 1H, major rotamer), 4.06(s, 1H, minor rotamer),
5 3.97(s, 1H, minor rotamer), 2.66(m, 1H, minor rotamer), 2.51(s, 1H, major rotamer), 2.27(s, 1H, major rotamer), 2.15(s, 1H, minor rotamer)

¹³C NMR (carbonyl and/or amidine protons; 100MHz): Δ 173.6, 173.1, 172.6, 170.5, 164.9

Example 6

10 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab x TFA

(i) 2-Nitro-5-trifluoromethoxybenzoic acid

To a solution of 3-trifluoromethoxybenzoic acid (49.0 g, 0.24 mol) in sulfuric acid (500 mL) at less than 0°C (ice-MeOH bath) was added a solution of potassium nitrate (31.3 g,
15 0.31 mol) in sulfuric acid (200 mL) over 20 minutes. The resulting solution was stirred at 0°C for 2 hours, then warmed to room temperature and stirred for 18 hours. The reaction was poured into ice and the resulting acidic solution was extracted with EtOAc (5x). The combined organics were washed with H₂O (1x), brine (2x), H₂O (1x) and brine (1x), dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound (65.7 g)
20 as a solid contaminated with HOAc. The crude sub-title compound was dissolved in EtOAc and toluene and concentrated *in vacuo* to give a HOAc free solid (58.4 g, 97%) that was used in the next step without further purification.

25 ¹H NMR (300 MHz, CDCl₃): Δ 10.10 (br s, 1H), 8.02 (d, 1H, J = 8 Hz), 7.69 (d, 1H, J = 2 Hz), 7.54 (dd, 1H, J = 2 Hz, J = 8 Hz)

(ii) 2-Amino-5-trifluoromethoxybenzoic acid

To a solution of 2-nitro-5-trifluoromethoxybenzoic acid (56.8 g, 0.23 mol; see step (i) above) in EtOH (1000 mL) was added 10% Pd/C (5.7 g). The resulting solution was
30 flushed with H₂ for 5 h, filtered through Celite® and concentrated *in vacuo* to give the crude sub-title compound (49.7 g, 98%) as a solid that was used in the next step without further purification.

¹H NMR (300 MHz, CD₃OD): δ 7.66 (m, 1H), 7.17 (d, 1H, J = 8 Hz), 6.77 (d, 1H, J = 8 Hz)

5 (iii) 2-Amino-3-chloro-5-trifluoromethoxybenzoic acid

To a solution of 2-amino-5-trifluoromethoxybenzoic acid (49.0 g, 0.22 mol; see step (ii) above) in HOAc (1200 mL) was slowly added sulfuryl chloride (41.8 g, 0.31 mol). Gas evolution was observed. The resulting heterogeneous mixture was stirred at room temperature for 1 h. Additional HOAc (300 mL) was added to aid stirring, followed by
10 sulfuryl chloride in 5 mL portions until the starting material was consumed based on TLC analysis. The reaction was concentrated *in vacuo* to give solids that were flushed on a rotary evaporator with EtOAc (2x) followed by Et₂O (1x) to remove the HOAc. The resulting solids were further dried to give the HCl salt of the crude sub-title compound (60.5 g, 94%), which was used in the next step without further purification.

15

¹H NMR (300 MHz, CD₃OD): δ 7.72 (s, 1H), 7.44 (s, 1H), 7.22 (s, exchangeables)

(iv) 3-Chloro-5-trifluoromethoxybenzoic acid

To a solution of 2-amino-3-chloro-5-trifluoromethoxybenzoic acid (60.5 g, assume 0.22
20 mol; see step (iii) above) in 1,4-dioxane (1000 mL) was added 6N HCl (750 mL). Some organics oiled out of solution. The dioxane solution was cooled to less than 0°C (ice-MeOH bath). A solution of sodium nitrite (18.2 g, 0.26 mol) in H₂O (250 mL) was added over 15 minutes *via* an addition funnel. The resulting solution was stirred for 45 min. Hypophosphorous acid (221.5 mL of 50 wt% in H₂O, 291.2 g,
25 2.20 mol) was added slowly *via* an addition funnel. The solution was stirred at 0°C for 1.5 hours, then warmed to room temperature (gas evolution observed) and stirred for 18 hours. The crude solution was transferred to a separating funnel and extracted with Et₂O (4x). The combined organics were extracted with aqueous NaHCO₃ (3x). The basic aqueous layer was cautiously acidified with 6N HCl and extracted with CH₂Cl₂ (3x). The CH₂Cl₂
30 extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound (26.5 g, 46% from 3-trifluoromethoxybenzoic acid) as a solid that was used in the next step without further purification.

¹H NMR (300 MHz, CD₃OD): δ 7.98 (s, 1H), 7.83 (s, 1H), 7.58 (s, 1H)

(v) 3-Chloro-5-trifluoromethoxybenzyl alcohol

5 To a solution of 3-chloro-5-trifluoromethoxybenzoic acid (22.5 g, 93.5 mmol; see step (iv) above) in anhydrous THF (1200 mL) under a N₂ atmosphere at room temperature was added a solution of BH₃·THF complex (140 mL of 1M in THF; 140.3 mmol). The solution was refluxed for 2 h, cooled to room temperature and stirred for 18 hours, quenched cautiously with H₂O and concentrated *in vacuo* to remove most of
10 the THF. The residue was diluted with EtOAc and the organics were washed with brine (3x), dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound (21.2 g, 100%) as an oil that was used without further purification.

¹H NMR (300 MHz, CDCl₃): δ 7.33 (s, 1H), 7.17 (s, 1H), 7.14 (s, 1H), 4.72 (s, 2H), 2.05
15 (br s, 1H)

(vi) 3-Chloro-5-trifluoromethoxybenzaldehyde

A solution of DMSO (16.1 g, 205.9 mmol) in anhydrous CH₂Cl₂ (300 mL) was cooled to -78°C. Oxalyl chloride (13.1 g, 103.0 mmol) was added slowly *via* a syringe (gas evolution
20 was observed). The resulting solution was stirred at -78°C for 15 minutes. A solution of 3-chloro-5-trifluoromethoxybenzyl alcohol (21.2 g, 93.6 mmol; see step (v) above) in CH₂Cl₂ (200 mL) was added *via* an addition funnel over a period of 15 minutes. The cloudy solution was stirred at -78°C for 40 minutes and DIPEA (60.5 g, 468.0 mmol) was added *via* an addition funnel over 10 minutes. The resulting homogeneous solution was
25 stirred at -78°C for 1.5 hours, then warmed to room temperature and stirred 18 hours. The crude solution was concentrated *in vacuo*, the residue diluted with EtOAc and washed with H₂O (1x), 2N HCl (1x), brine (1x), aqueous NaHCO₃ (1x) and brine (1x). The organics were dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound (19.9 g, 95%) which was used in the next step without further purification.

30 ¹H NMR (300 MHz, CDCl₃): δ 10.00 (s, 1H), 7.83 (s, 1H), 7.66 (s, 1H), 7.51 (s, 1H)

(vii) Ph(3-Cl)(5-OCF₃)-(R,S)CH(OTMS)CN

To a solution of 3-chloro-5-trifluoromethoxybenzaldehyde (19.9 g, 88.6 mmol; see step (vi) above) in CH₂Cl₂ (600 mL) at 0°C was added ZnI₂ (1.4 g, 4.4 mmol) and trimethylsilyl cyanide (9.7 g, 97.5 mmol). After stirring at 0°C for 1.5 hours, and at room temperature for 2 hours, TLC analysis showed only the starting material. ZnI₂ was added portion-wise until the reaction proceeded (over 30.0 g of ZnI₂ was added in total). After stirring at room temperature for 18 h, the reaction was quenched with water and the organics were separated. The organics were dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound (27.7 g, 96%) as a liquid that was used without further purification.

¹H NMR (300 MHz, CDCl₃): δ 7.43 (s, 1H), 7.28 (s, 1H), 7.25 (s, 1H), 5.49 (s, 1H), 0.38 (s, 9H)

(viii) Ph(3-Cl)(5-OCF₃)-(R,S)CH(OH)C(O)OH

A suspension of Ph(3-Cl)(5-OCF₃)-(R,S)CH(OTMS)CN (27.7 g, 85.6 mmol; see step (vii) above) in concentrated HCl (300 mL) was refluxed for 3 hours. The resulting brown heterogeneous mixture was cooled to room temperature and extracted with Et₂O (2x). The initial organics were extracted with 2N NaOH (2x), then the basic layer was acidified with 2N HCl and extracted with Et₂O. The Et₂O was dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound (4.9 g, 21%). TLC analysis of the initial organics showed the sub-title compound was still present so the basic extraction/acidification was repeated using 6N NaOH to afford additional crude sub-title compound (2.8 g, 12%). TLC analysis of the initial organics showed the sub-title compound was still present so the organics were dried (Na₂SO₄) and concentrated *in vacuo* to give the sodium salt of the sub-title compound (18.3 g) as an oil. The salt was then redissolved in Et₂O and the organics acidified with 2N HCl and washed with brine. The resulting organics were dried (Na₂SO₄), treated with activated charcoal, filtered through Celite® and concentrated *in vacuo* to give the crude sub-title compound (14.3 g, 62%) as a solid that was used in the next step without further purification.

¹H NMR (300 MHz, CD₃OD): δ 7.53 (s, 1H), 7.38 (s, 1H), 7.29 (s, 1H), 5.23 (s, 1H)

(ix) Ph(3-Cl)(5-OCF₃)-(R,S)CH(OH)C(O)OH (a) and Ph(3-Cl)(5-OCF₃)-
(S)CH(OAc)C(O)OH (b)

A mixture of Ph(3-Cl)(5-OCF₃)-(R,S)CH(OH)C(O)OH (7.7 g, 28.5 mmol; see step (viii)
5 above) and Lipase PS "Amano" (3.8 g) in MTBE (100 mL) and vinyl acetate (50 mL) was
stirred at 60°C for 26 hours. The reaction was cooled and filtered through Celite® and the
filter cake washed with EtOAc. The combined organics were concentrated *in vacuo*. Flash
chromatography on silica gel eluting with CHCl₃:MeOH:concentrated NH₄OH (6:3:1)
afforded a mixture of the ammonium salts of sub-title compound (a) and sub-title
10 compound (b) (6.7 g) and a pure sample of the ammonium salt of sub-title compound (a)
(1.2 g) with less than 95% e.e. The respective fractions were dissolved in Et₂O and
washed with 2N HCl (1x) and brine (1x), dried (Na₂SO₄), filtered and concentrated to give
the corresponding carboxylic acids (6.7 g and 1.1 g respectively). These fractions were
then separately re-submitted to the resolution conditions and re-purified as necessary *via*
15 chromatography on silica gel eluting with CHCl₃:MeOH:concentrated NH₄OH (6:3:1 or
75:20:5 or 145:45:10) as needed. The purified sub-title compound (a) was acidified with
aqueous HCl or aqueous citric acid prior to further use. The ammonium salt of sub-title
compound (b) was used without characterization.

20 For sub-title compound (a)

¹H NMR (300 MHz, CD₃OD): δ 7.53 (s, 1H), 7.38 (s, 1H), 7.29 (s, 1H), 5.23 (s, 1H)

¹³C NMR (75 MHz, CD₃OD): δ 174.9, 150.9, 145.4, 136.3, 126.8, 122.0, 120.6, 118.9,
72.9

MS (m/z) 269 (M - 1)

25

(x) Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

A solution of Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)OH (0.73 g, 2.70 mmol; see step (ix)
above) in DMF (40 mL) under a nitrogen atmosphere was cooled to 0°C. To the solution
30 was added H-Aze-Pab(Teoc) (1.46 g, 3.24 mmol), collidine (0.82 g, 6.75 mmol) and PyBOP (1.83 g, 3.51 mmol). The solution
was stirred at 0°C for 2 h, warmed to room temperature and stirred 18 hours, quenched

with water and concentrated *in vacuo*. The residue was diluted with EtOAc and washed with H₂O (1x), aqueous NaHCO₃ (1x), aqueous citric acid (1x) and brine (1x), dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the crude sub-title compound. Flash chromatography on silica gel (2x) eluting with EtOAc:MeOH (30:1) then CH₂Cl₂:MeOH (93:7) afforded the sub-title compound (0.73 g, 43%) as a crushable foam.

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers): Δ 7.78-7.82 (d, 2H, *J* = 8 Hz), 7.25-7.54 (m, 5H), 5.25 and 5.16 (s, 1H), 5.22 and 4.79 (m, 1H), 3.92-4.58 (m, 6H), 2.20-2.76 (m, 2H), 1.04-1.13 (m, 2H), 0.08 (s, 9H)

MS (m/z) 629 (M + 1)⁺

(xi) Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab

Trifluoroacetic acid (1.0mL) was added to a stirred ice/water-cooled solution of Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (101 mg; 160 μ mol; see step (x) above), in methylene chloride (10 mL). The cooling bath was removed after 1 hour. After 1.5 hours at room temperature, acetonitrile (30 mL) was added and the solvents were carefully removed under reduced pressure. The residue was dissolved in water and freeze dried to afford 90 mg (92%) of the title compound as its TFA salt.

MS (m/z) 483 (M - 1)⁻; 485 (M + 1)⁺

¹H NMR (300 MHz; CD₃OD): (complex due to diastereomers/rotamers): Δ 7.70-7.80 (m, 2H), 7.45-7.58 (m, 3H), 7.24-7.38 (m, 2H), 5.26 (s, 1H), 5.17 (m, 1H, minor rotamer), 4.82 (m, 1H, major rotamer), 4.35-4.6 (m, 3H), 4.22 (m, 1H, major rotamer), 3.92-4.12 (m, 2H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.55 (m, 1H, major rotamer), 2.30 (m, 1H, major rotamer), 2.16 (m, 1H, minor rotamer)

¹³C NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers): Δ 173.7, 173.4, 173.0, 172.8, 168.1

Example 7

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OMe)

HATU (71 mg; 0.19 mmol) was added to a stirred ice/water-cooled solution of Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)OH (39 mg; 0.14 mmol; see Example 6(ix) above) in DMF (3mL).

After 30 minutes, a solution of H-Aze-Pab(OMe) x 2HCl (69 mg; 0.21 mmol; see international patent application WO 00/42059) and 2,4,6-collidine (0.080 mL; 0.58 mmol) in DMF (1.5 mL) was added. The reaction mixture was left overnight and the temperature was allowed to rise slowly to ambient. The solvents were removed *in vacuo* and the crude product was purified using reverse-phase HPLC (acetonitrile: 0.1M aq. ammonium acetate) to afford, after freeze drying the appropriate fractions, the title compound (61 mg, 97%) as a colourless solid.

MS (m/z) 513 (M - 1)⁻, 515 (M + 1)⁺

¹H NMR (500 MHz; CD₃OD): δ 7.97 (bt, 1H), 7.53 (d, 2H), 7.27 (t, 1H), 7.22 (d, 2H), 7.19 (t, 1H), 7.11 (t, 2H), 6.77 (s, 1H), 4.92 (s, 1H), 4.9 (bs, 3H), 4.81 (m, 2H), 4.40 (m, 2H), 4.09 (m, 1H) 3.87 (s, 3H), 2.58 (m, 1H), 2.37(m, 1H)

¹³C NMR (125 MHz; CD₃OD): (carbonyl and/or amidine carbons): δ 171.8, 169.9, 156.8

Example 8

Parallel Synthesis of Alkoxyamidines

This synthesis was performed in a 96-well Robbins block. To wells containing an appropriate amount of O-substituted hydroxylamine (specified below; all of which are commercially available or were prepared using well known literature procedures) was added a solution of Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (10 mg; 17 μ mol; see Example 6(x) above) in acetonitrile (1.0 mL). The block was sealed and the reaction mixture was rotated overnight in an oven at 60°C. After cooling and filtration, the solids were washed with acetonitrile (3 x 0.3 mL). The combined liquid fractions were concentrated in a vacuum centrifuge. The residue was partitioned between water (0.4 mL) and ethyl acetate (0.4 mL). After liquid-liquid extraction was finished, everything was filtered through a column of HydromatrixTM. After washing three times with ethyl acetate, the combined filtrates were concentrated in a vacuum centrifuge. Deprotection was performed by addition of methylene chloride (0.1 mL) and trifluoroacetic acid (0.3 mL). After stirring at room temperature for 3 hours, the solvents were removed *in vacuo*. The residue was partitioned between aqueous saturated sodium hydrogen carbonate (0.5 mL) and ethyl acetate (0.5 mL). After extraction, filtration through HydromatrixTM and

concentration (*vide infra*) the residue was dissolved in isopropanol/water (7/3) (1 mL). About 2% of this solution was removed and diluted with isopropanol/water (7/3) (1 mL) for LC-MS analysis. After removal of the solvents *in vacuo* the solid residue was transferred to a 96-well plate using acetonitrile and ethyl acetate to dissolve the compound.
5 The solvents were evaporated in a vacuum centrifuge to afford the following title compounds:

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂-3-(5-Me-isoxazole))

(from 3-[(aminooxy)methyl]-5-methylisoxazole x HCl (18 mg; 0.11 mmol)). Yield:
10 3.64mg (35%) (MS (m/z) 596 (M + 1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂-3-pyridine)

(from 3-[(aminooxy)methyl]pyridine x 2 HCl (19 mg; 96 μmol). Yield: 5.14 mg (50%)
(MS (m/z) 592 (M + 1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OiBu)

15 (from *O*-isobutyl hydroxylamine x HCl (17 mg; 140 μmol). Yield: 4.4 mg (45%). MS
(m/z) 557 (M + 1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OEt)

(from *O*-ethyl hydroxylamine x HCl (14 mg; 140 μmol). Yield: 4.04 mg (42%). MS (m/z)
529 (M+1)⁺);

20 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OBn)

(from *O*-benzylhydroxylamine x HCl (17 mg; 110 μmol). Yield: 3.22 mg (29%). MS
(m/z) 591 (M + 1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OcHexyl)

25 (from *O*-cyclohexyl hydroxylamine x HCl (15 mg; 99 μmol). Yield: 2.9 mg (26%). MS
(m/z) 583 (M + 1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OcBu)

(from *O*-cyclobutyl hydroxylamine x HCl (17 mg; 140 μmol). Yield: 3.3 mg (30%). MS
(m/z) 555 (M + 1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂CH₂OPh(3-CF₃))

30 (from *O*-[2-[3-(trifluoromethyl)phenoxy]ethyl]hydroxylamine x HCl (24 mg; 93 μmol).
Yield: 6.52 mg (46%). MS (m/z) 689 (M+1)⁺);

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OBn(4-Cl))

(from *O*-(4-chlorobenzyl)hydroxylamine x HCl (16 mg; 82 μ mol). Yield: 3.47 mg (29%).

MS (m/z) 625 ($M + 1$)⁺;

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OBn(3-MeO))

(from *O*-(3-methoxybenzyl)hydroxylamine x HCl (18 mg; 94 μ mol). Yield: 4.33 mg

5 (36%). MS (m/z) 621 ($M + 1$)⁺;

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OBn(2-Br))

(from *O*-(2-bromobenzyl)hydroxylamine x HCl (23 mg; 96 μ mol). Yield: 3.87 mg (30%).

MS (m/z) 671 ($M + 1$)⁺;

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OBn(4-Me))

10 (from *O*-(4-methylbenzyl)hydroxylamine x HCl (14 mg; 81 μ mol). Yield: 2.91 mg (25%).

MS (m/z) 605 ($M + 1$)⁺; and

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(O-4-heptyl)

(from *O*-(4-heptyl)hydroxylamine x HCl (15 mg; 89 μ mol). Yield: 17 mg (100%). MS
(m/z) 599 ($M + 1$)⁺.

15

Example 9

Ph(3-Cl)(5-OCHF₂)-(S)CH(CH₂OH)C(O)-Aze-Pab x HOAc

(i) 3-Chloro-5-methoxybenzoic acid

20 Magnesium turnings (Fluka purum for Grignard reactions) were pre-treated in the following way: The turnings were placed in a glass sintered funnel and 0.1 M of hydrochloric acid was poured onto them. The turnings were stirred with a glass rod for a few seconds and then the acid was washed away with 3 portions of water. Finally, the turnings were washed with 2 portions of acetone and bottled. Tetrahydrofuran (100 mL,
25 99.95%) was dried by adding RedAl (1 g, 70% wt. in toluene). Pre-treated magnesium turnings (5 g, 200 mmol) were placed in a round bottomed flask, and were flushed with nitrogen 3 times. Dichloroanisole (26 g, 146 mmol) was dissolved in THF (100 mL, RedAl-dried) and dibromoethane (1.8 g, 10 mmol) was added. The reaction mixture was flushed with nitrogen and then refluxed
30 for 2 hours. Heating was interrupted and dry ice (10 g) was added portionwise over 2 minutes. When all of the dry ice was dissolved, the reaction mixture was poured into ice containing hydrochloric acid

(400 mL, 2 M). Extractive work up (ether, 300 mL) gave 11.2 g, 60.2 mmol (yield: 41%) of the sub-title compound.

¹H-NMR (500 MHz; acetone-d₆): Δ 7.57 (m, 1H), 7.49 (m, 1H), 7.23 (m, 1H), 3.91 (s, 3H)

5

(ii) 3-Chloro-5-hydroxybenzoic acid

Alumina (1.65 g, 60 mmol) and iodine (21 g, 82 mmol) were refluxed in toluene (200 mL) for 2 hours. Then, 3-chloro-5-methoxybenzoic acid (11.2 g, 60.2 mmol; see step (i) above) dissolved in toluene (50 mL) was added, together with tetrabutylammonium iodide (1.5 g, 4 mmol), and the mixture was refluxed for another 10 2 hours. After cooling to ambient temperature, extractive work up gave 8.7 g, 50 mmol (yield: 83%) of the sub-title compound.

¹H-NMR (300 MHz; acetone-d₆): Δ 9.27 (s, 1H), 7.48 (m, 1H), 7.44 (m, 1H), 7.11 (m, 1H)

15

(iii) 3-Chloro-5-difluoromethoxybenzoic acid

3-Chloro-5-hydroxybenzoic acid (6.4 g, 37.2 mmol; see step (ii) above) dissolved in chloroform (200 mL) was transferred to a 500 mL three-necked round-bottomed flask fitted with a dry ice condenser and a gas inlet tube. Sodium hydroxide (100 mL, 5 M) was added and with vigorous stirring. Chlorodifluoromethane (Freon 22; 25 g, 290 mmol) was added portionwise through the gas inlet tube at ambient temperature. After 2 hours, the reaction was complete. Extractive work up gave 6.2 g, 28 mmol (yield: 75%) of the sub-title compound.

25 ¹H-NMR (500 MHz; acetone-d₆): Δ 7.87 (m, 1H), 7.74 (m, 1H), 7.54 (m, 1H), 7.19 (t, 1H, J_{H-F} 73 Hz)

30 (iv) 3-Chloro-5-difluoromethoxy-N-methoxy-N-methylbenzamide

3-Chloro-5-difluoromethoxybenzoic acid (1.8 g, 8 mmol; see step (iii) above) and oxalyl chloride (1.5 g, 11.8 mmol) were dissolved in methylene chloride (50 mL). DMF (2 drops)

was added and the reaction mixture was stirred at ambient temperature for 30 minutes. Then, *N,O*-dimethylhydroxylamine (1 g, 10.2 mmol) and triethylamine (3 g, 30 mmol) were added and after another 10 minutes stirring at ambient temperature, the reaction mixture was concentrated at reduced pressure. The residue was taken up in ether (100 mL) and water (50 mL). After separation, the organic phase was washed with brine, dried over sodium sulphate, filtered and concentrated. This residue was chromatographed on silica (hexane/ethyl acetate 2:1) which gave 2 g, 7.5 mmol (93%) of the sub-title compound.

¹H-NMR (400 MHz; CDCl₃): δ 7.54 (m, 1H), 7.37 (m, 1H), 7.27 (m, 1H), 6.53 (t, 1H, J_{H-F} 73 Hz)

(v) 3-Chloro-5-difluoromethoxyacetophenone

3-Chloro-5-difluoromethoxy-N-methoxy-N-methylbenzamide (2 g, 7.5 mmol; see step (iv) above) was dissolved in ether (100 mL) and cooled under nitrogen to -70°C. Methylolithium (7 mL, 11 mmol, 1.6 M in ether) was added dropwise with a syringe to the stirred reaction mixture over 1 minute. The dry ice bath was removed and the mixture was allowed to reach ambient temperature before the reaction was quenched with ammonium chloride solution (50 mL, 5% NH₄Cl in water). The organic phase was washed with brine, dried over sodium sulphate, filtered and concentrated at reduced pressure. The residue was chromatographed on silica (hexane:ethyl acetate 2:1) which gave 1.5 g, 6.8 mmol (yield: 90%) of the sub-title compound.

¹H-NMR (600 MHz; CDCl₃): δ 7.77 (m, 1H), 7.59 (m, 1H), 7.35 (m, 1H), 6.56 (t, 1H, J_{H-F} 73 Hz), 2.60 (s, 3H)

(vi) 3-Chloro-5-difluoromethoxyphenylacetic acid methyl ester

3-Chloro-5-difluoromethoxyacetophenone (1.5 g, 6.8 mmol; see step (v) above) was dissolved in methylene chloride (200 mL). Thallium(III) nitrate x 3MeOH on K-10 montmorillonite (6 g, 10 mmol (ca 0.6 mmol/g); see *J. Am. Chem. Soc.*, 98, 6750 (1976)) was added and the mixture was stirred at ambient temperature for 20 hours. The mixture was filtered and the filtrate was washed with sodium bicarbonate (100 mL, 0.5 M), dried over sodium sulphate, filtered and concentrated at reduced pressure. The residue was

chromatographed on silica (hexane/ethyl acetate 2:1) which gave 1 g, 4 mmol (yield: 56%) of the sub-title compound.

¹H-NMR (500 MHz; CDCl₃): δ 7.14 (m, 1H), 7.06 (m, 1H), 6.96 (m, 1H), 6.50 (t, 1H, J_{H-F} 73 Hz), 3.72 (s, 3H), 3.60 (s, 1H)

(vii) I-Formyl(3-chloro-5-difluoromethoxyphenyl)acetic acid methyl ester

3-Chloro-5-difluoromethoxyphenylacetic acid methyl ester (1 g, 4 mmol; see step (vi) above) and methyl formate (1 g, 16 mmol) were dissolved in ether (100 mL) and cooled in an ice-bath (ca. 2°C). Then, finely cut sodium (180 mg, 7.8 mmol) and methanol (1 mL) were added and the mixture was left in the ice-bath with stirring overnight. Water (100 mL) was added carefully and the phases were separated. The water containing phase was acidified with hydrochloric acid (2 M) to pH 1 and extracted with ether (2 x 100 mL). The extract was dried over sodium sulphate, filtered and concentrated at reduced pressure. The residue was chromatographed on silica (hexane:ethyl acetate (1:1)) which gave 400 mg, 1.4 mmol (yield: 36%) of the sub-title compound.

¹H-NMR (400 MHz): δ 12.10 (d, 1H), 7.32 (d, 1H), 7.11 (m, 1H), 7.07 (m, 1H), 6.94 (m, 1H), 6.51 (t, 1H, J_{F-H} 73), 3.83 (s, 3H)

(viii) 3-Chloro-5-difluoromethoxytropic acid

I-Formyl(3-chloro-5-difluoromethoxyphenyl)acetic acid methyl ester (400 mg, 1.4 mmol; see step (vii) above) was dissolved in THF:methanol (50 mL, 9:1). Sodium borohydride was added and the mixture was stirred at ambient temperature for 30 minutes. Water was added and the mixture was concentrated to produce an aqueous suspension, which was taken up in ethyl acetate and water. The phases were separated and the organic phase was washed with sodium chloride (15% in water), dried over sodium sulphate, filtered and concentrated at reduced pressure. The residue was dissolved in methanol (30 mL) and hydrolyzed with sodium hydroxide (1 mL, 10 M) at ambient temperature for 10 minutes. Extractive work up gave 180 mg, 0.68 mmol (yield: 48%) of the sub-title compound.

¹H-NMR (500 MHz; CDCl₃): Δ 7.18 (m, 1H), 7.10 (m, 1H), 7.00 (m, 1H), 6.50 (t, 1H, J_{F-H} 73), 4.11 (m, 1H), 3.90 (m, 1H), 3.84 (m, 1H)

(ix) Ph(3-Cl)(5-OCHF₂)-(S)CH(CH₂OH)C(O)-Aze-Pab x HOAc

- 5 3-Chloro-5-difluoromethoxytropic acid (180 mg, 0.7 mmol; see step (viii) above), H-Aze-Pab(Teoc) x HCl (450 mg, 1 mmol) and PyBOP (530 mg, 1 mmol) were dissolved in DMF (10 mL), whereafter DIPEA (550 mg, 3.9 mmol) was added. The mixture was stirred at ambient temperature for 1 h before it was diluted with brine (20 mL, 15% NaCl) and extracted with ethyl acetate (40 mL). The extract was dried over sodium sulphate, filtered and evaporated to dryness.
- 10 The residue was dissolved in methylene chloride (5 mL) and trifluoroacetic acid (5 mL) was added. After 1 h at ambient temperature, the mixture of diastereomers was evaporated to dryness and the residue was chromatographed on a reverse phase column (acetonitrile:water (30:70), buffer: ammonium acetate 0.1 M). Freeze drying gave 36 mg, 0.067 mmol (yield: 10.4%) of the title compound.

15

MS (ES) 481 (M + 1)⁺

¹H-NMR (400 MHz; CDCl₃): Δ 7.77 (d, 2H), 7.57 (d, 2H), 7.30 (m, 1H), 7.13 (m, 2H), 6.87 (t, 1H, J_{F-H} 73 Hz), 4.76 (m, 1H), 4.55 (s, 2H), 4.37 (m, 1H), 4.03 (m, 2H), 3.82 (m, 1H), 3.72 (m, 1H), 2.53 (m, 1H), 2.28 (m, 1H), 1.92 (s, 1.5H)

- 20 ¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons) Δ 172.3, 171.9, 167.2

Example 10

Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab x TFA

- 25 (i) 3-Chloro-5-trifluoromethoxybenzyl mesylate

To a solution of 3-chloro-5-trifluoromethoxybenzyl alcohol (6.1 g, 26.9 mmol; see Example 6(v) above) in CH₂Cl₂ (250 mL) at 0°C under a nitrogen atmosphere was added DIPEA (4.2 g, 32.3 mmol) and methanesulfonyl chloride (3.4 g, 29.6 mmol). The solution was stirred at 0°C for 1.5 hours and quenched with H₂O. The organics were separated and then washed with H₂O (1x), 1N HCl (1x), H₂O (1x) and aqueous NaHCO₃ (1x) and then dried (Na₂SO₄), filtered and concentrated to afford the sub-title compound (8.2 g, 99%) as an oil.

30

¹H NMR (300 MHz, CDCl₃): δ 7.37 (s, 1H), 7.28 (s, 1H), 7.18 (s, 1H), 5.23 (s, 2H), 3.07 (s, 3H)

5. (ii) 3-Chloro-5-trifluoromethoxybenzyl cyanide

To a solution of 3-chloro-5-trifluoromethoxybenzyl mesylate (8.2 g, 26.8 mmol; see step (i) above) in DMSO (50 mL) was added sodium cyanide (2.6 g, 53.6 mmol). The resulting heterogeneous solution was warmed to 50°C and sonicated for 1 hour. The reaction was cooled and partitioned between Et₂O and H₂O. The organics were washed with H₂O (2x) and brine (2x). The combined aqueous phases were extracted with Et₂O (1x). The combined organics were dried (Na₂SO₄), filtered and concentrated under a low heat and partial vacuum to afford the sub-title compound (6.3 g, 100%) as a reddish volatile oil which was used in the next step without further purification.

15 ¹H NMR (300 MHz, CDCl₃) δ 7.32 (s, 1H), 7.24 (s, 1H), 7.12 (s, 1H), 3.78 (s, 2H)

(iii) 3-Chloro-5-trifluoromethoxyphenylacetic acid

To a solution of 3-chloro-5-trifluoromethoxybenzyl cyanide (6.3 g, 26.7 mmol; see step (ii) above) in 2-propanol (100 mL) was added water (200 mL) and potassium hydroxide (7.5 g, 133.5 mmol). The solution was refluxed for 18 h, cooled to room temperature, and the 2-propanol was removed *in vacuo*. The aqueous phase was washed with CH₂Cl₂ (2x) and the washings discarded. The basic aqueous phase was acidified with 2N HCl and extracted with CH₂Cl₂ (3x). The CH₂Cl₂ extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (5.2 g, 76%) as an oil which was used in the next step without further purification.

¹H NMR (300 MHz, CDCl₃): δ 7.25 (s, 1H), 7.19 (s, 1H), 7.08 (s, 1H), 3.68 (s, 2H)

(iv) Ethyl 3-chloro-5-trifluoromethoxyphenylacetate

30 To a solution of 3-chloro-5-trifluoromethoxyphenylacetic acid (5.2 g, 20.4 mmol; see step (iii) above) in EtOH (600 mL) was added sulfuric acid (several drops). The solution was refluxed for 18 h, cooled to room temperature, neutralized with solid

NaHCO₃ and the EtOH removed *in vacuo*. The residue was diluted with EtOAc then washed with H₂O (1x), aqueous NaHCO₃ (1x) and brine (1x). The organics were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (5.5 g, 96%) as an oil which was used in the next step without further purification.

¹H NMR (300 MHz, CDCl₃) Δ 7.24 (s, 1H), 7.16 (s, 1H), 7.07 (s, 1H), 4.13-4.22 (q, *J* = 8 Hz, 2H), 3.63 (s, 2H), 1.24-1.32 (t, *J* = 8 Hz, 3H)

(v) Ph(3-Cl)(5-OCF₃)-(R,S)CH(CHO)C(O)OEt

To a solution of ethyl 3-chloro-5-trifluoromethoxyphenylacetate (4.5 g, 15.9 mmol; see step (iv) above) in anhydrous THF (400 mL) under a nitrogen atmosphere at less than 0°C (ice-MeOH bath) was added sodium ethoxide (4.5 g, 63.6 mmol). The cold solution was stirred for 40 minutes and ethyl formate (8.1 g, 111.3 mmol) was added. The solution was stirred at 0°C for 30 minutes, warmed to room temperature and stirred for 2 hours. Then, the THF was removed *in vacuo*. The residue was diluted with Et₂O and extracted with H₂O (1x) and 0.5M NaOH (3x). The aqueous extracts were acidified with 2N HCl and extracted with CH₂Cl₂ (3x). The combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the crude sub-title compound (3.9 g). Flash chromatography on silica gel eluting with Hex:EtOAc (4:1) afforded the sub-title compound (3.0 g, 61%) as an oil.

¹H NMR (300 MHz, CDCl₃, mixture of isomers): Δ 12.30 and 12.25 (s, 1H), 7.39 and 7.34 (s, 1H), 7.21 (s, 1H), 7.17 (s, 1H), 7.08 (s, 1H), 4.27-4.37 (q, *J* = 8 Hz, 2H), 1.28-1.38 (t, *J* = 8 Hz, 3H)

(vi) Ph(3-Cl)(5-OCF₃)-(R,S)CH(CH₂OH)C(O)OEt

To a solution of Ph(3-Cl)(5-OCF₃)-(R,S)CH(CHO)C(O)OEt (3.0 g, 9.66 mmol; see step (v) above) in MeOH (200 mL) at -10°C (ice-MeOH bath) was added sodium borohydride (0.7 g, 19.32 mmol) portion-wise over 5 min. The solution was stirred at -10°C for 45 minutes and additional sodium borohydride (0.4 g) was added. After another 15 minutes, the reaction was quenched with aqueous ammonium chloride, made weakly acidic with 2N HCl and the MeOH was removed *in vacuo*. The residue was diluted with EtOAc and washed with H₂O (1x), aqueous NaHCO₃ (1x) and brine (1x). The

organics were dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the crude sub-title compound. Flash chromatography on silica gel eluting with Hex:EtOAc (5:1) afforded the sub-title compound (2.0 g, 66%) as an oil.

5 ^1H NMR (300 MHz, CDCl_3): δ 7.26 (s, 1H), 7.19 (s, 1H), 7.07 (s, 1H), 4.16-4.28 (m, 2H), 4.04-4.15 (m, 1H), 3.76-3.94 (m, 2H), 2.33 (t, J = 6 Hz, 1H), 1.18-1.30 (t, J = 8 Hz, 3H)

(vii) $\text{Ph}(3\text{-Cl})(5\text{-OCF}_3)\text{-(R,S)CH}(\text{CH}_2\text{OH})\text{C(O)OH}$

To a solution of $\text{Ph}(3\text{-Cl})(5\text{-OCF}_3)\text{-(R,S)CH}(\text{CH}_2\text{OH})\text{C(O)OEt}$ (2.0 g, 6.24 mmol; see step (vi) above) in THF (50 mL) and H_2O (25 mL) was added lithium hydroxide monohydrate (0.5 g, 12.48 mmol). The solution was stirred at room temperature for 1 hour and the THF was removed *in vacuo*. The residue was diluted with H_2O then washed with CHCl_3 (2x) and the washings discarded. The basic aqueous layer was acidified with 2N HCl and extracted with CHCl_3 (4x). The CHCl_3 extracts were dried 15 (Na_2SO_4), filtered and concentrated *in vacuo* to afford the crude sub-title compound (1.5 g) as an oil. Flash chromatography on silica gel eluting with CHCl_3 :MeOH:concentrated NH_4OH (gradient from 7.0:2.5:0.5 to 6:3:1) afforded the ammonium salt of the sub-title compound (1.1 g). The ammonium salt was partitioned between 1N HCl and CHCl_3 . The organics were dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title 20 compound (also known as 3-chloro-5-trifluoromethoxytropic acid) as an oil (1.1 g, 62%).

^1H NMR (300 MHz, CD_3OD): δ 7.41 (s, 1H), 7.27 (s, 1H), 7.24 (s, 1H), 4.03 (m, 1H), 3.75-3.87 (m, 2H)

25 (viii) $\text{Ph}(3\text{-Cl})(5\text{-OCF}_3)\text{-(S)CH}(\text{CH}_2\text{OH})\text{C(O)-Aze-Pab(Teoc)}$ (a) and $\text{Ph}(3\text{-Cl})(5\text{-OCF}_3)\text{-(R)CH}(\text{CH}_2\text{OH})\text{C(O)-Aze-Pab(Teoc)}$ (b)

To a solution of $\text{Ph}(3\text{-Cl})(5\text{-OCF}_3)\text{-(R,S)CH}(\text{CH}_2\text{OH})\text{C(O)OH}$ (0.65 g, 2.28 mmol; see step (vii) above) in DMF at less than 0°C (ice-MeOH bath) was added H-Aze-Pab(Teoc) (0.90 g, 2.39 mmol), collidine (0.71 g, 5.70 mmol) and PyBOP (1.31 g, 2.51 mmol). The resulting solution was stirred at less than 0°C for 1 h, warmed to room 30 temperature and stirred for 1 hour. The DMF was then removed *in vacuo*. The residue was diluted with EtOAc and washed with dilute aqueous HCl (1x), brine (1x), aqueous

NaHCO₃ (1x) and brine (1x). The organics were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the crude sub-title compound (2.1 g) as a mixture of diastereomers. Flash chromatography (3x) on silica gel eluting first with EtOAc:MeOH (95:5) then with CH₂Cl₂:MeOH (97:3) and last with CH₂Cl₂:MeOH (95:5) afforded the sub-title compounds
5 diastereomer (a) (0.51 g, 35%) and diastereomer (b) (0.45 g, 31%) as crushable foams.

For sub-title compound diastereomer (a)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) Δ 7.79-7.85 (d, *J* = 8 Hz, 2H), 7.22-7.49 (m, 5H), 5.17-4.77 (m, 1H), 4.53-4.18 (m, 4H), 3.58-4.11 (m, 5H), 2.47-2.73 (m,
10 1H), 2.11-2.34 (m, 1H), 1.08-1.12 (m, 2H), 0.07 (s, 9H)

MS (*m/z*) 643 (*M* + 1)⁺

(ix) Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab x TFA

15 Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(Teoc), (78 mg, 0.121 mmol; see step (viii) above - diastereomer (a)), was dissolved in 5 mL of trifluoroacetic acid. After 10 minutes, the reaction was over and the solvent was evaporated. The residue was freeze dried from water and acetonitrile to give the desired product. Yield: 70 mg (94%).

20 MS (*m/z*) 483 (*M* - 1)⁻; 485 (*M* + 1)⁺

¹H-NMR(400 MHz; D₂O) rotamers 1:1: δ 8.83 (bt, 1H), 7.79 (d, 1H), 7.72 (d, 1H), 7.54 (d, 1H), 7.43 (d, 2H), 7.35 (m, 1H, rotamer), 7.28 (m, 1H, rotamer), 7.20 (m, 1H, rotamer), 7.05 (m, 1H, rotamer), 5.22 (m, 1H, rotamer), 4.83 (m, 1H, rotamer), 4.57 (m, 2H, rotamer), 4.38 (m, 2H, rotamer), 4.3-3.7 (m, 5H), 2.77 (m, 1H, rotamer), 2.55 (m, 1H, rotamer), 2.27 (m, 1H)

25 ¹³C-NMR (100 MHz; D₂O): (carbonyl and/or amidine carbons, rotamers) δ 172.9, 172.2, 172.0, 171.8, 166.9

Example 11

30 Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(OMe)

(i) Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(OMe, Teoc)

Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(Teoc) (100 mg, 0.155 mmol; see Example 10(viii) above), was dissolved in 12 mL of tetrahydrofuran. *O*-Methylhydroxylamine hydrochloride (44 mg, 0.53 mmol), was added and the reaction was heated at 50°C overnight. The reaction mixture was evaporated and the residue purified by preparative HPLC (CH₃CN/0.1 M NH₄OAc (70/30)). The pertinent fractions were evaporated and the residue dissolved in a small amount of acetonitrile and water and freeze dried. The freeze drying was repeated once. Yield: 80 mg (76%) of pure material.

¹H-NMR(400 MHz; CD₃OD) rotamers: δ 7.5-7.4 (m, 3H), 7.35-7.2 (m, 4H), 5.15 (m, 1H, minor rotamer), 4.74 (m, 1H, major rotamer), 4.5-4.25 (m, 3H), 4.2-3.95 (m, 4H), 3.91 (b, 3H), 3.9-3.6 (m, 2H), 2.63 (m, 1H, minor rotamer), 2.50 (m, 1H, major rotamer), 2.3-2.1 (m, 1H), 0.95 (m, 2H), 0.02 (s, 9H, major rotamer), 0.01 (s, 9H, minor rotamer)

(ii) Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(OMe)

Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(OMe, Teoc), (80 mg, 0.12 mmol; see step (i) above), was dissolved in 1 mL of methylene chloride and cooled in an ice bath. Trifluoroacetic acid, 3 mL, was added and the reaction flask was kept in the ice bath for two hours. The mixture was evaporated and dissolved in ethyl acetate and washed three times with NaHCO₃ (aq) then with water and brine. The organic phase was dried (Na₂SO₄), filtered and evaporated. The residue was freeze dried from a small amount of acetonitrile and water. Yield: 60 mg (95%) of pure title product.

MS (m/z) 528 (M - 1)⁻; 531 (M + 1)⁺

¹H-NMR(500 MHz; CD₃OD) rotamers: δ 7.65-7.55 (m, 3H, rotamers), 7.45 (m, 1H, major rotamer), 7.4-7.2 (m, 4H), 5.15 (m, 1H, minor rotamer), 4.74 (m, 1H, major rotamer), 4.5-4.3 (m, 3H), 4.05-3.95 (m, 2H), 3.85 (m, 1H, major rotamer), 3.82 (s, 3H, major rotamer), 3.81 (s, 3H, minor rotamer), 3.73 (m, 1H, major rotamer), 3.67 (m, 1H, minor rotamer), 3.62 (m, 1H, minor rotamer), 2.63 (m, 1H, minor rotamer), 2.50 (m, 1H, major rotamer), 2.24 (m, 1H, major rotamer), 2.16 (m, 1H, minor rotamer)

¹³C-NMR (125 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 174.0, 173.2, 172.7, 172.6, 155.1

Example 12Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)5 (i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe, Teoc)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.40 g, 0.65 mmol; see Example 1(ix) above), was dissolved in 20 mL of acetonitrile and 0.50 g (6.0 mmol) of O-methyl hydroxylamine hydrochloride was added. The mixture was heated at 70°C for 2 h. The solvent was evaporated and the residue was partitioned between water and ethyl acetate.
10 The aqueous phase was extracted twice more with ethyl acetate and the combined organic phase was washed with water, brine, dried (Na₂SO₄), filtered and evaporated. Yield: 0.41 g (91%).

¹H-NMR (400 MHz; CDCl₃) : δ 7.83 (bt, 1H), 7.57 (bs, 1H), 7.47 (d, 2H), 7.30 (d, 2H),
15 7.20 (m, 1H), 7.14 (m, 1H), 7.01 (m, 1H), 6.53 (t, 1H), 4.89 (s, 1H), 4.87 (m, 1H), 4.47 (m, 2H), 4.4-4.2 (b, 1H), 4.17-4.1 (m, 3H), 3.95 (s, 3H), 3.67 (m, 1H), 2.68 (m, 1H), 2.42 (m, 1H) 0.97 (m, 2H), 0.01 (s, 9H).

(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)

20 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe, Teoc) (0.40 g, 0.62 mmol; see step (i) above), was dissolved in 5 mL of TFA and allowed to react for 30 min. TFA was evaporated and the residue was partitioned between ethyl acetate and NaHCO₃ (aq.). The aqueous phase was extracted twice more with ethyl acetate and the combined organic phase was washed with water, brine, dried (Na₂SO₄), filtered and evaporated. The product
25 was freeze dried from water/acetonitrile. No purification was necessary. Yield: 0.28 g (85%).

¹H-NMR (600 MHz; CDCl₃) : δ 7.89 (bt, 1H), 7.57 (d, 2H), 7.28 (d, 2H), 7.18 (m, 1H),
7.13 (m, 1H), 6.99 (m, 1H), 6.51 (t, 1H), 4.88 (s, 1H), 4.87 (m, 1H), 4.80 (bs, 2H), 4.48 (dd,
30 1H), 4.43 (dd, 1H), 4.10 (m, 1H), 3.89 (s, 3H), 3.68 (m, 1H), 2.68 (m, 1H), 2.40 (m, 1H).
¹³C-NMR (125 MHz; CDCl₃): (carbonyl and/or amidine carbons, rotamers) δ 172.9, 170.8, 152.7, 152.6

MS (m/z) 495 (M - 1)⁻, 497 (M + 1)⁺

Example 13

Ph(3-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab x HOAc

5 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe) (13 mg, 0.026 mmol; see Example 12 above) was dissolved in abs. ethanol (5 mL) and 30 mg of 10% Pd/C was added. Finally acetic acid (5 µL) was added and the mixture was hydrogenated at atmospheric pressure for 20h. The mixture was filtered through Celite®, evaporated, and purified by reversed phase HPLC (0.1 M aq. ammonium acetate/MeCN). The appropriate fractions
10 were freeze-dried to afford the title compound as a white solid: 8.5 mg (66%).

¹H-NMR(400 MHz; CD₃OD) rotamers: δ 7.73-7.78 (m, 2H), 7.55 (d, 2H), 7.19-7.43 (m, 3H), 7.06-7.13 (m, 1H), 6.83 (t, 1H, J_{HF} = 74Hz, major rotamer), 6.81 (t, 1H, major rotamer), 5.20 (s, 1H, major rotamer), 5.19 (m, 1H, minor rotamer), 5.15 (s, 1H, minor rotamer), 4.78 (m, 1H, major rotamer), 4.4-4.6 (several peaks, 2H), 4.35 (m, 1H, major rotamer), 4.08 (m, 1H), 3.99 (m, 1H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.52 (m, 1H, major rotamer), 2.30 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer), 1.89 (s, 3H).
15

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 173.7, 172.9, 168.3.
20

MS (m/z) 433 (M+1)⁺; 431 (M-1)⁻

Example 14

Ph(3-OCF₃)-(R)CH(OH)C(O)-Aze-Pab x TFA

25 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab x TFA (34 mg, 0.057 mmol, from Example 6) was dissolved in 5 mL of ethanol and 20 mg of 10% Pd/C was added. The mixture was hydrogenated at atmospheric pressure overnight. The mixture was filtered through Celite®, evaporated, and freeze dried from water/acetonitrile.

30 ¹H-NMR(400 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.55 (m, 2H), 7.5-7.2 (m, 4H), 5.24 (s, 1H, major rotamer), 5.23 (m, 1H, minor rotamer), 5.18 (s, 1H, minor rotamer),

4.77 (m, 1H, major rotamer), 4.6-4.45 (m, 2H), 4.36 (m, 1H, major rotamer), 4.08 (m, 1H), 3.99 (m, 1H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.52 (m, 1H, major rotamer), 2.30 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 174.1, 173.9, 173.5, 172.9, 168.2.

¹⁹F NMR (282 MHz; CD₃OD): -59.8 and -59.9 (3F, minor and major rotamer respectively), -77.4 (3F) indicates that the salt is TFA.

MS (m/z) 451.3 (M+1)⁺

Example 15

Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab x TFA

(i) 3-Chloro-5-trifluoroethoxybenzaldehyde

To a magnetically stirred solution of 3-chloro-5-hydroxybenzaldehyde (2.0 g, 12.8 mmol; see Example 1(ii) above) and potassium carbonate (2.3 g, 16.6 mmol) in DMF (35 mL) under nitrogen was added 2,2,2-trifluoroethyl *p*-toluenesulfonate (4.2 g, 16.6 mmol) at room temperature. The mixture was heated to 110°C for 7 h and then stirred overnight at room temperature. The reaction was cooled to 0°C, poured into ice-cold 2 N HCl (100 mL) and extracted with EtOAc (2 x 75 mL). The combined organic extracts were washed with 0.5 N HCl (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The brown oil was chromatographed on silica gel eluting with Hex:EtOAc (6:1) to afford the sub-title compound (1.9 g, 61%) as a yellow oil.

^1H NMR (300 MHz, CDCl_3) Δ 9.44 (s, 1H), 7.56 (s, 1H), 7.33 (s, 1H), 7.28 (s, 1H), 4.42 (q, $J = 8$ Hz, 2H)

(ii) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CF}_3)\text{-(R,S)CH(OTMS)CN}$

- 5 To a solution of 3-chloro-5-trifluoroethoxybenzaldehyde (5.2 g, 21.7 mmol; see step (i) above) and zinc iodide (1.7 g, 5.4 mmol) in CH_2Cl_2 (200 mL) under nitrogen was added trimethylsilyl cyanide (4.3 g, 43.3 mmol) dropwise *via* syringe at 0°C . The mixture was stirred at 0°C for 3 h then diluted with H_2O (150 mL). The organic layer was separated, dried (Na_2SO_4), filtered, and concentrated *in vacuo* to afford the sub-title compound (6.9 g, 10 95%) as a yellow oil which was used without further purification.

^1H NMR (300 MHz, CDCl_3) Δ 7.27 (s, 1H), 6.98 (s, 2H), 5.44 (s, 1H), 4.38 (q, $J = 8$ Hz, 2H), 0.30 (s, 9 H)

15 (iii) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CF}_3)\text{-(R,S)CH(OH)C(O)OH}$

- Concentrated hydrochloric acid (170 mL) was added to $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CF}_3)\text{-(R,S)CH(OTMS)CN}$ (6.9 g, 20.4 mmol; see step (ii) above) and stirred at 100°C for 1 h. After cooling to room temperature, the reaction was further cooled to 0°C and basified slowly with 3 N NaOH (300 mL). This mixture was washed with Et_2O (2 x 100 mL) and 20 the aqueous layer was acidified with 2 N HCl (50 mL). The aqueous layer was then extracted with EtOAc (2 x 100 mL), dried (Na_2SO_4), filtered, and concentrated *in vacuo* to afford the sub-title compound (5.3 g, 92%) as a pale yellow oil which was used without further purification.

- 25 ^1H NMR (300 MHz, CD_3OD) Δ 7.18 (s, 1H), 7.07 (s, 1H), 7.02 (s, 1H), 5.13 (s, 1H), 4.58 (q, $J = 8$ Hz, 2H)

(iv) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CF}_3)\text{-(R)CH(OH)C(O)OH}$ (a) and $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CF}_3)\text{-(S)CH(OAc)C(O)OH}$ (b)

- 30 A solution of $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CF}_3)\text{-(R,S)CH(OH)C(O)OH}$ (7.06 g, 24.8 mmol; see step (iii) above) and Lipase PS "Amano" (4.30 g) in vinyl acetate (250 mL) and MTBE (250 mL) was heated at 70°C under nitrogen for 40 h. The reaction was cooled to room

temperature, the enzyme was removed by filtration washing with EtOAc, and the filtrate concentrated *in vacuo*. Chromatography on silica gel eluting with $\text{CHCl}_3:\text{MeOH}:\text{Et}_3\text{N}$ (92:6:2) afforded the triethylamine salt of the sub-title compound (a) (3.02 g) as a yellow oil. The salt of sub-title compound (a) was dissolved in H_2O (150 mL), acidified with 2 N HCl and extracted with EtOAc (2 x 75 mL). The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated *in vacuo* to yield the sub-title compound (a) (2.18 g) as an off-white solid. In addition, the triethylamine salt of sub-title compound (b) (4.73 g) was obtained from the column chromatography mentioned above.

10 Data for sub-title compound (a):

mp: 98-103°C

¹H NMR (300 MHz, CD₃OD) Δ 7.18 (s, 1H), 7.07 (s, 1H), 7.02 (s, 1H), 5.13 (s, 1H), 4.58 (q, *J* = 8 Hz, 2H).

15 ¹³C NMR (75 MHz, CD₃OD) Δ 175.4, 159.6, 144.6, 136.2, 125.0 (q, *J* = 277 Hz), 121.8, 115.9, 113.1, 73.3, 67.0 (q, *J* = 35 Hz)

HPLC Analysis: 98.6%, >99% ee, Chiralcel OD Column (97:3:0.5 Hex:EtOH:TFA mobile phase)

$[\alpha]_D^{25} = -81.5^\circ$ (c = 1.0, MeOH)

APCI-MS: (M - 1) = 283 m/z

20

(v) Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

To a solution of Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)OH (0.50 g, 1.8 mmol; see step (iv) above (compound (a))) in DMF (20 mL) under nitrogen was added H-Aze-Pab(Teoc) x HCl (1.03 g, 2.3 mmol), PyBOP (1.01 g, 1.9 mmol), and DIPEA (0.57 g, 4.4 mmol) at 0°C. The reaction was stirred at 0°C for 2 h and then at room temperature for 20 h. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first with $\text{CHCl}_3:\text{EtOH}$ (10:1) and then with EtOAc:EtOH (10:1) to afford the sub-title compound (0.55 g, 48%) as a crushable white foam.

30 mp: 90-95°C

$R_f = 0.42$ (10:1 $\text{CHCl}_3:\text{EtOH}$)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.78-7.81 (m, 2H), 7.38-7.41 (m, 2H), 7.12-7.16 (m, 1H), 7.00-7.06 (m, 2H), 5.09-5.22 and 4.75-4.79 (m, 2H), 3.94-4.61 (m, 8H), 2.09-2.75 (m, 2H), 1.04-1.11 (m, 2H), 0.70 (s, 9H)

APCI-MS: (M + 1) = 643 m/z

5

(vi) Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.066 g, 0.103 mmol; see step (v) above), was dissolved in 3 mL of TFA and allowed to react for 30 min. TFA was evaporated and the residue was freeze dried from water/acetonitrile to yield 0.060 g (94%)

10 of the title compound as its TFA salt.

¹H-NMR (400 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.6-7.5 (m, 2H), 7.2-7.0 (m, 3H), 5.21 (m, 1H, minor rotamer), 5.17 (s, 1H, major rotamer), 5.11 (s, 1H, minor rotamer), 4.81 (m, 1H, major rotamer), 4.6-4.4 (m, 4H), 4.37 (m, 1H, major rotamer), 4.16 (m, 1H, major rotamer), 4.06 (m, 1H, minor rotamer), 3.99 (m, 1H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.54 (m, 1H, major rotamer), 2.29 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer)

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 172.2, 171.8, 171.7, 167.0.

20 MS (m/z) 499.3 (M+1)⁺

Example 16

Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab(OMe)

To a solution of Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)OH (0.48 g, 1.7 mmol; see Example 15(iv) above (compound (a)) in DMF (20 mL) under nitrogen was added H-Aze-Pab(OMe) x 2HCl (0.74 g, 2.2 mmol), PyBOP (0.97 g, 1.9 mmol), and DIPEA (0.55 g, 4.2 mmol) at 0°C. The reaction was stirred at 0°C for 2 h and then at room temperature for 20 h. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first with CHCl₃:EtOH (10:1) and second with EtOAc:EtOH (10:1) to afford the title compound (0.62 g, 69%) as a crushable white foam.

30

mp: 75-80°C

R_f = 0.43 (10:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.57-7.60 (m, 2H), 7.32-7.36 (m, 2H), 7.13-7.17 (m, 1H), 7.00-7.06 (m, 2H), 5.09-5.19 and 4.74-4.80 (m, 2H), 3.93-4.62 (m, 6H), 3.81 (s, 3H), 2.10-2.73 (m, 2H)

APCI-MS: (M + 1) = 529 m/z

5

Example 17

Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA

(i) 2,2-Difluoroethyl ester methanesulfonic acid

- 10 To a magnetically stirred solution of 2,2-difluoroethanol (1.52 g, 18.5 mmol) in CH₂Cl₂ (20 mL) under nitrogen was added triethylamine (5.61 g, 55.5 mmol) and methanesulfonyl chloride (2.54 g, 22.2 mmol) at 0°C. The mixture was stirred at 0°C for 1.5 h, diluted with CH₂Cl₂ (50 mL), and washed with 2 N HCl (50 mL). The aqueous layer was extracted with CH₂Cl₂ (30 mL) and the combined organic extracts washed with brine (30 mL), dried
15 (Na₂SO₄), filtered, and concentrated *in vacuo* to afford the sub-title compound (2.52 g, 85%) as a yellow oil which was used without further purification.

¹H NMR (300 MHz, CDCl₃) δ 6.02 (tt, *J* = 3 Hz, *J* = 55 Hz, 1H), 4.39 (dt, *J* = 3 Hz, *J* = 13 Hz, 2H), 3.13 (s, 3H)

20

(ii) 3-Chloro-5-difluoroethoxybenzaldehyde

- To a solution of 3-chloro-5-hydroxybenzaldehyde (1.50 g, 9.6 mmol; see Example 1(ii) above) and potassium carbonate (1.72 g, 12.5 mmol) in DMF (10 mL) under nitrogen was added a solution of 2,2-difluoroethyl ester methanesulfonic acid (2.0 g, 12.5 mmol; see
25 step (i) above) in DMF (10 mL) dropwise at room temperature. The mixture was heated to 100°C for 6 h and then stirred overnight at room temperature. The reaction was cooled to 0°C, poured into ice-cold 2 N HCl (100 mL), and extracted with EtOAc (2 x 75 mL). The combined organic extracts were washed with 0.5 N HCl (2 x 50 mL), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The brown oil was chromatographed on silica gel
30 eluting with Hex:EtOAc (5:1) to afford the sub-title compound (1.35 g, 64%) as a yellow oil.

^1H NMR (300 MHz, CDCl_3) Δ 9.92 (s, 1H), 7.52 (s, 1H), 7.31 (s, 1H), 7.22 (s, 1H), 6.12 (tt, $J = 3$ Hz, $J = 55$ Hz, 1H), 4.26 (dt, $J = 3$ Hz, $J = 15$ Hz, 2H)

5 (iii) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R,S)CH(OTMS)CN}$

To a solution of 3-chloro-5-difluoroethoxybenzaldehyde (1.35 g, 6.1 mmol; see step (ii) above) and zinc iodide (0.48 g, 1.5 mmol) in CH_2Cl_2 (50 mL) was added trimethylsilyl cyanide (1.21 g, 12.2 mmol) dropwise at 0°C under nitrogen. The mixture was stirred at 0°C for 3 h, then diluted with H_2O (50 mL). The organic layer was separated, dried
10 (Na_2SO_4), filtered, and concentrated *in vacuo* to afford the sub-title compound (1.85 g, 95%) as a brown oil which was used without further purification.

^1H NMR (300 MHz, CDCl_3) Δ 7.13 (s, 1H), 6.94 (s, 2H), 6.10 (tt, $J = 3$ Hz, $J = 55$ Hz, 1H), 5.43 (s, 1H), 4.20 (dt, $J = 3$ Hz, $J = 15$ Hz, 2H), 0.28 (s, 9H)

15

(iv) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R,S)CH(OH)C(O)OH}$

Concentrated hydrochloric acid (60 mL) was added to $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R,S)CH(OTMS)CN}$ (1.85 g, 5.8 mmol; see step (iii) above) and stirred at 100°C for 1 h. After cooling to room temperature, the reaction was further cooled to 0°C , basified slowly
20 with 3 N NaOH (~180 mL) and washed with Et_2O (2 x 75 mL). The aqueous layer was acidified with 2 N HCl (20 mL) and extracted with EtOAc (2 x 75 mL). The combined organic extracts were dried (Na_2SO_4), filtered, and concentrated *in vacuo* to afford the sub-title compound (1.50 g, 97%) as a pale yellow solid which was used without further purification.

25

^1H NMR (300 MHz, CD_3OD) Δ 7.15 (s, 1H), 7.05 (s, 1H), 6.98 (s, 1H), 6.19 (tt, $J = 4$ Hz, $J = 55$ Hz, 1H), 5.12 (s, 1H), 4.25 (dt, $J = 4$ Hz, $J = 17$ Hz, 2H)

(v) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(S)CH(OAc)C(O)OH}$ (a) and $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R)CH(OH)C(O)OH}$ (b)

30

A solution of $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R,S)CH(OH)C(O)OH}$ (3.90 g, 14.6 mmol; see step (iv) above) and Lipase PS "Amano" (2.50 g) in vinyl acetate (140 mL) and MTBE (140

mL) was heated at 70°C under nitrogen for 40 h. The reaction was cooled to room temperature, the enzyme removed by filtration washing with EtOAc, and the filtrate concentrated *in vacuo*. Chromatography on silica gel eluting with CHCl₃:MeOH:Et₃N (92:6:2) afforded the triethylamine salt of the sub-title compound (a) as a yellow oil. In addition, the triethylamine salt of the sub-title compound (b) (1.47 g) was obtained and the salt was dissolved in H₂O (100 mL), acidified with 2 N HCl and extracted with EtOAc (2 x 75 mL). The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated *in vacuo* to yield the sub-title compound (b) (1.00 g) as an off-white solid.

10 Data for sub-title compound (b):

mp: 103–106°C

R_f = 0.39 (90:8:2 CHCl₃:MeOH:Et₃N)

¹H NMR (300 MHz, CD₃OD) Δ 7.13 (s, 1H), 7.04 (s, 1H), 6.97 (s, 1H), 6.17 (tt, *J* = 4 Hz, *J* = 55 Hz, 1H), 5.12 (s, 1H), 4.24 (dt, *J* = 4 Hz, *J* = 8 Hz, 2H).

15 ¹³C NMR (75 MHz, CD₃OD) Δ 175.5, 160.3, 144.5, 136.1, 121.3, 115.7, 115.3, (t, *J* = 240 Hz), 112.9, 73.4, 68.6 (t, *J* = 29 Hz)

HPLC Analysis: 96.2%, >95.0% ee, ChiralPak AD Column (95:5:0.5 Hex:EtOH:TFA mobile phase)

[α]_D²⁵ = -84.0° (c = 0.85 MeOH)

20 APCI-MS: (*M* - 1) = 265 m/z

(vi) Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

To a solution of Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)OH (0.35 g, 1.3 mmol; see step (v) above (compound (b))) in DMF (18 mL) under nitrogen was added H-Aze-Pab(Teoc) x HCl (0.76 g, 1.7 mmol), PyBOP (0.75 g, 1.4 mmol), and DIPEA (0.43 g, 3.3 mmol) at 0°C. The reaction was stirred at 0°C for 2 h and then at room temperature for 20 h. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first with CHCl₃:EtOH (10:1), and then with EtOAc:EtOH (10:1) to afford the sub-title compound (0.69 g, 84%) as a crushable white foam.

30

mp: 108–118°C

R_f = 0.48 (10:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.78-7.81 (m, 2H), 7.40-7.43 (m, 2H), 7.09-7.12 (m, 1H), 6.96-7.02 (m, 2H), 6.16 (t, $J = 57$ Hz, 1H), 5.09-5.20 and 4.75-4.80 (m, 2H), 3.95-4.55 (m, 8H), 2.10-2.75 (m, 2H), 1.04-1.11 (m, 2H), 0.07 (s, 9H)
APCI-MS: (M + 1) = 625 m/z

5

(vii) Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.086 g, 0.138 mmol; see step (vi) above), was dissolved in 3 mL of TFA and allowed to react for 1 h. TFA was evaporated and the residue was freeze dried from water/acetonitrile to yield 0.080 g (98%)
10 of the title compound as its TFA salt.

¹H-NMR (300 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.6-7.5 (m, 2H), 7.15-6.95 (m, 3H), 6.35-5.95 (m, 1H), 5.20 (m, 1H, minor rotamer), 5.14 (s, 1H, major rotamer), 5.10 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.6-4.0 (m, 6H), 2.70 (m, 1H, minor rotamer),
15 2.53 (m, 1H, major rotamer), 2.29 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 174.0, 173.8, 173.4, 172.9, 168.2

MS (m/z) 481.2 (M+1)⁺

20

Example 18

Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)

To a solution of Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)OH (0.30 g, 1.7 mmol; see Example 17(v) above (compound (b))) in DMF (15 mL) under nitrogen was added H-Aze-Pab(OMe) x 2HCl (0.49 g, 1.5 mmol), PyBOP (0.65 g, 1.2 mmol), and DIPEA (0.36 g, 2.8 mmol) at 0°C. The reaction was stirred at 0°C for 2 h and then at room temperature for 20 h. The mixture was concentrated *in vacuo* and the residue chromatographed three times on silica gel, eluting first with CHCl₃:EtOH (10:1), then with EtOAc:EtOH (10:1), and finally with CHCl₃:MeOH (20:1) to afford the title compound (0.47 g, 81%) as a crushable white
25 foam.
30

mp: 65-75°C

$R_f = 0.37$ (10:1 CHCl_3 :EtOH)

^1H NMR (300 MHz, CD_3OD , complex mixture of rotamers) Δ 7.58-7.60 (m, 2H), 7.32-7.35 (m, 2H), 7.09-7.12 (m, 1H), 6.96-7.02 (m, 2H), 6.16 (t, $J = 55$ Hz, 1H), 5.08-5.18 and
5 4.74-4.80 (m, 2H), 3.96-4.50 (m, 6H), 3.80 (s, 3H), 2.10-2.75 (m, 2H)

APCI-MS: $(M + 1) = 511$ m/z.

Example 19

Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab x TFA

10

(i) Ph(3-Cl)(5-TMSO)-(R,S)CH(OTMS)CN

To a solution of 3-chloro-5-hydroxybenzaldehyde (9.8 g, 62.6 mmol; see Example 1(ii) above) and ZnI_2 (5.0 g, 15.7 mmol) in anhydrous CH_2Cl_2 (500 mL) at 0°C was added trimethylsilyl cyanide (13.7 g, 138 mmol). The reaction mixture was allowed to warm to
15 room temperature and stirred overnight. Water (250 mL) was added, and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (2 x 300 mL). The combined organic extracts were dried (Na_2SO_4), filtered, and concentrated *in vacuo* to afford the sub-title compound (16.9 g, 83%) as a yellow oil that was used without further purification.

20 $R_f = 0.42$ (3:1 Hex:EtOAc)

^1H NMR (300 MHz, CDCl_3) δ 7.06 (s, 1H), 6.86 (s, 2H), 5.40 (s, 1H), 0.30 (s, 9 H), 0.24 (s, 9 H).

(ii) Ph(3-Cl)(5-OH)-(R,S)CH(OH)C(O)OH

25 A solution of Ph(3-Cl)(5-OTMS)-(R,S)CH(OTMS)CN (22.6 g, 68.8 mmol; see step (i) above) in concentrated HCl (200 mL) was refluxed under nitrogen for 3 h. The reaction was cooled to 0°C and basified slowly with 2N NaOH. The mixture was washed with Et_2O (3 x 100 mL) to remove the organic impurities. The aqueous layer was acidified with 2N HCl and extracted with EtOAc (3 x 200 mL). The combined organic extracts were
30 dried (Na_2SO_4), filtered, and concentrated *in vacuo* to afford the sub-title compound (9.3 g, 67%) as a brown oil that was used without further purification.

$R_f = 0.23$ (6:3:1 CHCl_3 :MeOH:concentrated NH_4OH)

^1H NMR (300 MHz, CD_3OD) δ 7.05 (s, 1H), 6.94 (s, 1H), 6.73 (s, 1H), 5.03 (s, 1H).

(iii) $\text{Ph}(3\text{-Cl})(5\text{-OH})\text{-(R,S)CH(OH)C(O)OEt}$

5 To a solution of $\text{Ph}(3\text{-Cl})(5\text{-OH})\text{-(R,S)CH(OH)C(O)OH}$ (9.3 g, 46.0 mmol; see step (ii) above) in absolute EtOH (200 mL) was added concentrated sulfuric acid (0.25 mL) and the reaction was refluxed under nitrogen for 4 h. The reaction was cooled to 0°C and solid NaHCO_3 (0.2 g) was added. The reaction was concentrated *in vacuo* and partitioned with saturated NaHCO_3 (100 mL) and Et_2O (3 x 50 mL). The combined organic extracts were
10 dried (Na_2SO_4), filtered, and concentrated *in vacuo* to give the sub-title compound (6.9 g, 65%) as a yellow oil which was used without further purification.

$R_f = 0.62$ (6:3:1 CHCl_3 :MeOH:concentrated NH_4OH).

^1H NMR (300 MHz, CDCl_3) δ 6.99 (s, 1H), 6.81 (s, 2H), 5.07 (s, 1H), 4.16-4.32 (m, 2H),
15 1.23 (t, $J = 7$ Hz, 3H).

(iv) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{F})\text{-(R,S)CH(OH)C(O)OEt}$

To a solution of $\text{Ph}(3\text{-Cl})(5\text{-OH})\text{-(R,S)CH(OH)C(O)OEt}$ (6.1 g, 26.8 mmol; see step (iii) above) in DMF (100 mL) in a sealed flask under nitrogen at 0°C was added cesium carbonate (13.1 g, 40.2 mmol). The reaction mixture was stirred at 0°C for 15 minutes
20 followed by the addition of potassium iodide (0.5 g, 2.7 mmol). The reaction was cooled to -78°C and chlorofluoromethane (18.4 g, 268 mmol) was bubbled into the vessel. The sealed flask was then allowed to warm to room temperature and stirred for 18 h. The reaction mixture was cooled to 0°C , vented carefully to remove any excess
25 chlorofluoromethane, and partitioned with H_2O (20 mL) and Et_2O (3 x 50 mL). The combined organics were washed with brine (2 x 50 mL), dried (Na_2SO_4), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (gradient from 9:1 to 3:1) afforded the sub-title compound (2.4 g, 35%) as a light yellow oil. Note: The compound is faintly uv-visible on TLC. It can be visualised by staining the
30 TLC with bromocresol green.

$R_f = 0.46$ (2:1 Hex:EtOAc)

^1H NMR (300 MHz, CDCl_3) δ 7.21 (s, 1H), 7.08 (s, 1H), 7.05 (s, 1H), 5.70 (d, $J_{\text{H-F}} = 54$ Hz, 2H), 5.12 (d, $J = 5$ Hz, 1H), 3.80–4.35 (m, 2H), 3.50 (d, $J = 5$ Hz, 1H), 1.26 (t, $J = 7$ Hz, 3H).

5 (v) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{F})\text{-(R,S)CH(OH)C(O)OH}$

To a solution of $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{F})\text{-(R,S)CH(OH)C(O)OEt}$ (1.8 g, 6.8 mmol; see step (iv) above) in $\text{H}_2\text{O}:\text{THF}$ (30 mL, 1:2) at 0°C under nitrogen was added lithium hydroxide monohydrate (0.40 g, 10.3 mmol). The mixture was stirred at 0°C for 2 h. The reaction mixture was concentrated *in vacuo* and partitioned with H_2O (5 mL) and Et_2O (2 x 20 mL).
10 The aqueous layer was acidified carefully with 0.2N HCl at 0°C and extracted with EtOAc (3 x 30 mL). The combined organics were dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (1.4 g, 87%) as a colourless oil which solidified to a white solid upon standing.

15 $R_f = 0.43$ (6:2:1 $\text{CHCl}_3:\text{MeOH}:\text{Et}_3\text{N}$)

^1H NMR (300 MHz, CD_3OD) δ 7.24 (s, 1H), 7.17 (s, 1H), 7.07 (s, 1H), 5.78 (d, $J_{\text{H-F}} = 54$ Hz, 2H), 5.13 (s, 1H).

20 (vi) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{F})\text{-(R)CH(OH)C(O)OH}$ (a) and $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{F})\text{-(S)CH(OAc)C(O)OH}$ (b)

A mixture of $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{F})\text{-(R,S)CH(OH)C(O)OH}$ (3.2 g, 13.9 mmol; see step (v) above) and Lipase PS "Amano" (1.9 g) in vinyl acetate (150 mL) and MTBE (150 mL) was heated at 70°C under a nitrogen atmosphere for 3 d. The reaction mixture was cooled, filtered through Celite® and the filter cake washed with EtOAc . The filtrate was
25 concentrated *in vacuo* and subjected to flash chromatography on a silica gel eluting with $\text{CHCl}_3:\text{MeOH}:\text{Et}_3\text{N}$ (15:1:0.5) to afford the triethylamine salt of the sub-title compound (a) (0.50 g, 21%) that was used without neutralisation. In addition, the triethylamine salt of the sub-title compound (b) (0.46 g, 20%) was obtained.

30 Data for Sub-Title Compound (a):

$R_f = 0.19$ (15:1:0.5 $\text{CHCl}_3:\text{MeOH}:\text{Et}_3\text{N}$)

¹H NMR (300 MHz, CD₃OD) δ 7.26 (s, 1H), 7.18 (s, 1H), 6.97 (s, 1H), 5.74 (d, *J*_{H-F} = 54 Hz, 2H), 4.81 (s, 1H), 3.17 (q, *J* = 7 Hz, 6H), 1.28 (t, *J* = 7 Hz, 9H).

Data for Sub-Title Compound (b)

5 *R*_f = 0.33 (15:1:0.5 CHCl₃:MeOH:Et₃N)

¹H NMR (300 MHz, CD₃OD) δ 7.28 (s, 1H), 7.19 (s, 1H), 7.09 (s, 1H), 5.76 (d, *J*_{H-F} = 54 Hz, 2H), 5.75 (s, 1H), 3.17 (q, *J* = 7 Hz, 6H), 2.16 (s, 3H), 1.28 (t, *J* = 7 Hz, 9H).

(vii) Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

10 To a solution of the triethylamine salt of Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)OH (0.50 g, 1.50 mmol; see step (vi) above) and HAZE-Pab(Teoc)•HCl (0.87 g, 1.90 mmol) in dry DMF (15 mL) under nitrogen at 0°C was added PyBOP (0.85 g, 2.60 mmol) and DIPEA (0.48 g, 3.70 mmol). The reaction was allowed to warm to room temperature and stirred overnight. The reaction mixture was concentrated *in vacuo* and flash chromatographed
15 twice on silica gel, eluting first with CHCl₃:EtOH (9:1) and second with EtOAc:EtOH (20:1) to afford the sub-title compound (0.23 g, 26%) as a crushable white foam.

Mp: 88-92°C

*R*_f = 0.61 (9:1 CHCl₃:EtOH)

20 ¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.81 (d, *J* = 8 Hz, 2H), 7.40-7.42 (m, 2H), 7.06-7.23 (m, 3H), 5.76 (d, *J*_{H-F} = 51 Hz, 2H), 5.10-5.16 and 4.77-4.83 (m, 2H), 3.80-4.49 (m, 6H), 2.30-2.53 (m, 2H), 1.08 (t, *J* = 7 Hz, 2H), 0.08 (s, 9H).

APCI-MS (*M* + 1) = 593 *m/z*

25 (viii) Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.051 g, 0.086 mmol; see step (vii) above), was dissolved in 3 mL of TFA and allowed to react for 20 min. TFA was evaporated and the residue was freeze dried from water/acetonitrile. The product was 95% pure with 5% of defluoromethylated material. Attempts to purify it by preparative RPLC
30 with CH₃CN:0.1M NH₄OAc failed, and the material, partially as an acetate, was dissolved in 5 mL of TFA, evaporated and freeze dried to yield 26 mg (51%) of the title compound as its TFA salt. Purity: 95%.

¹H-NMR (600 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.6-7.5 (m, 2H), 7.21 (s, 1H, major rotamer), 7.17 (s, 1H, minor rotamer), 7.13 (s, 1H, major rotamer), 7.09 (s, 1H, minor rotamer), 7.07 (m, 1H, major rotamer), 7.04 (m, 1H, minor rotamer), 5.73 (d, 2H),
5 5.18 (m, 1H, minor rotamer), 5.16 (s, 1H, major rotamer), 5.09 (s, 1H, minor rotamer),
4.78 (m, 1H, minor rotamer), 4.56 (d, 1H, major rotamer), 4.50 (d, 1H, minor rotamer),
4.46 (d, 1H, minor rotamer), 4.45 (d, 1H, major rotamer), 4.35 (m, 1H, major rotamer),
4.14 (m, 1H, major rotamer), 4.05 (m, 1H, minor rotamer), 3.97 (m, 1H, minor rotamer),
2.68 (m, 1H, minor rotamer), 2.52 (m, 1H, major rotamer), 2.28 (m, 1H, major rotamer),
10 2.19 (m, 1H, minor rotamer).
¹³C-NMR (150 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 173.9,
173.3, 172.9, 168.2.
ESI-MS+: (M+1) = 449 (m/z)

15 Example 20

Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(OMe)

To a solution of the triethylamine salt of Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)OH (0.60 g, 1.80 mmol; see Example 19(vi)) and Haze-Pab(OMe)•2HCl (0.79 g, 2.30 mmol) in DMF (15 mL) under nitrogen at 0°C was added PyBOP (1.04 g, 1.90 mmol) and DIPEA (0.58 g,
20 4.50 mmol). The reaction was allowed to warm to room temperature and stirred overnight. The reaction mixture was concentrated *in vacuo* and flash chromatographed three times on silica gel, eluting first with CHCl₃:EtOH (9:1) and then twice with EtOAc:EtOH (20:1) to afford the title compound (0.22 g, 26%) as a crushable white foam.

25 Mp: 66-70°C

R_f = 0.45 (9:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.59 (d, *J* = 8 Hz, 2H), 7.32 (d, *J* = 7 Hz, 2H), 7.06-7.23 (m, 3H), 5.75 (s, *J*_{H-F} = 54 Hz, 1H), 5.10-5.16 and 4.78-4.84 (m, 2H), 4.11-4.45 (m, 4H), 3.80 (s, 3H), 2.10-2.75 (m, 2H).

30 ¹³C-NMR (150 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 173.0, 170.8, 170.7, 152.5.

APCI-MS: (M + 1) = 479 m/z

Example 21Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab x TFA(i) (2-Monofluoroethyl) methanesulfonate

To a magnetically stirred solution of 2-fluoroethanol (5.0 g, 78.0 mmol) in CH₂Cl₂ (90 mL) under nitrogen at 0°C was added triethylamine (23.7 g, 234 mmol) and methanesulfonyl chloride (10.7 g, 93.7 mmol). The mixture was stirred at 0°C for 1.5 h, diluted with CH₂Cl₂ (100 mL) and washed with 2N HCl (100 mL). The aqueous layer was extracted with CH₂Cl₂ (50 mL) and the combined organic extracts washed with brine (75 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (9.7 g, 88%) as a yellow oil which was used without further purification.

¹H NMR (300 MHz, CDCl₃) δ 4.76 (t, *J* = 4 Hz, 1H), 4.64 (t, *J* = 4 Hz, 1H), 4.52 (t, *J* = 4 Hz, 1H), 4.43 (t, *J* = 4 Hz, 1H), 3.09 (s, 3H).

(ii) 3-Chloro-5-monofluoroethoxybenzaldehyde

To a solution of 3-chloro-5-hydroxybenzaldehyde (8.2 g, 52.5 mmol; see Example 1(ii) above) and potassium carbonate (9.4 g, 68.2 mmol) in DMF (10 mL) under nitrogen was added a solution of (2-monofluoroethyl) methanesulfonate (9.7 g, 68.2 mmol; see step (i) above) in DMF (120 mL) dropwise at room temperature. The mixture was heated to 100°C for 5 h and then stirred overnight at room temperature. The reaction was cooled to 0°C, poured into ice-cold 2N HCl and extracted with EtOAc. The combined organic extracts were washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The brown oil was chromatographed on silica gel eluting with Hex:EtOAc (4:1) to afford the sub-title compound (7.6 g, 71%) as a yellow oil.

¹H NMR (300 MHz, CDCl₃) δ 9.92 (s, 1H), 7.48 (s, 1H), 7.32 (s, 1H), 7.21 (s, 1H), 4.87 (t, *J* = 4 Hz, 1H), 4.71 (t, *J* = 3 Hz, 1H), 4.33 (t, *J* = 3 Hz, 1H), 4.24 (t, *J* = 3 Hz, 1H).

(iii) Ph(3-Cl)(5-OCH₂CH₂F)-(R,S)CH(OTMS)CN

To a solution of 3-chloro-5-monofluoroethoxybenzaldehyde (7.6 g, 37.5 mmol; see step (ii) above) and zinc iodide (3.0 g, 9.38 mmol) in CH_2Cl_2 (310 mL) was added trimethylsilyl cyanide (7.4 g, 75.0 mmol) dropwise at 0°C under nitrogen. The mixture was stirred at 0°C for 3 h and at room temperature overnight. The reaction was diluted with H_2O (300 mL), the organic layer was separated, dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (10.6 g, 94%) as a brown oil that was used without further purification or characterisation.

(iv) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R,S)CH(OH)C(O)OH}$

Concentrated hydrochloric acid (100 mL) was added to $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R,S)CH(OTMS)CN}$ (10.6 g, 5.8 mmol; see step (iii) above) and the solution stirred at 100°C for 3 h. After cooling to room temperature, the reaction was further cooled to 0°C , basified slowly with 3N NaOH (~300 mL) and washed with Et_2O (3 x 200 mL). The aqueous layer was acidified with 2N HCl (80 mL) and extracted with EtOAc (3 x 300 mL). The combined EtOAc extracts were dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (8.6 g, 98%) as a pale yellow solid that was used without further purification.

$R_f = 0.28$ (90:8:2 CHCl_3 :MeOH:concentrated NH_4OH)

^1H NMR (300 MHz, CD_3OD) δ 7.09 (s, 1H), 7.02 (s, 1H), 6.93 (s, 1H), 5.11 (s, 1H), 4.77-4.81 (m, 1H), 4.62-4.65 (m, 1H), 4.25-4.28 (m, 1H), 4.15-4.18 (m, 1H).

(v) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(S)CH(OAc)C(O)OH}$ (a) and $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R)CH(OH)C(O)OH}$ (b)

A solution of $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R,S)CH(OH)C(O)OH}$ (8.6 g, 34.5 mmol; see step (iv) above) and Lipase PS "Amano" (4.0 g) in vinyl acetate (250 mL) and MTBE (250 mL) was heated at 70°C under nitrogen for 3 d. The reaction was cooled to room temperature and the enzyme removed by filtration through Celite®. The filter cake was washed with EtOAc and the filtrate concentrated *in vacuo*. Chromatography on silica gel eluting with CHCl_3 :MeOH: Et_3N (90:8:2) afforded the triethylamine salt of sub-title compound (a) as a yellow oil. In addition, the triethylamine salt of sub-title compound (b) (4.0 g) was obtained. The salt of sub-title compound (b) was dissolved in H_2O (250 mL), acidified

with 2N HCl and extracted with EtOAc (3 x 200 mL). The combined organic extracts were dried (Na_2SO_4), filtered and concentrated *in vacuo* to yield the sub-title compound (b) (2.8 g, 32%) as a yellow oil.

5 Data for Sub-Title Compound (b):

$R_f = 0.28$ (90:8:2 CHCl_3 :MeOH:concentrated NH_4OH)

^1H NMR (300 MHz, CD_3OD) δ 7.09 (s, 1H), 7.02 (s, 1H), 6.93 (s, 1H), 5.11 (s, 1H), 4.77-4.81 (m, 1H), 4.62-4.65 (m, 1H), 4.25-4.28 (m, 1H), 4.15-4.18 (m, 1H).

10 (vi) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R)CH(OH)C(O)-Aze-Pab(Teoc)}$

To a solution of $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R)CH(OH)C(O)OH}$ (940 mg, 3.78 mmol; see step (v) above) in DMF (30 mL) under nitrogen at 0°C was added HAze-Pab(Teoc)HCl (2.21 g, 4.91 mmol), PyBOP (2.16 g, 4.15 mmol), and DIPEA (1.22 g, 9.45 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature for 4 h. The mixture was
15 concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first with CHCl_3 :EtOH (15:1) and second with EtOAc:EtOH (20:1) to afford the sub-title compound (450 mg, 20%) as a crushable white foam.

Mp: $80\text{-}88^\circ\text{C}$

20 $R_f = 0.60$ (10:1 CHCl_3 :EtOH)

^1H NMR (300 MHz, CD_3OD , complex mixture of rotamers) δ 7.79 (d, $J = 8$ Hz, 2H), 7.42 (d, $J = 8$ Hz, 2H), 7.05-7.08 (m, 1H), 6.93-6.99 (m, 2H), 5.08-5.13 (m, 1H), 4.75-4.80 (m, 2H), 4.60-4.68 (m, 1H), 3.95-4.55 (m, 8H), 2.10-2.75 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

25 APCI-MS: $(M + 1) = 607$ m/z.

(vii) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R)CH(OH)C(O)-Aze-Pab} \times \text{TFA}$

$\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-(R)CH(OH)C(O)-Aze-Pab(Teoc)}$ (0.357 g, 0.589 mmol; see step (vi) above), was dissolved in 10 mL of TFA and allowed to react for 40 min. TFA was
30 evaporated and the residue was freeze dried from water/acetonitrile to yield 0.33 g (93%) of the title compound as its TFA salt.

¹H-NMR (600 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.54 (d, 2H), 7.08 (s, 1H, major rotamer), 7.04 (s, 1H, minor rotamer), 6.99 (s, 1H, major rotamer), 6.95 (s, 1H), 6.92 (s, 1H, minor rotamer), 5.18 (m, 1H, minor rotamer), 5.14 (s, 1H, major rotamer), 5.08 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.73 (m, 1H), 4.65 (m, 1H), 4.6-4.4 (m, 2H),
5 4.35 (m, 1H, major rotamer), 4.21 (doublet of multiplets, 2H), 4.12 (m, 1H, major rotamer), 4.06 (m, 1H, minor rotamer), 3.99 (m, 1H, minor rotamer), 2.69 (m, 1H, minor rotamer), 2.53 (m, 1H, major rotamer), 2.29 (m, 1H, major rotamer), 2.14 (m, 1H, minor rotamer).

¹³C-NMR (150 MHz; CD₃OD): (carbonyl and/or amidine carbons) δ 172.8, 172.1, 167.4.

10 ESI-MS+: (M+1) = 463 (m/z)

Example 22

Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OMe)

To a solution of Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)OH (818 mg, 3.29 mmol; see
15 Example 21(v) above) in DMF (30 mL) under nitrogen at 0°C was added Haze-Pab(OMe)•2HCl (1.43 g, 4.27 mmol), PyBOP (1.89 g, 3.68 mmol), and DIPEA (1.06 g, 8.23 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed two
20 times on silica gel, eluting first with CHCl₃:EtOH (15:1) and second with EtOAc:EtOH (20:1) to afford the title compound (880 mg, 54%) as a crushable white foam.

Mp: 65-72°C

R_f = 0.60 (10:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.58-7.60 (d, *J* = 8 Hz, 2H),
25 7.34 (d, *J* = 7 Hz, 2H), 7.05-7.08 (m, 2H), 6.95-6.99 (m, 1H), 5.08-5.13 (m, 1H), 4.77-4.82 (m, 1H), 4.60-4.68 (m, 1H), 3.99-4.51 (m, 7H), 3.82 (s, 3H), 2.10-2.75 (m, 2H).

¹³C-NMR (150 MHz; CD₃OD): (carbonyl and/or amidine carbons) δ 173.3, 170.8, 152.5.

APCI-MS: (M + 1) = 493 m/z.

Example 23Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab x TFA(i) 1,3-Difluoroisopropyl methanesulfonate

5 To a magnetically stirred solution of 1,3-difluoro-2-propanol (7.0 g, 72.8 mmol) in CH₂Cl₂ (100 mL) under nitrogen at 0°C was added triethylamine (22.1 g, 219 mmol) and methanesulfonyl chloride (10.0 g, 87.4 mmol). The mixture was stirred at 0°C for 3 h. The mixture was washed with 2N HCl (150 mL) and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (200 mL) and the combined organic extracts
10 washed with brine (100 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (11.5 g, 91%) as a yellow oil which was used without further purification.

15 ¹H NMR (300 MHz, CDCl₃) δ 4.97-5.08 (m, 1H), 4.75-4.77 (m, 2H), 4.59-4.61 (m, 2H), 3.12 (s, 3H).

(ii) Ph(3-Cl)(5-OCH(CH₂F)₂)CHO

To a solution of 3-chloro-5-hydroxybenzaldehyde (8.0 g, 50.7 mmol; see Example 1(ii) above) and potassium carbonate (9.1 g, 66.0 mmol) in DMF (75 mL) under nitrogen was
20 added a solution of 1,3-difluoroisopropyl methanesulfonate (11.5 g, 66.0 mmol; see step (i) above) in DMF (75 mL) dropwise at room temperature. The mixture was heated to 110°C for 18 h. The reaction was cooled to 0°C, poured into ice-cold 2N HCl (200 mL) and extracted with EtOAc (3 x 250 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo*. The brown oil was chromatographed on silica gel
25 eluting with Hex:EtOAc (4:1) to afford the sub-title compound (4.4 g, 37%) as a yellow oil.

¹H NMR (300 MHz, CDCl₃) δ 9.92 (s, 1H), 7.51 (s, 1H), 7.36 (s, 1H), 7.26 (s, 1H), 4.70-4.89 (m, 3H), 4.63-4.68 (m, 2H).

(iii) Ph(3-Cl)(5-OCH(CH₂F)₂)-(R,S)CH(OTMS)CN

To a solution of $\text{Ph(3-Cl)(5-OCH(CH}_2\text{F)}_2\text{)CHO}$ (4.4 g, 18.7 mmol; see step (ii) above) and zinc iodide (1.5 g, 4.67 mmol) in CH_2Cl_2 (200 mL) at 0°C under nitrogen was added trimethylsilyl cyanide (3.7 g, 37.3 mmol) dropwise. The mixture was stirred at 0°C for 3 h and overnight at room temperature, then diluted with H_2O (200 mL). The organic layer
5 was separated, dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (5.5 g, 87%) as a brown oil that was used without further purification.

$^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.12 (s, 1H), 7.00 (s, 2H), 5.42 (s, 1H), 4.70-4.80 (m, 3H), 4.59-4.64 (m, 2H), 0.26 (s, 9H).

10

(iv) $\text{Ph(3-Cl)(5-OCH(CH}_2\text{F)}_2\text{)-(R,S)CH(OH)C(O)OH}$

Concentrated hydrochloric acid (50 mL) was added to $\text{Ph(3-Cl)(5-OCH(CH}_2\text{F)}_2\text{)-(R,S)CH(OTMS)CN}$ (5.5 g, 16.3 mmol; see step (iii) above) and the solution stirred at 100°C for 1.5 h. After cooling to room temperature, the reaction was further cooled to
15 0°C , basified slowly with 3N NaOH (~200 mL) and washed with Et_2O (3 x 200 mL). The aqueous layer was acidified with 2N HCl (75 mL) and extracted with EtOAc (3 x 200 mL). The combined EtOAc extracts were dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (4.6 g, 100%) as a brown oil that was used without further purification.

20

$^1\text{H NMR}$ (300 MHz, CD_3OD) δ 7.14 (s, 1H), 7.08 (s, 1H), 7.02 (s, 1H), 5.12 (s, 1H), 4.70-4.90 (m, 3H), 4.52-4.67 (m, 2H).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

(v) Ph(3-Cl)(5-OCH(CH₂F)₂)-(S)CH(OAc)C(O)OH (a) and Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)OH (b)

A solution of Ph(3-Cl)(5-OCH(CH₂F)₂)-(R,S)CH(OH)C(O)OH (4.6 g, 16.4 mmol; see step (iv) above) and Lipase PS "Amano" (3.0 g) in vinyl acetate (150 mL) and MTBE (150 mL) was heated at 70°C under nitrogen for 2.5 d. The reaction was cooled to room temperature, the enzyme removed by filtration through Celite®. The filter cake was washed with EtOAc and the filtrate concentrated *in vacuo*. Chromatography on silica gel eluting with CHCl₃:MeOH:Et₃N (90:8:2) afforded the triethylamine salt of the sub-title compound (a) as a yellow oil. In addition, the triethylamine salt of the sub-title compound (b) (2.2 g) was obtained and the salt was dissolved in H₂O (100 mL), acidified with 2N HCl and extracted with EtOAc (3 x 200 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to yield the sub-title compound (b) (1.4 g, 29%) as a yellow oil.

15 Data for Sub-Title Compound (b):

¹H NMR (300 MHz, CD₃OD) δ 7.14 (s, 1H), 7.08 (s, 1H), 7.02 (s, 1H), 5.12 (s, 1H), 4.70-4.90 (m, 3H), 4.52-4.67 (m, 2H).

(vi) Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

20 To a solution of Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)OH (824 mg, 2.94 mmol; see step (v) above) in DMF (30 mL) under nitrogen at 0°C was added HAze-Pab(Teoc)•HCl (1.71 g, 3.81 mmol), PyBOP (1.68 g, 3.23 mmol), and DIPEA (949 mg, 7.34 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first with CHCl₃:EtOH (15:1), and second with EtOAc:EtOH (20:1) to afford the sub-title compound (720 mg, 38%) as a crushable white foam.

Mp: 78-84°C

R_f = 0.62 (10:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.79 (d, *J* = 8 Hz, 2H), 7.42 (d, *J* = 8 Hz, 2H), 7.00-7.12 (m, 3H), 5.08-5.20 (m, 1H), 3.97-4.80 (m, 12H), 2.10-2.75 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

30 APCI-MS: (*M* + 1) = 639 *m/z*.

(vii) Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.129 g, 0.202 mmol; see step (vi) above), was dissolved in 3 mL of TFA and allowed to react for 20 min. TFA was evaporated and the residue was freeze dried from water/acetonitrile to yield 0.123 g (100%) of the title compound as its TFA salt.

¹H-NMR (400 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.55 (d, 2H), 7.2-7.0 (m, 3H), 5.18 (m, 1H, minor rotamer), 5.15 (s, 1H, major rotamer), 5.08 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer partly obscured by the CD₃OH peak), 4.75-4.4 (m, 7H), 4.38 (m, 1H, major rotamer), 4.15 (m, 1H, major rotamer), 4.1-3.9 (m, 2H, 2 signals from minor rotamer), 2.70 (m, 1H, minor rotamer), 2.53 (m, 1H, major rotamer), 2.30 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 172.9, 172.6, 172.2, 171.7, 167.1.

ESI-MS+: (M+1) = 495 (m/z)

Example 24

Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)

To a solution of Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)OH (513 mg, 1.83 mmol; see Example 23(v) above) in DMF (30 mL) under nitrogen at 0°C was added Haze-Pab(OMe)·2HCl (797 mg, 2.38 mmol), PyBOP (1.04 g, 2.01 mmol), and DIPEA (591 mg, 4.57 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed two times on silica gel, eluting first with CHCl₃:EtOH (15:1) and second with EtOAc:EtOH (20:1) to afford the title compound (370 mg, 39%) as a crushable white foam.

Mp: 58-63°C

R_f = 0.66 (10:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.58-7.60 (d, *J* = 8 Hz, 2H), 7.34 (d, *J* = 8 Hz, 2H), 7.00-7.12 (m, 3H), 5.08-5.20 (m, 1H), 4.65-4.82 (m, 3H), 4.28-4.65 (m, 5H), 3.92-4.18 (m, 2H), 3.82 (s, 3H), 2.10-2.75 (m, 2H).

^{13}C -NMR (150 MHz; CD_3OD): (carbonyl and/or amidine carbons) δ 173.2, 170.8, 152.5.

APCI-MS: $(M + 1) = 525$ m/z.

Example 25

5 $\text{Ph}(3\text{-F})(5\text{-OCHF}_2)\text{-(R)CH(OH)C(O)-Aze-Pab} \times \text{TFA}$

(i) 1-Bromo-3-fluoro-5-benzyloxybenzene

Sodium hydride (60% dispersion in oil, 24.0 g, 0.48 mol) was added portionwise to a stirred solution of anhydrous benzyl alcohol (64.5 g, 0.60 mol) in THF (1.0 L). After the
10 mixture was stirred for 1 h, a solution of 1-bromo-3,5-difluorobenzene (76.8 g, 0.40 mmol) in THF (100 mL) was added dropwise over a period of 1 h. The reaction was stirred at room temperature for 2 d. Water (400 mL) was added and the THF was removed *in vacuo*. The aqueous layer was extracted with hexane (3 x 150 mL). The combined organic
15 extracts were washed with 2N NaOH (2 x 100 mL) then, dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (110.7 g, 98%) as a light yellow oil that was used without further purification.

$R_f = 0.47$ (Hex)

^1H NMR (300 MHz, CDCl_3) δ 7.36-7.41 (m, 5H), 6.94 (bs, 1H), 6.87 (d,
20 $J_{\text{H-F}} = 8$ Hz, 1H), 6.63 (d, $J_{\text{H-F}} = 10$ Hz, 1H), 5.03 (s, 2H).

(ii) 3-Bromo-5-fluorophenol

To a solution of 1-bromo-3-fluoro-5-benzyloxybenzene (110.0 g, 0.39 mol; see step (i) above) and *N,N*-dimethylaniline (474.0 g, 3.92 mol) in anhydrous CH_2Cl_2 (1.0 L) at 0°C
25 was added aluminium chloride (156.0 g, 1.17 mol). After 10 min, the ice bath was removed and the stirring was continued for 2 h. The reaction was quenched by the cautious addition of 3N HCl (600 mL). The layers were separated, and the aqueous layer was extracted with CH_2Cl_2 (2 x 150 mL). The combined organic extracts were washed with 2N HCl (250 mL) and H_2O (3 x 250 mL). To the organic layer was added 15% KOH
30 (500 mL), and the layers were separated. The organic layer was further extracted with 2 N KOH (2 x 70 mL). The combined aqueous layers were washed with CH_2Cl_2 (3 x 100 mL) and then acidified with 4N HCl. The aqueous layer was extracted with Et_2O (3 x 125 mL)

then, the combined Et₂O extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (69.0 g, 92%) as a brown oil that was used without further purification.

5 Mp: 33-35°C

R_f = 0.25 (CHCl₃)

¹H NMR (300 MHz, DMSO-*d*₆) δ 10.38 (s, 1H), 6.90 (dd, *J*_{H-F} = 11 Hz, *J* = 2 Hz, 1H), 6.81 (s, 1H), 6.59 (dt, *J*_{H-F} = 11 Hz, *J* = 2 Hz, 1H).

APCI-MS: (M-1) = 189 m/z

10 (iii) 1-Bromo-3-fluoro-5-difluoromethoxybenzene

A mixture of 3-bromo-5-fluorophenol (6.1 g, 31.0 mmol; see step (ii) above) and chlorodifluoromethane (13.0 g, 150.0 mmol) in *i*-PrOH (100 mL) and 30% KOH (80 mL) was heated in a sealed flask for 18 h at 80-85°C. The reaction mixture was cooled to room
15 temperature and the layers were separated. The organic layer was concentrated *in vacuo* to afford a colourless oil. The aqueous layer was extracted with Et₂O (3 x 30 mL). The crude oil and the combined organic extracts were washed with 2N NaOH (3 x 30 mL) and H₂O (3 x 30 mL). The organics were then dried (Na₂SO₄), filtered through a small silica gel plug, and concentrated *in vacuo* to afford the sub-title compound (6.1 g, 79%) as a
20 colourless oil that was used without further purification.

¹H NMR (300 MHz, CDCl₃) δ 7.11-7.14 (m, 2H), 6.84 (dt, *J* = 9 Hz, *J* = 2 Hz, 1H), 6.50 (t, *J*_{H-F} = 72 Hz, 1H).

25 (iv) 1-Fluoro-3-difluoromethoxy-5-vinylbenzene

Tri(butyl)vinylstannane (7.0 g, 22.2 mmol) was added to a suspension of 1-bromo-3-fluoro-5-difluoromethoxybenzene (4.9 g, 20.2 mmol; see step (iii) above), dichlorobis(triphenylphosphine)palladium(II) (1.42 g, 2.02 mmol) and anhydrous lithium chloride (0.90 g, 20.2 mmol) in THF (40 mL) under nitrogen at 65°C and the mixture was
30 stirred for 5 h. The reaction mixture was cooled to 0°C and 1N NaOH (90 mL) was added. The biphasic mixture was vigorously stirred for 1 h then the layers were separated. The aqueous layer was extracted with Et₂O (3 x 70 mL). The combined organic layers were

washed with 2N NaOH (2 x 40 mL) and H₂O (40 mL) then dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with hexane afforded the sub-title compound (2.2 g, 57%) as a colourless oil.

5 $R_f = 0.47$ (Hex)

¹H NMR (300 MHz, CDCl₃) δ 6.93-6.99 (m, 2H), 6.73-6.78 (m, 1H), 6.67 (dd, $J = 18$ Hz, $J = 11$ Hz, 1H), 6.51 (t, $J_{H-F} = 73$ Hz, 1H), 5.77 (d, $J = 18$ Hz, 1H), 5.36 (d, $J = 11$ Hz, 1H).

(v) Ph(3-F)(5-OCHF₂)-(R)CH(OH)CH₂OH

10 2-Methyl-2-propanol (140 mL), H₂O (140 mL), and AD-mix- β (39.2 g) were combined together and cooled to 0°C. 1-Fluoro-3-difluoromethoxy-5-vinylbenzene (5.0 g, 26.4 mmol; see step (iv) above) dissolved in a small amount of 2-methyl-2-propanol was added at once, and the heterogeneous slurry was vigorously stirred at 0°C until TLC revealed the absence of the starting material. The reaction was quenched at 0°C by addition of sodium
15 sulfite (42.0 g) and then warmed to room temperature and stirred for 60 min. The reaction mixture was extracted with Et₂O (3 x 120 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with CHCl₃:EtOAc (3:2) afforded the sub-title compound (5.8 g, 98%) as a colourless oil.

20 $R_f = 0.41$ (3:2 CHCl₃:EtOAc)

¹H NMR (300 MHz, CDCl₃) δ 6.96-6.99 (m, 2H), 6.77-6.82 (m, 1H), 6.51 (t, $J_{H-F} = 73$ Hz, 1H), 4.79-4.85 (m, 1H), 3.76-3.84 (m, 1H), 3.58-3.66 (m, 1H), 2.66 (d, $J = 3$ Hz, 1H), 2.00 (t, $J = 6$ Hz, 1H).

HPLC Analysis: 89.2%, >99% ee, ChiralPak AD Column (95:5 Hex:EtOH mobile phase).

(vi) Ph(3-F)(5-OCHF₂)-(R)CH(OH)CH₂OTBS

A solution of Ph(3-F)(5-OCHF₂)-(R)CH(OH)CH₂OH (5.5 g, 24.7 mmol; see step (v) above), 4-(dimethylamino)pyridine (121 mg, 1.0 mmol) and triethylamine (3.0 g, 29.6 mmol) in anhydrous CH₂Cl₂ (100 mL) was cooled to 0°C. A 1.0 M solution of *tert*-butyldimethylsilyl chloride in CH₂Cl₂ (26.0 mL, 26.0 mmol) was added dropwise, and the reaction mixture was allowed to warm to room temperature and stirred overnight. Saturated ammonium chloride solution (60 mL) was added, and the layers were separated. The organic layer was washed with saturated ammonium chloride solution (60 mL) and H₂O (2 x 35 mL) then dried (Na₂SO₄), filtered, and concentrated *in vacuo*. Flash chromatography on silica gel eluting with CHCl₃:Hex (3:1) afforded the sub-title compound (7.9 g, 85%) as a yellow oil.

R_f = 0.47 (3:1 CHCl₃:Hex)

¹H NMR (300 MHz, CDCl₃) δ 6.95-6.98 (m, 2H), 6.76-6.79 (m, 1H), 6.51 (t, J_{H-F} = 73 Hz, 1H), 4.71-4.74 (m, 1H), 3.75-3.80 (m, 1H), 3.48-3.54 (m, 1H), 2.99 (bs, 1H), 0.91 (s, 9H), 0.05 (s, 3H), 0.00 (s, 3H).

(vii) Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)CH₂OTBS

To a solution of Ph(3-F)(5-OCHF₂)-(R)CH(OH)CH₂OTBS (7.9 g, 0.51 mmol; see step (vi) above) and DIPEA (4.9 g, 48.1 mmol) in anhydrous CH₂Cl₂ (50 mL) at 0°C under nitrogen was added dropwise 2-methoxyethoxymethyl chloride (6.6 g, 48.1 mmol). The mixture was stirred for 24 h. Saturated ammonium chloride solution (70 mL) was added, and the layers were separated. The organic layer was washed with saturated ammonium chloride solution (70 mL) and H₂O (3 x 60 mL) then dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (8.8 g, 99%) as a yellow oil that was used without further purification.

R_f = 0.41 (4:1 CHCl₃:EtOAc)

¹H NMR (300 MHz, CDCl₃) δ 7.20 (s, 1H), 7.06 (s, 1H), 7.02 (s, 1H), 6.50 (t, J_{H-F} = 73 Hz, 1H), 4.79-4.81 (m, 1H), 4.66-4.68 (m, 2H), 3.47-3.82 (m, 6H), 3.36 (s, 3H), 0.85 (s, 9H), 0.01 (s, 3H), 0.00 (s, 3H).

(viii) Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)CH₂OH

To a solution of $\text{Ph(3-F)(5-OCHF}_2\text{)-(R)CH(OMEM)CH}_2\text{OTBS}$ (9.3 g, 21.9 mmol; see step (vii) above) in THF (60 mL) at room temperature was added a 1.0 M solution of tetrabutylammonium fluoride in THF (70.0 mL, 70.0 mmol) and the mixture was stirred overnight under nitrogen. The reaction was concentrated *in vacuo*. The yellow residue was dissolved in Et_2O (100 mL) and hexane (100 mL) and washed successively with saturated ammonium chloride solution (2 x 150 mL) and H_2O (3 x 70 mL). The organic layer was dried (Na_2SO_4), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (1:1) afforded the sub-title compound (4.2 g, 62%) as a yellow oil.

$R_f = 0.42$ (1:1 Hex:EtOAc)

$^1\text{H NMR}$ (300 MHz, CDCl_3) δ 6.91-6.95 (m, 2H), 6.75-6.81 (m, 1H), 6.51 (t, $J_{\text{H-F}} = 73$ Hz, 1H), 4.80-4.82 (m, 1H), 4.70-4.74 (m, 2H), 3.88-3.93 (m, 1H), 3.67-3.71 (m, 3H), 3.53-3.56 (m, 2H), 3.39 (s, 3H), 2.96-2.99 (m, 1H).

(ix) $\text{Ph(3-F)(5-OCHF}_2\text{)-(R)CH(OMEM)C(O)OH}$

A solution of $\text{Ph(3-F)(5-OCHF}_2\text{)-(R)CH(OMEM)CH}_2\text{OH}$ (4.2 g, 13.4 mmol; see step (viii) above) in acetone (100 mL) was added to an aqueous 5% NaHCO_3 solution (35 mL). This magnetically stirred heterogeneous mixture was cooled to 0°C and potassium bromide (159 mg, 1.3 mmol) and 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical (2.2 g, 14.1 mmol) were added. Sodium hypochlorite (5.25%, 30 mL) was then added dropwise over a period of 20 min while the mixture was vigorously stirred and maintained at 0°C . After 1 h, additional sodium hypochlorite (30 mL) and 5% NaHCO_3 solution (35 mL) were added and stirring was continued at 0°C for 2 h. The acetone was removed *in vacuo*. The aqueous layer was washed with Et_2O (4 x 40 mL). The aqueous layer was acidified to pH 3.5 with 10% citric acid and extracted with EtOAc (4 x 50 mL). The combined EtOAc extracts were successively washed with H_2O (4 x 30 mL) and brine (60 mL) then, dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (4.3 g, 98%) as a colourless oil which was used without further purification.

$R_f = 0.74$ (8.0:1.5:0.5 CHCl_3 :MeOH: Et_3N)

^1H NMR (300 MHz, acetone- d_6) δ 7.16-7.18 (m, 2H), 7.16 (t, $J_{\text{H-F}} = 89$ Hz, 1H), 7.00-7.03 (m, 1H), 5.30 (s, 1H), 4.88 (d, $J = 7$ Hz, 1H), 4.80 (d, $J = 7$ Hz, 1H), 3.54-3.75 (m, 2H), 3.46-3.49 (m, 2H), 3.28 (s, 3H).

5 (x) Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(Teoc)

To a solution of Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)C(O)OH (1.1 g, 3.4 mmol; see step (ix) above) in DMF (20 mL) under nitrogen at 0°C was added HAZE-Pab(Teoc)•HCl (2.0 g, 4.4 mmol), PyBOP (1.9 g, 3.7 mmol), and DIPEA (1.1 g, 8.4 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first
10 with CHCl₃:EtOH (15:1) and second with EtOAc:EtOH (20:1) to afford the sub-title compound (1.3 g, 56%) as a crushable white foam.

$R_f = 0.65$ (15:1 CHCl₃:EtOH)

15 ^1H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.80-7.84 (m, 2H), 7.40-7.46 (m, 2H), 6.95-7.16 (m, 3H), 6.92 and 6.88 (t, $J_{\text{H-F}} = 73$ Hz, 1H), 5.28 and 5.08 (s, 1H), 5.18-5.22 and 4.70-4.78 (m, 1H), 4.50-4.75 (m, 1H), 4.30-4.49 (m, 2H), 4.21-4.26 (m, 3H), 3.97-4.08 (m, 1H), 3.35-3.72 (m, 6H), 3.30 (s, 3H), 2.10-2.75 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

20

(xi) Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

A mixture of Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(Teoc) (590 mg, 0.87 mmol; see step (x) above) and carbon tetrabromide (287 mg, 0.87 mmol) in 2-propanol (20 mL) was refluxed for 1.5 h. The mixture was concentrated *in vacuo* then, partitioned with
25 H₂O (50 mL) and EtOAc (3 x 50 mL). The aqueous layer was extracted with additional EtOAc (2 x 10 mL). The combined organic extracts were washed with brine (30 mL) then dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with CHCl₃:EtOH (15:1) afforded the sub-title compound (60 mg, 12%) as a crushable white foam.

30

$R_f = 0.46$ (15:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.74 (d, *J* = 8 Hz, 2H), 7.35-7.37 (m, 2H), 6.97-7.07 (m, 2H), 6.80-6.84 (m, 1H), 6.82 and 6.80 (t, *J*_{H-F} = 73 Hz, 1H), 5.10 and 5.06 (s, 1H), 4.68-4.70 (m, 1H), 3.97-4.60 (m, 6H), 2.10-2.75 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

5 APCI-MS: (*M* + 1) = 595 *m/z*

(xii) Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.053 g, 0.089 mmol; see step (xi) above), was dissolved in 3 mL of TFA and allowed to react for 80 min while cooled on an ice bath. TFA was evaporated and the residue was freeze dried from water/acetonitrile to yield 0.042 g (80%) of the title compound as its TFA salt.

¹H-NMR (300 MHz; CD₃OD) rotamers: δ 7.7-7.6 (m, 2H), 7.5-7.4 (m, 2H), 7.1-6.6 (m, 4H), 5.2-5.0 (m, 1H plus minor rotamer of 1H), ca 4.8 (major rotamer of previous signal obscured by the CD₃OH signal), 4.6-4.3 (m, 2H), 4.26 (m, 1H, major rotamer), 4.10 (m, 1H, major rotamer), 3.96 (m, 1H, minor rotamer), 3.89 (m, 1H, minor rotamer), 2.60 (m, 1H, minor rotamer), 2.44 (m, 1H, major rotamer), 2.19 (m, 1H, major rotamer), 2.05 (m, 1H, minor rotamer).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons) δ 172.8, 172.0, 167.0.

20 ESI-MS+: (*M*+1) = 451 (*m/z*)

Example 26

Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)

25 (i) Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(OMe)

To a solution of Ph(3-F)(5-OCHF₂)-(R)CH(OMEM)C(O)OH (1.0 g, 3.1 mmol; see Example 25(ix) above) in DMF (30 mL) under nitrogen at 0°C was added HAze-Pab(OMe)•2HCl (1.4 g, 4.1 mmol), PyBOP (1.8 g, 3.4 mmol), and DIPEA (1.0 g, 7.8 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight.

30 The mixture was concentrated *in vacuo* and the residue chromatographed two times on silica gel, eluting first with CHCl₃:EtOH (15:1) and second with EtOAc to afford the subtitle compound (1.5 g, 79%) as a crushable white foam.

$R_f = 0.24$ (EtOAc)

^1H NMR (300 MHz, CD_3OD , complex mixture of rotamers) δ 7.58-7.62 (m, 2H), 7.32-7.38 (m, 2H), 7.03-7.16 (m, 3H), 6.92 and 6.88 (d, $J_{\text{H-F}} = 73$ Hz, 1H), 5.27 and 5.08 (s, 1H), 5.22-5.15 and 4.75-4.80 (m, 1H), 4.38-4.65 (m, 5H), 3.92-4.27 (m, 1H), 3.82 (s, 3H), 3.43-3.68 (m, 4H), 3.29 (s, 3H), 2.28-2.85 (m, 2H).

(ii) $\text{Ph}(3\text{-F})(5\text{-OCHF}_2)\text{-(R)CH(OH)C(O)-Aze-Pab(OMe)}$

A mixture of $\text{Ph}(3\text{-F})(5\text{-OCHF}_2)\text{-(R)CH(OMEM)C(O)-Aze-Pab(OMe)}$ (828 mg, 2.33 mmol; see step (i) above) and carbon tetrabromide (525 mg, 2.33 mmol) in 2-propanol (20 mL) was refluxed for 8 h and then stirred at room temperature overnight. The mixture was concentrated *in vacuo* and the residue partitioned with H_2O (70 mL) and EtOAc (50 mL). The aqueous layer was extracted with EtOAc (2 x 25 mL). The combined organic extracts were washed with brine (35 mL) then, dried (Na_2SO_4), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with $\text{CHCl}_3\text{:EtOH}$ (15:1) afforded the title compound (520 mg, 74%) as a crushable white foam.

Mp: 73-81°C

$R_f = 0.43$ (15:1 $\text{CHCl}_3\text{:EtOH}$)

^1H NMR (300 MHz, CD_3OD , complex mixture of rotamers) δ 7.59 (d, $J = 8$ Hz, 2H), 7.32-7.37 (m, 2H), 7.05-7.14 (m, 2H), 6.87-6.92 (m, 1H), 6.90 and 6.86 (t, $J_{\text{H-F}} = 73$ Hz, 1H), 5.13-5.18 and 4.75-4.85 (m, 2H), 4.15-4.45 (m, 4H), 3.81 (s, 3H), 2.10-2.75 (m, 2H).

^{13}C -NMR (100 MHz; CD_3OD): (carbonyl and/or amidine carbons) δ 172.0, 171.4, 153.9.

APCI-MS: $(M + 1) = 481$ m/z

Example 27Ph(3-Br)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab x TFA(i) 1,3-Dibromo-5-benzyloxybenzene

- 5 Sodium hydride (9.9 g, 0.414 mol, 95% dry) was added in portions to a stirred solution of benzyl alcohol (41.0 g, 0.394 mol) in THF (1.0 L) at room temperature under a nitrogen atmosphere and stirred for 1 h. To this solution was added dropwise 1,3-dibromo-5-fluorobenzene (100.0 g, 0.394 mol). After stirring overnight, the mixture was partitioned with H₂O (600 mL) and EtOAc (4 x 600 mL). The combined organic extracts were dried
10 (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with hexanes afforded the sub-title compound (101.3 g, 75%) as a yellow oil.

¹H NMR (300 MHz, CDCl₃) δ 7.30-7.48 (m, 5H), 7.18 (s, 1H), 7.06 (s, 2H), 4.99 (s, 2H).

15 (ii) 3,5-Dibromophenol

- Aluminium chloride (11.7 g, 87.6 mmol) was added in portions to a solution of 1,3-dibromo-5-benzyloxybenzene (10.0 g, 29.2 mmol; see step (i) above) and *N,N*-dimethylaniline (35.4 g, 292 mmol) in CH₂Cl₂ (100 mL) at room temperature under a nitrogen atmosphere. After 30 min, the mixture was partitioned with 1N HCl (300 mL)
20 and EtOAc (5 x 150 mL). The combined organic extracts were washed with saturated NaHCO₃ (150 mL) and brine (150 mL) then, dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (9:1) afforded the sub-title compound (6.1 g, 82%) as a white solid.

- 25 ¹H NMR (300 MHz, CDCl₃) δ 7.21 (s, 1H), 6.97 (s, 2H), 5.88 (bs, 1H).

(iii) 1,3-Dibromo-5-monofluoromethoxybenzene

To a tared, sealed 350 mL round-bottomed pressure flask containing a suspension of 3,5-dibromophenol (10.0 g, 39.7 mmol; see step (ii) above) and Cs_2CO_3 (20.7 g, 63.5 mmol) in DMF (150 mL) at -78°C was added chlorofluoromethane via bubbling for 5 min through the septum. The septum was replaced with a Teflon stopper and the flask was then sealed and allowed to warm to room temperature where the flask was weighed and determined to contain 9.0 g (131 mmol) of chlorofluoromethane. The solution was heated in an oil bath set at 70°C overnight. The flask was cooled to room temperature, the pressure cautiously released and the contents diluted with water (100 mL). The aqueous layer was extracted with Et_2O (3 x 200 mL) then, the combined organics were dried (Na_2SO_4), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with hexanes afforded the sub-title compound (7.9 g, 71%) as a white solid.

^1H NMR (300 MHz, CDCl_3) δ 7.40 (s, 1H), 7.18 (s, 2H), 5.67 (d, $J_{\text{H-F}} = 53$ Hz, 2H).

(iv) 1-Bromo-3-monofluoromethoxy-5-vinylbenzene

Tri(butyl)vinyltin (10.0 g, 31.4 mmol) was added dropwise to a solution of 1,3-dibromo-5-monofluoromethoxybenzene (8.5 g, 29.9 mmol; see step (iii) above), tetrakis(triphenylphosphine)palladium(0) (690 mg, 0.599 mmol), and 2,6-di-*tert*-butyl-4-methylphenol (spatula tip) in toluene (100 mL) under nitrogen. The mixture was stirred at 70°C for 8 h. The mixture was cooled to 0°C and 1N NaOH (70 mL) was added. After 1 h, the mixture was extracted with CH_2Cl_2 (3 x 300 mL) then, the combined organics were dried (Na_2SO_4), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with hexanes afforded the sub-title compound (4.3 g, 57%) as a colourless oil.

^1H NMR (300 MHz, CDCl_3) δ 7.30 (s, 1H), 7.16 (s, 1H), 7.01 (s, 1H), 6.60 (dd, $J = 6$ Hz, $J = 11$ Hz, 1H), 5.74 (d, $J = 16$ Hz, 1H), 5.67 (d, $J_{\text{H-F}} = 53$ Hz, 2H), 5.32 (d, $J = 8$ Hz, 1H).

(v) Ph(3-Br)(5-OCH₂F)-(R)CH(OH)CH₂OH

2-Methyl-2-propanol (100 mL), H_2O (100 mL), and AD-mix- β (27.5 g) were combined together and cooled to 0°C . 1-Bromo-3-monofluoromethoxy-5-vinylbenzene (4.3 g, 17.3 mmol; see step (iv) above) was added at once, and the heterogeneous slurry was vigorously stirred at 0°C until TLC revealed the absence of the starting material. The reaction was

quenched at 0°C by addition of saturated sodium sulfite (200 mL) and then warmed to room temperature and stirred for 60 min. The reaction mixture was extracted with EtOAc (3 x 150 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (4.9 g, 100%) as a colourless oil that was used without further purification.

¹H NMR (300 MHz, CD₃OD) δ 7.30 (s, 1H), 7.15 (s, 1H), 7.11 (s, 1H), 5.70 (d, *J*_{H-F} = 53 Hz, 2H), 4.62-4.70 (m, 1H), 3.52-3.70 (m, 2H).

HPLC Analysis: 92.1%, 96.9% ee, ChiralPak AD Column (95:5 Hex:EtOH mobile phase).

(vi) Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)CH₂OTBS

To a solution of Ph(3-Br)(5-OCH₂F)-(R)CH(OH)CH₂OH (4.9 g, 18.6 mmol; see step (v) above), 4-(dimethylamino)pyridine (453 mg, 3.71 mmol), and DIPEA (8.9 g, 93.0 mmol) in anhydrous CH₂Cl₂ (200 mL) was added dropwise a 1.0 M solution of *tert*-butyldimethylsilyl chloride in CH₂Cl₂ (22.3 mL, 22.3 mmol). The reaction mixture was stirred 10 h at room temperature. To the mixture was added DIPEA (8.9 g, 93.0 mmol) and 2-methoxyethoxymethyl chloride (13.9 g, 111 mmol) dropwise. After 16 h, additional 2-methoxyethoxymethyl chloride (2.2 g) was added and the reaction stirred overnight. The mixture was diluted with H₂O (100 mL), and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (3 x 200 mL) then, the combined organic layers were dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (5:1) afforded the sub-title compound (4.8 g, 55%) as a colourless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.29 (s, 1H), 7.22 (s, 1H), 7.05 (s, 1H), 5.74 (d, *J*_{H-F} = 53 Hz, 2H), 4.84 (d, *J* = 7 Hz, 1H), 4.70-4.74 (m, 2H), 3.50-3.91 (m, 6H), 3.42 (s, 3H), 0.90 (s, 9H), 0.05 (s, 3H), 0.01 (s, 3H).

(vii) Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)CH₂OH

To a solution of Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)CH₂OTBS (4.7 g, 10.1 mmol; see step (vi) above) in THF (100 mL) was added a 1.0 M solution of tetrabutylammonium fluoride in THF (13.1 mL, 13.1 mmol) at room temperature and the mixture was stirred 1 h. The mixture was partitioned with H₂O (100 mL) and EtOAc (3 x 100 mL) then, the

combined organics were dried (Na_2SO_4), filtered and concentrated *in vacuo* to afford the sub-title compound (3.3 g, 92%) as a colourless oil that was used without further purification.

- 5 ^1H NMR (300 MHz, CD_3OD) δ 7.22 (s, 1H), 7.14 (s, 1H), 7.03 (s, 1H), 5.71 (d, $J_{\text{H-F}} = 53$ Hz, 2H), 4.80-4.82 (m, 1H), 4.58-4.66 (m, 2H), 3.71-3.77 (m, 1H), 3.39-3.65 (m, 5H), 3.27 (s, 3H).

100711

(viii) Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)C(O)OH

A solution of Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)CH₂OH (2.1 g, 6.0 mmol; see step (vii) above) in acetone (40 mL) was added to an aqueous 5% NaHCO₃ solution (15 mL). This magnetically stirred heterogeneous mixture was cooled to 0°C and potassium bromide (70 mg, 0.60 mmol) and 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical (976 mg, 5.8 mmol) were added. Sodium hypochlorite (5.25%, 15 mL) was then added dropwise over a period of 10 min while the mixture was vigorously stirred and maintained at 0°C. After 1 h, additional sodium hypochlorite (10 mL) and NaHCO₃ solution (20 mL) were added and stirring was continued at 0°C for an additional 4 h. The acetone was removed on a rotary evaporator. The aqueous layer was diluted with 10% NaHCO₃ solution (30 mL) and was washed with Et₂O (3 x 20 mL). The aqueous layer was acidified to pH 3.5 with 10% citric acid and extracted with EtOAc (3 x 40 mL). The combined EtOAc extracts were washed with H₂O (3 x 50 mL) and brine (50 mL) then, dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (1.7 g, 78%) as a colourless oil which was used without further purification.

¹H NMR (300 MHz, CD₃OD) δ 7.38 (s, 1H), 7.25 (s, 1H), 7.18 (s, 1H), 5.76 (d, *J*_{H-F} = 53 Hz, 2H), 5.21 (s, 1H), 4.83 (d, *J* = 7 Hz, 1H), 4.75 (d, *J* = 7 Hz, 1H), 3.62-3.78 (m, 2H), 3.48-3.52 (m, 2H), 3.32 (s, 3H).

(ix) Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)C(O)-Aze-Pab(Teoc)

To a solution of Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)C(O)OH (1.0 g, 2.72 mmol; see step (viii) above) in DMF (20 mL) under nitrogen at 0°C was added Haze-Pab(Teoc)•HCl (1.6 g, 3.5 mmol), PyBOP (1.6 g, 3.0 mmol), and DIPEA (880 mg, 6.81 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica gel, eluting first with CHCl₃:EtOH (15:1) and second with EtOAc:EtOH (20:1) to afford the sub-title compound (1.2 g, 62%) as a crushable white foam.

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.80-7.84 (m, 2H), 7.40-7.46 (m, 2H), 7.13-7.32 (m, 3H), 5.84-5.87 (m, 1H), 5.67-5.69 (m, 1H), 5.25 and 5.07 (s,

1H), 5.18-5.23 and 4.80-4.88 (m, 1H), 3.97-4.79 (m, 8H), 3.60-3.71 (m, 2H), 3.40-3.53 (m, 2H), 3.32 (s, 3H), 2.10-2.75 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

(x) Ph(3-Br)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

- 5 A mixture of Ph(3-Br)(5-OCH₂F)-(R)CH(OMEM)C(O)-Aze-Pab(Teoc) (347 mg, 0.478 mmol; see step (ix) above) and carbon tetrabromide (159 mg, 0.478 mmol) in 2-propanol (10 mL) was refluxed for 1.5 h. The mixture was concentrated *in vacuo* then partitioned with H₂O (20 mL) and EtOAc (3 x 30 mL). The combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with
10 CHCl₃:EtOH (15:1) afforded the sub-title compound (59 mg, 19%) as a crushable white foam.

Mp: 81-87°C

R_f = 0.58 (9:1 CHCl₃:EtOH)

- 15 ¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.84 (d, *J* = 8 Hz, 2H), 7.40-7.48 (m, 2H) 7.18-7.30 (m, 3H), 5.80 (d, *J*_{H-F} = 53 Hz, 2H), 5.21 and 5.15 (s, 1H), 5.18-5.24 and 4.80-4.88 (m, 1H), 3.98-4.54 (m, 6H), 2.10-2.70 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

APCI-MS: (M + 1) = 637 m/z

(xi) Ph(3-Br)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-Br)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.073 g, 0.11 mmol; see step (x) above), was dissolved in 5 mL of TFA and allowed to react for 90 min while being cooled on an ice bath. TFA was evaporated and the residue purified by prep RPLC with
5 CH₃CN:0.1M NH₄OAc (30:70). The pertinent fractions were evaporated and freeze dried from water/acetonitrile to yield 49 mg (77%) of the title compound as its acetate salt.

¹H-NMR (300 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.54 (m, 2H), 7.37 (s, 1H, major rotamer), 7.33 (s, 1H, minor rotamer), 7.25-7.1 (m, 2H), 5.75 (d, 2H), 5.22 (m, 1H, minor rotamer), 5.18 (s, 1H, major rotamer), 5.11 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.6-4.4 (m, 2H), 4.37 (m, 1H, major rotamer), 4.16 (m, 1H, major rotamer), 4.1-3.9 (m, 2H, two signals from minor rotamer), 2.70 (m, 1H, minor rotamer), 2.52 (m, 1H, major rotamer), 2.30 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer), 1.89 (s, 3H).

15 ESI-MS+: (M+1) = 493/495 (m/z)

Example 28

Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA

20 (i) 1,3-Dibromo-5-difluoromethoxybenzene

To a tared, sealed 350 mL round-bottomed pressure flask containing a solution of 3,5-dibromophenol (10.0 g, 39.7 mmol; see Example 27(ii) above) in 2-propanol (100 mL) and 30% KOH (80 mL) at -78°C was added chlorodifluoromethane *via* bubbling for 15 min through the septum. The septum was replaced with a Teflon stopper and the flask was then
25 sealed and allowed to warm to room temperature where the flask was weighed and determined to contain 12.0 g (138 mmol) of chlorodifluoromethane. The solution was refluxed overnight in an oil bath set at 80°C. The flask was cooled to room temperature, the pressure cautiously released and the contents diluted with H₂O (200 mL). The aqueous layer was extracted with CHCl₃ (2 x 150 mL), then the combined organics were dried
30 (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by Kugelrohr distillation at 80°C at 0.2 mm Hg to afford the sub-title compound (9.6 g, 80%) as clear liquid.

¹H NMR (300 MHz, CDCl₃) δ 7.55 (s, 1H), 7.26 (s, 2H), 6.52 (t, *J*_{H-F} = 68 Hz, 1H).

(ii) 1-Bromo-3-difluoromethoxy-5-vinylbenzene

5 Tri(ethyl)vinyltin (10.5 g, 33.1 mmol) was added dropwise to a solution of 1,3-dibromo-5-difluoromethoxybenzene (9.1 g, 30.1 mmol; see step (i) above), tetrakis(triphenylphosphine)palladium(0) (700 mg, 0.60 mmol), and 2,6-di-*tert*-butyl-4-methylphenol (spatula tip) in toluene (125 mL) under nitrogen. The mixture was stirred at 50°C overnight. The mixture was cooled to 0°C and 1N NaOH (70 mL) was added. After
10 1 h, the mixture was extracted with CH₂Cl₂ (3 x 300 mL) then, the combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with hexanes afforded the sub-title compound (5.1 g, 68%) as a colourless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.53 (s, 1H), 7.18 (s, 1H), 7.08 (s, 1H), 6.60 (dd, *J* = 6 Hz, *J* = 11 Hz, 1H), 6.57 (t, *J*_{H-F} = 68 Hz, 1H), 5.77 (d, *J* = 11 Hz, 1H), 5.36 (d, *J* = 8 Hz, 1H).
15

(iii) Ph(3-Br)(5-OCF₂)-(R)CH(OH)CH₂OH

2-Methyl-2-propanol (150 mL), H₂O (150 mL), and AD-mix-β (27.8 g) were combined together and cooled to 0°C. 1-Bromo-3-difluoromethoxy-5-vinylbenzene (4.6 g, 18.6
20 mmol; see step (ii) above) was added at once, and the heterogeneous slurry was vigorously stirred at 0°C until TLC indicated the absence of the starting material, then the solution was warmed to room temperature and stirred overnight. The reaction was quenched at 0°C by addition of saturated sodium sulfite (300 mL) and then warmed to room temperature and stirred for 60 min. The reaction mixture was extracted with EtOAc (3 x 200 mL). The
25 combined organic extracts were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (5.0 g, 95%) as a colourless oil that was used without further purification.

¹H NMR (300 MHz, CD₃OD) δ 7.43 (s, 1H), 7.23 (s, 1H), 7.16 (s, 1H), 6.86 (t, *J*_{H-F} = 75
30 Hz, 1H), 4.64-4.67 (m, 1H), 3.54-3.59 (m, 2H).

HPLC Analysis: 88.6%, 96.3% ee, ChiralPak AD Column (95:5 Hex:EtOH mobile phase).

(iv) Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)CH₂OTBS

To a solution of Ph(3-Br)(5-OCHF₂)-(R)CH(OH)CH₂OH (4.9 g, 17.3 mmol; see step (iii) above), 4-(dimethylamino)pyridine (420 mg, 3.5 mmol), and DIPEA (11.2 g, 86.3 mmol) in anhydrous CH₂Cl₂ (250 mL) was added dropwise a 1.0 M solution of *tert*-butyldimethylsilyl chloride in CH₂Cl₂ (20.7 mL, 20.7 mmol). The reaction mixture was stirred overnight at room temperature. To the mixture was added DIPEA (11.2 g, 86.3 mmol) and 2-methoxyethoxymethyl chloride (12.9 g, 104 mmol) dropwise. After 3 d, additional 2-methoxyethoxymethyl chloride (3.3 g) was added and the reaction stirred overnight. The mixture was diluted with water (250 mL), and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (2 x 250 mL), then the combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with Hex:EtOAc (4:1) afforded the sub-title compound (4.3 g, 51%) as a colourless oil.

¹H NMR (300 MHz, CDCl₃) δ 7.40 (s, 1H), 7.25 (s, 1H), 7.08 (s, 1H), 6.58 (t, *J*_{H-F} = 75 Hz, 1H), 4.84 (d, *J* = 7 Hz, 1H), 4.70-4.74 (m, 2H), 3.50-3.91 (m, 6H), 3.42 (s, 3H), 0.90 (s, 9H), 0.12 (s, 3H), 0.05 (s, 3H).

(v) Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)CH₂OH

To a solution of Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)CH₂OTBS (3.3 g, 6.9 mmol; see step (iv) above) in THF (60 mL) was added a 1.0 M solution of tetrabutylammonium fluoride in THF (9.0 mL, 9.0 mmol) at room temperature. The reaction was stirred for 45 min, then the mixture was partitioned with water (150 mL) and EtOAc (2 x 120 mL). The combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (2.5 g, 98%) as a yellow oil that was used without further purification.

¹H NMR (300 MHz, CD₃OD) δ 7.35 (s, 1H), 7.21 (s, 1H), 7.08 (s, 1H), 6.83 (t, *J*_{H-F} = 73 Hz, 1H), 4.73 (d, *J* = 7 Hz, 1H), 4.59-4.68 (m, 2H), 3.40-3.80 (m, 6H), 3.26 (s, 3H).

(vi) Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)OH

A solution of Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)CH₂OH (3.0 g, 8.1 mmol; see step (v) above) in acetone (60 mL) was added to an aqueous 5% NaHCO₃ solution (25 mL). This

magnetically stirred heterogeneous mixture was cooled to 0°C then potassium bromide (100 mg, 0.81 mmol) and 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical (1.3 g, 8.5 mmol) were added. Sodium hypochlorite (5.25%, 19 mL) was then added dropwise over a period of 10 min while the mixture was vigorously stirred and maintained at 0°C. After 1 h, additional sodium hypochlorite (17 mL) and NaHCO₃ solution (34 mL) were added and stirring was continued at 0°C for an additional 4 h. The acetone was removed on a rotary evaporator. The aqueous layer was diluted with 10% NaHCO₃ solution (30 mL) and was washed with Et₂O (3 x 20 mL). The aqueous layer was acidified to pH 3.5 with 10% citric acid and extracted with EtOAc (3 x 40 mL). The combined EtOAc layers were washed with H₂O (3 x 50 mL) and brine (50 mL), and then dried (Na₂SO₄), filtered and concentrated *in vacuo* to afford the sub-title compound (2.1 g, 66%) as a colourless oil which was used without further purification.

¹H NMR (300 MHz, CD₃OD) δ 7.51 (s, 1H), 7.32 (s, 1H), 7.24 (s, 1H), 6.88 (t, *J*_{H-F} = 73 Hz, 1H), 5.21 (s, 1H), 4.84 (d, *J* = 7 Hz, 1H), 4.76 (d, *J* = 7 Hz, 1H), 3.62-3.80 (m, 2H), 3.48-3.52 (m, 2H), 3.32 (s, 3H).

(vii) Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(Teoc)

To a solution of Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)OH (1.0 g, 2.62 mmol; see step (vi) above) in DMF (50 mL) under nitrogen at 0°C was added HAze-Pab(Teoc)•HCl (1.5 g, 3.38 mmol), PyBOP (1.5 g, 2.9 mmol), and DIPEA (840 mg, 6.50 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed on silica gel eluting with CHCl₃:EtOH (15:1) to afford the sub-title compound (1.1 g, 59%) as a crushable white foam.

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.79-7.83 (m, 2H), 7.26-7.52 (m, 5H), 6.94 and 6.91 (t, *J*_{H-F} = 73 Hz, 1H), 5.27 and 5.07 (s, 1H), 5.20-5.23 and 4.80-4.88 (m, 1H), 4.01-4.79 (m, 8H), 3.60-3.71 (m, 2H), 3.40-3.53 (m, 2H), 3.32 (s, 3H), 2.10-2.75 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

(viii) Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc)

A mixture of Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(Teoc) (369 mg, 0.496 mmol; see step (vii) above) and carbon tetrabromide (165 mg, 0.496 mmol) in 2-propanol (10 mL) was refluxed for 12 h. The mixture was concentrated *in vacuo*, then partitioned with H₂O (15 mL) and EtOAc (5 x 20 mL). The combined organics were dried (Na₂SO₄),
5 filtered and concentrated *in vacuo*. Flash chromatography on silica gel eluting with CHCl₃:EtOH (15:1) afforded the sub-title compound (134 mg, 41%) as a crushable white foam.

Mp: 92-98°C

10 R_f = 0.37 (9:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.80-7.86 (m, 2H), 7.40-7.48 (m, 2H) 7.10-7.33 (m, 3H), 6.92 and 6.88 (t, J_{H-F} = 73 Hz, 1H), 5.18 and 5.11 (s, 1H), 5.18-5.24 and 4.76-4.80 (m, 1H), 3.98-4.54 (m, 6H), 2.10-2.70 (m, 2H), 1.05-1.11 (m, 2H), 0.08 (s, 9H).

15 APCI-MS: (M + 1) = 655 m/z

(ix) Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab x TFA

Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (0.081 g, 0.124 mmol; see step (viii) above), was dissolved in 5 mL of TFA and allowed to react for 80 min while being cooled
20 on an ice bath. TFA was evaporated and the residue purified by prep RPLC with CH₃CN:0.1M NH₄OAc (30:70). The pertinent fractions were evaporated and freeze dried from water/acetonitrile to yield 59 mg (83%) of the title compound as its acetate salt.

¹H-NMR (300 MHz; CD₃OD) rotamers: δ 7.8-7.7 (m, 2H), 7.6-7.4 (m, 3H), 7.3-7.2 (m, 2H), 6.89 (t, 1H, major rotamer), 6.87 (t, 1H, minor rotamer), 5.23 (m, 1H, minor rotamer), 5.21 (s, 1H, major rotamer), 5.13 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.6-4.4 (m, 2H), 4.38 (m, 1H, major rotamer), 4.20 (m, 1H, major rotamer), 4.1-3.9 (m, 2H, two signals from minor rotamer), 2.70 (m, 1H, minor rotamer), 2.54 (m, 1H, major rotamer), 2.29 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer), 1.89 (s, 3H).

30 ¹³C-NMR (75 MHz; CD₃OD): (carbonyl and/or amidine carbons) δ 172.0, 171.7, 167.0.

MS (m/z) 511/513 (M+1)⁺

Example 29Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)(i) Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(OMe)

5 To a solution of Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)OH (957 mg, 2.48 mmol; see Example 28(vi) above) in DMF (30 mL) under nitrogen at 0°C was added Haze-Pab(OMe)•2HCl (1.1 g, 3.2 mmol), PyBOP (1.4 g, 2.7 mmol), and DIPEA (804 mg, 6.2 mmol). The reaction was stirred at 0°C for 2 h and then at room temperature overnight. The mixture was concentrated *in vacuo* and the residue chromatographed twice on silica
10 gel, eluting first with CHCl₃:EtOH (9:1) and second with EtOAc:EtOH (15:1) to afford the sub-title compound (1.1 g, 72%) as a crushable white foam.

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.59-7.65 (m, 2H), 7.20-7.55 (m, 5H), 6.95 and 6.91 (t, *J*_{H-F} = 73 Hz, 1H), 5.27 and 5.07 (s, 1H), 5.18-5.23 and
15 4.75-4.84 (m, 1H), 3.87-4.89 (m, 6H), 3.84 (s, 3H), 3.60-3.71 (m, 2H), 3.40-3.53 (m, 2H), 3.32 (s, 3H), 2.10-2.75 (m, 2H).

(ii) Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe)

A mixture of Ph(3-Br)(5-OCHF₂)-(R)CH(OMEM)C(O)-Aze-Pab(OMe) (1.1 g, 1.8 mmol; see step (i) above) and carbon tetrabromide (583 mg, 1.8 mmol) in 2-propanol (30 mL)
20 was refluxed for 2.5 d. During this time, additional carbon tetrabromide (5 portions of 50 mg at intervals for an additional 0.90 mmol) was added to ensure completion of the reaction. The mixture was concentrated *in vacuo*, then partitioned with H₂O (50 mL) and EtOAc (5 x 25 mL). The combined organics were dried (Na₂SO₄), filtered and
25 concentrated *in vacuo*. Flash chromatography on silica gel eluting with CHCl₃:EtOH (15:1) afforded the title compound (460 mg, 50%) as a crushable white foam.

Mp: 71-75°C

R_f = 0.63 (9:1 CHCl₃:EtOH)

¹H NMR (300 MHz, CD₃OD, complex mixture of rotamers) δ 7.59 (d, *J* = 8 Hz, 2H), 7.20-7.54 (m, 5H), 6.90 and 6.87 (t, *J*_{H-F} = 73 Hz, 1H), 5.18 and 5.11 (s, 1H), 4.76-4.80 (m, 1H),
30 3.98-4.54 (m, 4H), 3.82 (s, 3H), 2.10-2.70 (m, 2H).

^{13}C -NMR (100 MHz; CD_3OD): (carbonyl and/or amidine carbons, rotamers) δ 172.5, 172.1, 171.6, 154.1.

APCI-MS: $(M + 1) = 542$ m/z

5 Example 30

$\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R)CH(OH)C(O)-Aze-Pab(OH)}$

(i) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R)CH(OH)C(O)-Aze-Pab(Z)}$

Boc-Aze-Pab(Z) (see international patent application WO 97/02284, 92 mg, 0.197 mmol)
10 was dissolved in 10 mL of EtOAc saturated with HCl(g) and allowed to react for 10 min. The solvent was evaporated and the residue was mixed with $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R)CH(OH)C(O)OH}$ (50 mg, 0.188 mmol; see Example 17(v) above), PyBOP (109 mg, 0.209 mmol) and finally diisopropylethyl amine (96 mg, 0.75 mmol) in 2 mL of DMF. The mixture was stirred for 2 h and then poured into 50 mL of water and extracted three
15 times with EtOAc. The combined organic phase was washed with water, dried (Na_2SO_4) and evaporated. The crude product was flash chromatographed on silica gel with EtOAc:MeOH (9:1). Yield: 100 mg (87%).

^1H NMR (300 MHz, CD_3OD , mixture of rotamers) δ 7.85-7.75 (m, 2H), 7.45-7.25 (m, 7H),
20 7.11 (m, 1H, major rotamer), 7.08 (m, 1H, minor rotamer), 7.05-6.9 (m, 2H), 6.13 (bt, 1H), 5.25-5.05 (m, 3H), 4.77 (m, 1H, partially hidden by the CD_3OH signal), 4.5-3.9 (m, 7H), 2.64 (m, 1H, minor rotamer), 2.47 (m, 1H, major rotamer), 2.25 (m, 1H, major rotamer), 2.13 (m, 1H, minor rotamer)

25 (ii) $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R)CH(OH)C(O)-Aze-Pab(OH)}$

Hydroxylamine hydrochloride (65 mg, 0.94 mmol) and triethylamine (0.319 g, 3.16 mmol) were mixed in 8 mL of THF and sonicated for 1 h at 40°C . $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CHF}_2)\text{-(R)CH(OH)C(O)-Aze-Pab(Z)}$ (96 mg, 0.156 mmol; see step (i) above) was added with 8 mL more of THF. The mixture was stirred at 40°C for 4.5 days. The solvent was
30 evaporated and the crude product was purified by preparative RPLC with CH_3CN :0.1M NH_4OAc (40:60). Yield: 30 mg (38%). Purity: 99%.

¹H NMR (300 MHz, CD₃OD, mixture of rotamers) δ 7.6-7.55 (m, 2H), 7.35-7.3 (m, 2H), 7.12 (m, 1H, major rotamer), 7.09 (m, 1H, minor rotamer), 7.05-6.9 (m, 2H), 6.15 (triplet of multiplets, 1H), 5.15 (m, 1H, minor rotamer), 5.13 (s, 1H, major rotamer), 5.08 (s, 1H, minor rotamer), 4.77 (m, 1H, major rotamer), 4.5-4.2 (m, 5H), 4.08 (m, 1H, major rotamer), 3.97 (m, 1H, minor rotamer), 2.66 (m, 1H, minor rotamer), 2.50 (m, 1H major rotamer), 2.27 (m, 1H, major rotamer), 2.14 (m, 1H, minor rotamer).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 172.8, 172.2, 171.4, 159.1, 158.9, 154.2.

APCI-MS: (M + 1) = 497/499 m/z

10

Example 31

Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OH)

(i) Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(Z)

15 Boc-Aze-Pab(Z) (130 mg, 0.279 mmol) was dissolved in 15 mL of EtOAc saturated with HCl(g) and allowed to react for 10 min. The solvent was evaporated and the residue was mixed with Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)-C(O)OH (63 mg, 0.188 mmol; see Example 21(v) above) in 3 mL of DMF, PyBOP (147 mg, 0.279 mmol) and finally diisopropylethyl amine (134 mg, 1.03 mmol). The mixture was stirred for 130 min and
20 then poured into 75 mL of water and extracted three times with EtOAc. The combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. The crude product was flash chromatographed on silica gel with EtOAc/MeOH = 95/5. Yield: 119 mg (79%).

25 ¹H NMR (400 MHz, CDCl₃) δ 8.06 (bt, 1H), 7.67 (d, 2H), 7.45-7.25 (m, 5H), 7.18 (d, 2H), 6.89 (m, 1H), 6.84 (m, 1H), 6.76 (m, 1H), 5.16 (s, 2H), 4.84 (s, 1H), 4.79 (m, 1H), 4.66 (doublet of multiplets, 2H), 4.4-4.3 (m, 2H), 4.10 (doublet of multiplets, 2H), 4.02 (m, 1H), 3.67 (m, 1H), 2.46 (m, 1H), 2.28 (m, 1H).

30 (ii) Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OH)

Hydroxylamine hydrochloride (80 mg, 1.16 mmol) and triethylamine (0.392 g, 3.87 mmol) were mixed in 9 mL of THF and sonicated for 1 h at 40°C. Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(Z) (96 mg, 0.156 mmol; see step (i) above) was added with 9 mL more of THF. The mixture was stirred at 40°C for 48 h and 3 days at room temperature. The solvent was evaporated and the crude product was purified by preparative RPLC with CH₃CN:0.1M NH₄OAc (30:70). Yield: 72 mg (78%). Purity: 100%.

¹H NMR (400 MHz, CD₃OD, mixture of rotamers) δ 7.6-7.55 (m, 2H), 7.35-7.25 (m, 4H), 7.07 (m, 1H, major rotamer), 7.04 (m, 1H, minor rotamer), 7.0-6.9 (M, 2h), 5.12 (m, 1H, minor rotamer), 5.08 (s, 1H, minor rotamer), 5.04 (s, 1H), 4.78 (m, 1H, major rotamer), 4.68 (doublet of multiplets, 2 H), 4.5-4.25 (m, 3H), 4.20 (doublet of multiplets, 2H) 4.06 (m, 1H, major rotamer), 3.97 (m, 1H, minor rotamer), 2.65 (m, 1H, minor rotamer), 2.48 (m, 1H major rotamer), 2.27 (m, 1H, major rotamer), 2.14 (m, 1H, minor rotamer)

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 172.3, 171.5, 159.8, 154.3

APCI-MS: (M + 1) = 479/481 m/z

Example 32

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Pro-Pab

(i) Boc-Pro-Pab(Teoc)

Boc-Pro-Pab(Z) (see international patent application WO 97/02284, 15.0 g, 0.0321 mol) was dissolved in 150 mL of ethanol and 200 mg 10% Pd/C (50% moisture) was added. The mixture was stirred and hydrogenated at atmospheric pressure for 2 h, filtered through Hyflo and concentrated. The product was used without further purification. Of this product was taken 10 g (0.029 mol), which was dissolved in 300 mL of THF. Teoc-*p*-nitrophenyl carbonate (10 g, 0.035 mol) was added. A solution of potassium carbonate (5.2 g, 0.038 mol) in 50 mL of water was added over 3 min and the resulting solution was stirred for 3 days, concentrated and the remainder was extracted with EtOAc three times. The combined organic layer was washed with water, dried (Na₂SO₄) and evaporated. The crude product was flash chromatographed on silica gel using methylene chloride:acetone (4:1). Yield: 9.8 g (69%).

(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Pro-Pab(Teoc)

Boc-Pro-Pab(Teoc) (107 mg, 0.218 mmol; see step (i) above) was dissolved in 10 mL of EtOAc saturated with HCl(g) and allowed to react for 10 min. The solvent was evaporated and the residue was mixed with Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (50 mg, 0.198 mmol; see Example 1(viii) above) in 3 mL of DMF, PyBOP (115 mg, 0.218 mmol) and finally diisopropylethyl amine (104 mg, 0.80 mmol). The mixture was stirred for 2 h and then poured into 75 mL of water and extracted three times with EtOAc. The combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. The crude product was flash chromatographed on silica gel with EtOAc:MeOH (95:5). Yield: 89 mg (72%).

¹H NMR (400 MHz, CDCl₃) δ 7.54 (bt, 1H), 7.47 (d, 2H), 7.12 (m, 1H), 7.08 (d, 2H), 7.02 (m, 1H), 6.95 (m, 1H), 6.50 (t, 1H), 5.21 (s, 1H), 4.42 (m, 1H), 4.35-4.15 (m, 3H), 3.59 (m, 1H), 2.94 (m, 1H), 2.1-1.7 (m, 4H), 1.06 (m, 2H), 0.04 (s, 9H).

(iii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Pro-Pab x TFA

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Pro-Pab(Teoc) (85 mg, 0.136 mmol; see step (ii) above) was dissolved in 1 mL of methylene chloride and cooled on an ice bath. TFA (4 mL) was added and the reaction was stirred for 90 min. The TFA was evaporated and the residue was freeze-dried from water and acetonitrile. Yield: 79 mg (92%). Purity: 94%.

¹H NMR (400 MHz, CD₃OD, mixture of rotamers) δ 7.85-7.7 (m, 2H), 7.58 (d, 2H, major rotamer), 7.47 (d, 2H, minor rotamer), 7.35 (m, 1H, major rotamer), 7.27 (m, 1H, minor rotamer), 7.2-7.1 (m, 2H), 6.88 (t, 1H), 5.38 (s, 1H, major rotamer), 5.22 (s, 1H, minor rotamer), 4.58 (d, 1H), 4.5-4.2 (m, 2H), 3.8-3.5 (m, 1H), 3.35 (m, 1H), 2.2-1.8 (m, 4H).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons) δ 173.6, 171.1, 167.0.

APCI-MS: (M + 1) = 481/483 m/z

Example 33

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Pro-Pab(OMe)

(i) 4-Azidomethyl-N-methoxy-benzamidine

4-Azidomethylbenzonitrile (17.3 g, 0.109 mol; Nishiyama *et al*; *Chem. Lett.* (1982) 1477) was dissolved in 500 mL of toluene and 200 mL of absolute ethanol. The solution was cooled to -10°C and HCl(g) was bubbled through until saturation. The mixture was kept in the refrigerator for 2 days when most of the solvents were evaporated. Diethyl ether was added and was decanted off. The product was re-dissolved in a solution of O-methylhydroxyl amine (10.5 g, 0.125 mol) and triethyl amine (56 mL) in 200 mL of methanol. The mixture was allowed to stand for 3 days whence the methanol was evaporated with addition of EtOAc. The organic phase was washed with water, dilute HOAc and aqueous sodium bicarbonate, dried (Na₂SO₄) and diluted with more EtOAc to a total volume of 500 mL. A sample of 25 mL was evaporated to dryness. The remainder was 932 mg. Total yield: 18.6 g (83%).

(ii) 4-Aminomethyl-N-methoxy-benzamidine

To a solution of 4-azidomethyl-N-methoxy-benzamidine (11.3 g, 0.055 mol; see step (i) above) in 200 mL of ethanol was added 200 mg of PtO₂. The mixture was hydrogenated with constant bubbling of hydrogen for 4 h and subsequently filtered through Celite® and evaporated. Yield: 7.34 g (74%).

(iii) Boc-Pro-Pab(OMe)

To a suspension of Boc-Pro-OH (9.7 g, 0.045 mol), 4-aminomethyl-N-methoxy-benzamidine (7.34 g, 0.041 mol; see step (ii) above) and dimethylaminopyridine (7.8 g, 0.064 mol) in 300 mL of acetonitrile was added EDC base (11.7 mL, 0.068 mol). The mixture was stirred for 18 h, concentrated and partitioned between water and EtOAc. The organic layer was washed with water, aqueous sodium bicarbonate, dried (MgSO₄) and evaporated. The crude product was flash chromatographed on silica gel with EtOAc. Yield: 9.73 g (63%).

(iv) H-Pro-Pab(OMe) x 2 HCl

Boc-Pro-Pab(OMe) (9.7 g, 0.026 mol; see step (iii) above) was dissolved in 250 mL of EtOAc. The ice cooled solution was saturated with HCl(g) by bubbling for 5 min. The

product precipitated immediately and 125 mL of absolute ethanol was added. The mixture was sonicated until most of the material had solidified. Diethyl ether (200 mL) was added and the suspension was filtered. A few lumps that had not solidified were again treated with absolute ethanol and diethyl ether. The solid was dried. Yield: 7.57 g (86%).

5

^1H NMR (400 MHz, CD_3OD) δ 7.74 (d, 2H), 7.58 (d, 2H), 4.55 (s, 2H), 4.38 (m, 1H), 3.98 (s, 3H), 3.45-3.3 (m, 2H), 2.50 (m, 1H), 2.15-2.0 (m, 3H)

(v) $\text{Ph}(3\text{-Cl})(5\text{-OCHF}_2)\text{-(R)CH(OH)C(O)-Pro-Pab(OMe)}$

10 $\text{Ph}(3\text{-Cl})(5\text{-OCHF}_2)\text{-(R)CH(OH)C(O)OH}$ (50 mg, 0.198 mmol; see Example 1(viii) above), H-Pro-Pab(OMe) (76 mg, 0.218 mmol, see step (iv) above) and PyBOP (115 mg, 0.218 mmol) were dissolved in 2 mL of DMF. Diisopropylethyl amine (104 mg, 0.80 mmol) was added and the mixture was stirred for 2.5 h. The mixture was poured into 50 mL of water and extracted three times with EtOAc and the combined organic phase was
15 washed with brine, dried (Na_2SO_4) and evaporated. The residue was flash chromatographed on silica gel with EtOAc:MeOH (95:5). Yield: 37 mg (36%). Purity: 98%.


^1H NMR (400 MHz, CD_3OD , mixture of rotamers) δ 7.60 (d, 2H, major rotamer), 7.57 (d, 20 2H, minor rotamer), 7.4-7.1 (m, 5H), 6.89 (t, 1H, major rotamer), 6.87 (t, 1H, minor rotamer), 5.35 (s, 1H, major rotamer), 5.21 (s, 1H, minor rotamer), 4.72 (m, 1H, minor rotamer), 4.5-4.35 (m, 1H and 2H, major rotamer), 4.3-4.25 (m, 2H, minor rotamer), 3.814 (s, 3H, major rotamer), 3.807 (s, 3H, minor rotamer), 3.75-3.5 (m, 1H), 3.35 (m, 1H), 2.2-1.8 (m, 4H)

25 ^{13}C -NMR (100 MHz; CD_3OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 173.3, 173.2, 171.3, 171.0, 153.9, 152.4

APCI-MS: $(\text{M} + 1) = 511/513 \text{ m/z}$

Example 34

30 $\text{Ph}(3\text{-Cl})(5\text{-OCHF}_2)\text{-(R)CH(OH)C(O)-Aze-NH-CH}_2\text{-(2-amidino)-5-pyridinyl}$



(i) 6-Cyanonicotinic acid

To a solution of the nicotinic acid N-oxide (51 g, 0.37 mol) in 1.2 L of DMF, NaCN (54 g, 1.1 mol) was added, followed by triethylamine (255 mL, 1.83 mol) and TMSCl (185 mL). The reaction mixture was stirred at 110°C for 10 h, filtered and the filtrate was concentrated. The residue was dissolved in 100 mL of 2N HCl and extracted with methylene chloride. The organic layers were combined, concentrated and recrystallised from water to yield 12 g (22%) of the product.

(ii) 5-(Hydroxymethyl)pyridine-2-carbonitrile

To a solution of 6-cyanonicotinic acid (12g, 0.081mol; see step (i) above) in THF at 0°C, Et₃N (12.4 mL, 0.0892 mol) was added followed by ethyl chloroformate (8.53 mL, 0.0892 mol). The reaction mixture was stirred for 15 min and NaBH₄ (6.14 g, 0.162 mol) was added. Then the mixture was stirred at RT overnight, quenched with water and extracted with methylene chloride. The organic layer was concentrated and purified by column chromatography to yield 4g (20%) of the alcohol.

(iii) 5-(Azidomethyl)pyridine-2-carbonitrile

5-(Hydroxymethyl)pyridine-2-carbonitrile (4 g, 0.03 mol; see step (ii) above) was dissolved in 25 mL of methylene chloride and cooled in an ice bath. Mesyl chloride (2.32 mL, 0.0300 mol) and then triethylamine (4.6 mL, 0.033 mol) were added dropwise. The reaction mixture was stirred and after work up the crude mesylate was treated with NaN₃ (7.35 g, 0.113 mol) in 20 mL of DMF. The reaction mixture was stirred at 40°C for 2 h,

diluted with water and extracted with ethyl acetate. The organic layer was concentrated to yield 3.95g (83%) of the crude azide.

(iv) 5-(tert-Butoxycarbonylaminomethyl)pyridine-2-carbonitrile

5 To a solution of 5-(azidomethyl)pyridine-2-carbonitrile (3.95 g, 0.0248 mol; see step (iii) above) in 30 mL of THF and 10 mL of water, triphenyl phosphine (7.8 g, 0.0298 mol) was added and the resultant stirred for 24 h. Then, triethylamine (3.8 mL, 0.027 mol) was added, followed by Boc anhydride (5.4 g, 0.025 mol) and stirring for 2 h. The reaction mixture was partitioned between water and ethyl acetate. The organic layer was
10 concentrated and purified by column chromatography to yield 2.1 g (36%) of the sub-title compound.

¹H NMR (300 MHz, CDCl₃) δ 8.6 (s, 1H), 8.0 (d, 1H), 8.9 (d, 1H), 4.1 (m, 2H), 1.4 (s, 9H)

15

(v) 5-(Aminomethyl)pyridine-2-carbonitrile x 2 HCl

5-(tert-Butoxycarbonylaminomethyl)pyridine-2-carbonitrile (0.200 g, 0.86 mmol, see step (iv) above) was dissolved in 10 mL of EtOAc saturated with HCl(g) and was stirred for 30 min. The solvent was evaporated and 0.175 g (99%) of the sub-title compound was
20 obtained as its dihydrochloride salt.

¹H NMR (500 MHz, D₂O) δ 8.79 (s, 1H), 8.17 (d, 1H), 8.05 (d, 1H), 4.38 (s, 2H)

(vi) Boc-Aze-NH-CH₂-5-Py(2-CN)

25 To a mixture of 5-(aminomethyl)pyridine-2-carbonitrile x 2 HCl (0.175 g, 0.85 mmol; see step (v) above), Boc-Aze-OH (0.201 g, 1.00 mmol) and TBTU (0.321 g, 1.00 mmol) in 5 mL of DMF was added dimethylaminopyridine (0.367 g, 3.00 mmol). The mixture was stirred overnight and subsequently poured into water and extracted three times with EtOAc. The combined organic phase was washed with aqueous sodium bicarbonate, dried
30 (Na₂SO₄) and evaporated. The crude product started to crystallise and was used as such in the next step. Yield: 0.23 g (73%).

¹H NMR (500 MHz, CDCl₃) δ 8.66 (s, 1H), 8.2-7.8 (broad, 1H), 7.79 (d, 1H), 7.67 (d, 1H), 4.73 (m, 1H), 4.65-4.5 (m, 2H), 3.94 (m, 1H), 3.81 (m, 1H), 2.6-2.35 (m, 2H), 1.8 (broad, 1H), 1.45 (s, 9H)

5 (vii) H-Aze-NH-CH₂-5-Py(2-CN) x 2 HCl

Boc-Aze-NH-CH₂-5-Py(2-CN) (0.23 g, 0.73 mmol; see step (vi) above) was dissolved in 10 mL of EtOAc saturated with HCl(g) and was stirred for 30 min. The solvent was evaporated and 0.21 g (100%) of the sub-title compound was obtained as its dihydrochloride salt.

10 ¹H NMR (500 MHz, D₂O) δ 8.64 (s, 1H), 8.0-7.9 (m, 2H), 5.19 (m, 1H), 4.65-4.55 (m, 2H), 4.20 (m, 1H), 4.03 (m, 1H), 2.88 (m, 1H), 2.64 (m, 1H)

(viii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Aze-NH-CH₂-5-Py(2-CN)

15 To a mixture of H-Aze-NH-CH₂-5-Py(2-CN) x 2 HCl (0.206 g, 0.713 mmol; see step (vii) above), Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (0.180 g, 0.713 mmol; see Example 1(viii) above) and PyBOP (0.408 g, 0.784 mmol) in 5 mL of DMF was added dimethylaminopyridine (0.367 g, 3.00 mmol). The mixture was stirred overnight and subsequently poured into water and extracted three times with EtOAc. The combined
20 organic phase was washed with aqueous sodium bicarbonate, dried (Na₂SO₄) and evaporated. The crude product was flash chromatographed on silica gel with EtOAc gave a pure product. Yield: 0.197 g (61%).

¹H NMR (500 MHz, CDCl₃) δ 8.63 (m, 1H), 8.22 (bt, 1H), 7.78 (m, 1H), 7.67 (m, 1H),
25 7.21 (m, 1H), 7.16 (m, 1H), 7.04 (m, 1H), 6.56 (t, 1H), 4.97 (bd, 1H), 4.92 (m, 1H), 4.6-4.5 (m, 2H), 4.40 (bd, 1H), 4.18 (m, 1H), 3.80 (m, 1H), 2.69 (m, 1H), 2.46 (m, 1H), 1.92 (s, 1H)

APCI-MS: (M + 1) = 451/453 m/z

30 (ix) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Aze-NH-CH₂-((2-amidino)-5-pyridinyl) x HOAc

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Aze-NH-CH₂-5-Py(2-CN) (0.200 g, 0.444 mmol; see step (viii) above), ammonium acetate (1.00 g, 0.0130 mol) and N-acetylcysteine (2.00 g, 0.0122 mol) in 10 mL of methanol was heated at 50°C for 2 days. Preparative RPLC with CH₃CN:0.1M NH₄OAc (30:79) and running the appropriate fractions again with CH₃CN:0.1M NH₄OAc (5:95 – 40:60) gave 60 mg (26%) of pure title compound as its acetate salt after freeze drying from water and acetonitrile. Purity: 100%.


¹H NMR (500 MHz, D₂O, mixture of rotamers) δ 8.68 (s, 1H, major rotamer), 8.62 (s, 1H, minor rotamer), 8.05-7.9 (m, 2H), 7.33 (m, 1H, rotamer), 7.27 (m, 1H, rotamer), 7.22 (m, 1H, rotamer), 7.17 (m, 1H, rotamer), 7.01 (m, 1H, rotamer), 6.84 (t, 1H), 5.32 (s, 1H, major rotamer), 5.20 (m, 1H, minor rotamer), 5.13 (s, 1H, minor rotamer), 4.88 (m, 1H, major rotamer), 4.65-4.55 (m, 2H, major rotamer), 4.45-4.35 (m, 1H, rotamer plus 1H, minor rotamer), 4.31 (d, 1H, minor rotamer), 4.2-4.05 (m, 1H plus 1H, rotamer), 2.80 (m, 1H, minor rotamer), 2.61 (m, 1H, major rotamer), 2.33 (m, 1H, major rotamer), 2.24 (m, 1H, minor rotamer), 1.93 (s, 3H)

¹³C-NMR (100 MHz; D₂O): (carbonyl and/or amidine carbons, mixture of rotamers) δ 181.6, 173.3, 172.7, 172.6, 172.3, 162.6, 162.3

APCI-MS: (M + 1) = 468/470 m/z

20 Example 35

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((2-methoxyamidino)-5-pyridinyl)



(i) Boc-NH-CH₂-[(2-(amino(hydroxylimino)methyl))-5-pyridinyl]

25 5-(*tert*-Butoxycarbonylaminomethyl)pyridine-2-carbonitrile (1.00 g, 4.29 mmol; see Example 34(iv) above) was dissolved in 10 mL of ethanol and hydroxylamine hydrochloride (0.894 g, 0.0129 mol) and triethyl amine (1.30 g, 0.0129 mol) were added.

The mixture was stirred at room temperature for 6 days. The mixture was partitioned between water and methylene chloride. The aqueous layer was extracted with methylene chloride and the combined organic phase was washed with water, dried (Na_2SO_4) and evaporated. Yield: 0.96 g (84%).

^1H NMR (400 MHz, acetone- d_6) δ 9.01 (bs, 1H), 8.50 (bs, 1H), 7.87 (m, 1H), 7.70 (m, 1H), 6.58 (broad, 1H), 5.70 (broad, 2H), 4.31 (d, 2H), 1.41 (s, 9H)

(ii) Boc-Aze-NH-CH₂-(2-(amidino)-5-pyridinyl) x HOAc

This reaction was carried out according to the method described in Judkins *et al*, *Synth. Comm.* (1998) 4351. A suspension of Boc-NH-CH₂-[(2-(amino(hydroxylimino)methyl))-5-pyridinyl] (0.910 g, 3.42 mmol; see step (i) above), acetic anhydride (0.35 mL, 3.7 mmol) and 0.35 g of 10% Pd/C (50% moisture) in 100 mL of acetic acid was hydrogenated at a pressure of 5 atm. for 5 h. The mixture was filtered through Celite and concentrated. The residue was freeze-dried from water and acetonitrile to give 0.97 g (92%) of the sub-title compound.

^1H NMR (500 MHz, CD_3OD) δ 8.74 (s, 1H), 8.12 (d, 1H), 7.98 (d, 1H), 4.38 (s, 2H), 1.92 (s, 3H), 1.46 (s, 9H)

(iii) Boc-NH-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl)

To a suspension of Boc-NH-CH₂-(2-(amidino)-5-pyridinyl) x HOAc (0.96 g, 3.1 mmol; see step (ii) above) in 75 mL of THF was added a solution of potassium carbonate (1.07 g, 7.7 mmol) and Teoc-*p*-nitrophenyl carbonate (1.14g, 4.02 mmol) in 15 mL of water. The mixture was stirred overnight. An excess of glycine and potassium carbonate was added, and the reaction was continued for 2 h. The THF was evaporated and the remainder was extracted three times with EtOAc. The combined organic phase was washed with water, dried (Na_2SO_4) and evaporated. The product could be used without further purification.

^1H NMR (500 MHz, CDCl_3) δ 9.31 (broad, 1H), 8.52 (s, 1H), 8.41 (d, 1H), 8.35 (broad, 1H), 7.74 (d, 1H), 4.97 (broad, 1H), 4.39 (m, 2H), 4.26 (m, 2H), 1.46 (s, 9H), 1.14 (m, 2H), 0.07 (s, 9H)

(iv) H₂N-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl) x 2 HCl

Boc-NH-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl) (0.23 g, 0.58 mmol; see step (iii) above) was dissolved in 25 mL of EtOAc saturated with HCl(g) and stirred for 30 min. The solvent was evaporated and the product used without further purification. Yield: 0.21 g (98%).

¹H NMR (500 MHz, D₂O) δ 8.89 (s, 1H), 8.25 (s, 2H), 4.55 (m, 2H), 4.42 (s, 2H), 1.20 (m, 2H), 0.09 (s, 9H)

(v) Boc-Aze-NH-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl)

To a solution of H₂N-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl) x 2 HCl (0.21 g, 0.57 mmol; see step (iv) above), Boc-Aze-OH (0.127 g, 0.631 mmol), and TBTU (233 mg, 0.726 mmol) in 5 mL of DMF was added dimethylaminopyridine (269 mg, 2.20 mmol). The mixture was stirred overnight, poured into 100 mL of water and extracted with EtOAc three times. The combined organic phase was washed with aqueous sodium bicarbonate and water, dried (Na₂SO₄) and evaporated. The crude product was flash chromatographed on silica gel with EtOAc to give 170 mg (56%) of the desired product.

¹H NMR (500 MHz, CDCl₃) δ 9.33 (broad, 1H), 8.54 (s, 1H), 8.41 (d, 1H), 8.36 (broad, 1H), 7.75 (m, 1H), 4.72 (m, 1H), 4.56 (m, 2H), 4.26 (m, 2H), 3.93 (m, 1H), 3.80 (m, 1H), 2.6-2.4 (m, 2H), 1.42 (s, 9H), 1.14 (m, 2H), 0.07 (s, 9H)

(vi) H-Aze-NH-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl) x 2 HCl

Boc-Aze-NH-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl) (170 mg, 0.356 mmol; see step (v) above) was dissolved in 25 mL of EtOAc saturated with HCl(g)

and stirred for 30 min. The solvent was evaporated and the product used without further purification. Yield: 160 mg (100%).

¹H NMR (500 MHz, CD₃OD) δ 9.00 (m, 1H), 8.84 (m, 1H), 8.23 (d, 2H), 8.10 (m, 1H),
5.09 (m, 1H), 4.7-4.6 (m, 2H), 4.51 (m, 2H), 4.14 (m, 1H), 3.97 (m, 1H), 2.86 (m, 1H),
2.58 (m, 1H), 1.22 (m, 2H), 0.11 (s, 9H)

(vii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-(2-(amino(trimethylsilylethylimino)methyl)-5-pyridinyl)

To a solution of H-Aze-NH-CH₂-(2-(amino(trimethylsilylethylimino)-methyl)-5-pyridinyl)
x 2 HCl (160 mg, 0.462 mmol; see step (vi) above), Ph(3-Cl)(5-OCHF₂)-
(R)CH(OH)C(O)OH (131 mg, 0.462 mmol; see Example 1(viii) above) and PyBOP (263
mg, 0.505 mmol) in 5 mL of DMF was added diisopropylethyl amine (0.30 mL, 1.71
mmol). The mixture was stirred overnight, poured into 100 mL of water and extracted
three times with EtOAc. The combined organic phase was washed with aqueous sodium
bicarbonate and water, dried (Na₂SO₄) and evaporated. The crude product was flash
chromatographed on silica gel with EtOAc:MeOH (95:5) to give 148 mg (52%) of the
desired product.

(viii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-(2-(methoxyamino(trimethylsilylethylimino)methyl)-5-pyridinyl)

A suspension of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)-C(O)-Aze-NH-CH₂-(2-(methoxyamino(trimethylsilylethylimino)methyl)-5-pyridinyl) (148 mg, 0.242 mmol; see
step (vii) above) and O-methylhydroxyl amine (202 mg, 2.42 mmol) in 10 mL of
acetonitrile was heated at 70°C for 3 h. The mixture was partitioned between water and
EtOAc. The aqueous layer was extracted twice with EtOAc and the combined organic
phase was washed with water, dried (Na₂SO₄) and evaporated. The crude material was
flash chromatographed on silica gel with EtOAc:MeOH (95:5) to give 44 mg (28%) of
pure material.

¹H NMR (500 MHz, CDCl₃) δ 8.55 (m, 1H), 8.05 (bt, 1H), 7.70 (m, 1H), 7.58 (s, 1H), 7.56
(d, 1H), 7.22 (m, 1H), 7.16 (m, 1H), 7.03 (m, 1H), 6.50 (t, 1H), 4.92 (s, 1H), 4.89 (m, 1H),

4.55-4.45 (m, 2H), 4.38 (broad, 1H), 4.2-4.1 (m, 3H), 4.00 (s, 3H), 3.73 (m, 1H), 2.69 (m, 1H), 2.44 (m, 1H), 0.97 (m, 2H), 0.02 (s, 9H)

(ix) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((2-methoxy-amidino)-5-pyridinyl)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-(2-(methoxyamino(tri-methylsilylethylimino)methyl)-5-pyridinyl) (44 mg, 0.069 mmol; see step (viii) above) was dissolved in 2 mL of TFA and allowed to react for 1 h. The TFA was evaporated and the residue was partitioned between EtOAc and aqueous sodium bicarbonate. The aqueous layer was extracted with EtOAc and the combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. Yield: 30 mg (88%). Purity: >95%.


¹H NMR (500 MHz, CDCl₃) δ 8.44 (m, 1H), 8.03 (bt, 1H), 7.91 (m, 1H), 7.60 (m, 1H), 7.19 (m, 1H), 7.13 (m, 1H), 7.00 (m, 1H), 6.52 (t, 1H), 5.6-5.45 (broad, 2H), 4.90 (s, 1H), 4.89 (m, 1H), 4.55-4.4 (m, 2H), 4.27 (broad, 1H), 4.12 (m, 1H), 3.92 (s, 3 H), 2.68 (m, 1H), 2.41 (m, 1H)

¹³C-NMR (100 MHz; CDCl₃): (carbonyl and/or amidine carbons) δ 173.0, 170.9, 152.6

APCI-MS: (M + 1) = 498/500 m/z

Example 36

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-amidino)-2-pyrimidinyl)



(i) 2-Amino-2-iminoethylcarbamate•AcOH

N-Boc-aminoacetonitrile (40.2 g, 257.4 mmol) and N-acetylcysteine (42.0 g, 257.4 mmol) were dissolved in methanol (300 mL) at 60°C and ammonia was passed through for 18 h. The solvent was removed *in vacuo*. After ion exchange chromatography (Amberlite IRA-

400 (AcOH)) and recrystallisation from acetone, 28.4 g (53%) of the sub-title compound was obtained as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 4.41 (t, *J* = 4.9 Hz, 1H), 4.01 (s, 2H), 2.91 (d, *J* = 5.0 Hz, 2H), 2.01 (s, 3H), 1.46 (s, 9H)

(ii) 1,3-Bis(dimethylamino)-2-cyanotrimethinium perchlorate

A solution of 3-dimethylaminoacrylonitrile (25.0 g, 260.0 mmol) in chloroform (75 mL) was added dropwise to a solution of (chloromethylene)dimethylammonium chloride (50.0 g, 390.1 mmol) in chloroform (175 mL) at 0°C. The reaction mixture was stirred an additional 2 h at 0°C, then allowed to warm to room temperature overnight, then subsequently heated for 8 h under reflux. The solvent was removed *in vacuo*. The residue was added to a mixture of sodium perchlorate (110 g, 0.898 mmol) in water (150 mL) and ethanol (300 mL). The mixture was heated under reflux for 15 min then cooled and allowed to stand overnight in a refrigerator. The precipitate was collected and recrystallized from ethanol to yield 23.8 g (52%) of the sub-title compound as colorless needles.

mp: 140-141°C

¹H NMR (300 MHz, CDCl₃) δ 8.24 (s, 2H), 3.59 (s, 6H), 3.51 (s, 6H)

(iii) Boc-NH-CH₂-(5-cyano)-2-pyrimidine

A mixture of *t*-butyl 2-amino-2-iminoethylcarbamate•AcOH (5.0 g, 23.8 mmol; see step (i) above) and 1,3-bis(dimethylamino)-2-cyanotrimethinium perchlorate (6.0 g, 23.8 mmol; see step (ii) above) in pyridine (300 mL) was stirred under nitrogen at 70-75 °C for 16 h and then heated under reflux for 6 h. The mixture was cooled to room temperature and the solvent was removed *in vacuo*. The residue was extracted with a hot mixture (1:1) of ethyl acetate and chloroform, filtered through a small pad of silica, and concentrated to give the crude product. Flash chromatography on silica eluting with chloroform gave 4.0 g (71%) of the title compound as colorless oil, which solidified upon standing.

mp: 86-87 °C

$R_f = 0.77$ (silica, 3:2 Ethyl Acetate/Chloroform)

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 9.25 (s, 2H), 7.39 (bt, 1H), 4.39 (d, $J = 6$ Hz, 2H), 1.38 (s, 9H).

^{13}C NMR (750 MHz, $\text{DMSO}-d_6$) δ 170.4, 160.3, 155.8, 115.2, 106.9, 80.0, 46.3, 28.1

5 APCI-MS: $(M + 1) = 235$ m/z

(iv) Boc-Aze-NH-CH₂-((5-cyano)-2-pyrimidinyl)

10 Boc-NH-CH₂-(5-cyano)-2-pyrimidine (1.14 g, 4.87 mmol; see step (iii) above) was dissolved in 50 mL of EtOAc saturated with HCl(g) and allowed to react for 1 h and concentrated. The residue was dissolved in 20 mL of DMF and cooled in an ice bath. Diisopropylethyl amine (3.5 mL, 0.020 mol), Boc-Aze-OH (1.08 g, 5.37 mmol) and HATU (2.80 g, 5.38 mmol) were added and the reaction mixture was stirred at room
15 temperature overnight. The solvent was evaporated and the product was purified by preparative RPLC using CH₃CN:0.1M NH₄OAc (40:60). The acetonitrile was evaporated and the aqueous layer was extracted three times with EtOAc. The combined organic layer was dried (MgSO₄) and evaporated. Yield: 1.12 g (72%).

20 ^1H NMR (400 MHz, CDCl_3) δ 8.95 (s, 2H), 4.82 (d, 2H), 4.74 (m, 1H), 3.95 (m, 1H), 3.84 (m, 1H), 2.6-2.4 (m, 2H), 1.47 (s, 9H)

(v) Boc-Aze-NH-CH₂-((5-amidino)-2-pyrimidinyl) x HOAc

25 A solution of Boc-Aze-NH-CH₂-((5-cyano)-2-pyrimidinyl) (0.83 g, 2.6 mmol; see step (iv) above), N-acetylcysteine (0.43 g, 2.6 mmol) and ammonium acetate (0.60 g, 7.8 mmol) in 10 mL of methanol was heated at 60°C under nitrogen for 2 days. The solvent was evaporated and the crude material was purified by preparative RPLC using a gradient of CH₃CN:0.1M NH₄OAc (5:95 to 100:0). The fractions of interest were freeze dried to give 1.0 g (93%) of the desired material.

30

^1H NMR (300 MHz, D_2O , signals obscured by the HDO signal) δ 9.17 (s, 2H), 4.1-3.9 (m, 2H), 2.60 (m, 1H), 2.29 (m, 1H), 1.93 (s, 3H), 1.44 (s, 9H)

(vi) Boc-Aze-NH-CH₂-[(5-(amino(trimethylsilylethylimino)methyl))-2-pyrimidinyl]

5 To a suspension of Boc-Aze-NH-CH₂-((5-amidino)-2-pyrimidinyl) x HOAc (0.95 g, 2.41 mmol; see step (v) above) in 50 mL of THF was added a solution of Teoc-*p*-nitrophenyl carbonate (0.85 g, 3.0 mmol) and potassium carbonate (1.0 g, 7.2 mmol) in 10 mL of water. The mixture was stirred for 24 h, concentrated and partitioned between water and methylene chloride. The organic layer was washed twice with saturated aqueous sodium
10 bicarbonate, dried (Na₂SO₄) and evaporated. The crude product was flash chromatographed on silica gel with heptane:EtOAc (1:1). Yield: 1.04 g (90%).

¹H NMR (300 MHz, CDCl₃) δ 9.16 (s, 2H), 4.80 (d, 2H), 4.73 (m, 1H), 4.26 (m, 2H), 4.0-3.8 (m, 2H), 2.6-2.4 (m, 2H), 1.47 (s, 9H), 1.12 (m, 2H), 0.07 (s, 9H)

15

(vii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-[(5-(amino(trimethylsilylethylimino)methyl))-2-pyrimidinyl]

Boc-Aze-NH-CH₂-[(5-(amino(trimethylsilylethylimino)methyl))-2-pyrimidinyl] (0.209 g, 0.437 mmol; see step (vi) above) was dissolved in 25 mL of EtOAc saturated with HCl(g) and allowed to react for 15 min. The solvent was evaporated and the remainder was
20 dissolved in 4 mL of DMF. Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (0.100 g, 0.396 mmol; see Example 1(viii) above), PyBOP (0.231 g, 0.444 mmol) and diisopropylethyl amine (0.208 g, 1.61 mmol) were added, and the mixture was stirred for 80 min. The reaction mixture was poured into 100 mL of water and extracted three times with EtOAc.
25 The combined organic layer was washed with brine, dried (Na₂SO₄) and evaporated. The crude product was purified by preparative RPLC using CH₃CN:0.1M NH₄OAc (1:1). Yield: 63 mg (26%).

¹H NMR (400 MHz, CDCl₃, mixture of rotamers) δ 9.3 (broad, 1H), 9.03 (s, 2H, minor rotamer), 9.00 (s, 2H, major rotamer), 8.25 (m, 1H, major rotamer), 7.9 (broad, 1H), 7.80 (m, 1H, minor rotamer), 7.2-6.9 (m, 3H), 6.50 (t, 1H), 5.14 (s, 1H, minor rotamer), 5.08 (m, 1H, minor rotamer), 4.94 (s, 1H, major rotamer), 4.80 (m, 1H, major rotamer), 4.7-4.4

30

(m, 2H), 4.3-3.9 (m, 3H), 3.74 (m, 1H, major rotamer), 2.7-2.1 (m, 2H), 1.03 (m, 2H), 0.01 (s, 9H)

(viii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-amidino)-2-pyrimidinyl) x

5 TFA

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-[(5-amino(trimethylsilylethylimino)methyl))-2-pyrimidinyl] (21 mg, 0.034 mmol; see step (vii) above) was dissolved in 0.5 mL of methylene chloride and cooled in an ice bath. TFA (2 mL) was added and the mixture was stirred for 60 min and then concentrated. The product was
10 freeze-dried from water and acetonitrile. Yield: 20 mg (100%). Purity: 100%.

¹H NMR (400 MHz, CD₃OD, mixture of rotamer, signals obscured by the HDO signal) δ 9.08 (s, 2H), 7.4-7.1 (m, 3H), 6.88 (t, 1H, major rotamer), 6.85 (t, 1H, minor rotamer), 5.30 (m, 1H, minor rotamer), 5.22 (s, 1H, minor rotamer), 5.20 (s, 1H, major rotamer), 4.73 (m,
15 1H, major rotamer), 4.34 (m, 1H, rotamer), 4.21 (m, 1H, rotamer), 4.15-3.95 (m, 2H, rotamers), 2.73 (m, 1H, rotamer), 2.57 (m, 1H, rotamer), 2.45-2.25 (m, 2H, rotamers)


¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 173.0, 172.6, 172.1, 171.0, 163.4.

APCI-MS: (M + 1) = 469/471 m/z

20

Example 37

25 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-methoxyamidino)-2-pyrimidinyl)



(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-[(5-(methoxyamino-(trimethylsilylethylimino)methyl))-2-pyrimidinyl]

A suspension of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-[(5-(amino(trimethylsilylethylimino)methyl))-2-pyrimidinyl] (40 mg, 0.065 mmol; see
5 Example 36(vii) above) and O-methylhydroxyl amine (33 mg, 0.40 mmol) in 3 mL of acetonitrile was heated at 70°C for 3 h. The mixture was partitioned between water and EtOAc. The aqueous layer was extracted twice with EtOAc and the combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. Yield: 33 mg (79%).

¹H NMR (400 MHz, CDCl₃, mixture of rotamers) δ 8.76 (s, 2H, major rotamer), 8.70 (s, 2H, rotamer), 8.18 (m, 1H), 7.62 (s, 1H), 7.4-6.9 (m, 4H), 6.50 (bt, 1H), 5.3-4.5 (m, 4H), 4.2-4.05 (m, 3H), 3.96 (s, 3H), 3.68 (m, 1H), 2.8-2.2 (m, 2H), 2.1 (broad, 1H), 0.96 (m, 2H), 0.01 (s, 9H)


(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-[(5-methoxyamidino)-2-pyrimidinyl]

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-[(5-(methoxyamino-(trimethylsilylethylimino)methyl))-2-pyrimidinyl] (33 mg, 0.052 mmol; see step (i) above)
20 was dissolved in 0.5 mL of methylene chloride and cooled in an ice bath. TFA (2 mL) was added and the mixture was stirred for 2 h and then concentrated. The product was freeze dried from water and acetonitrile. Yield: 31 mg (81%). Purity: 100%.

¹H NMR (400 MHz, CD₃OD, mixture of rotamer, signals obscured by the HDO signal)
25 δ 8.96 (s, 2H, rotamer), 8.94 (s, 2H, rotamer), 7.4-7.3 (m, 1H), 7.2-7.1 (m, 2H), 6.88 (t, 1H, rotamer), 6.85 (t, 1H, rotamer), 5.29 (m, 1H, rotamer), 5.24 (s, 1H, rotamer), 5.20 (s, 1H, rotamer), 4.75-4.55 (m, 2H), 4.33 (m, 1H, rotamer), 4.19 (m, 1H, rotamer), 4.15-3.95 (m, 2H, rotamers), 3.88 (s, 3H, rotamer), 3.86 (s, 3H, rotamer), 2.72 (m, 1H, rotamer), 2.56 (m, 1H, rotamer), 2.45-2.25 (m, 2H, rotamers)

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ
30 172.8, 172.6, 172.1, 171.8, 167.8, 167.7, 155.1, 152.3, 152.1

APCI-MS: (M + 1) = 499/501 m/z

Example 38**Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(3-F)****(i) 2-Fluoro-4-vinylbenzonitrile**

A solution of 4-bromo-2-fluorobenzonitrile (4.92 g, 0.0246 mol), vinyltributyltin (0.78 g, 0.246 mol), and tetrakis(triphenyl)phosphine (0.67 g, 0.58 mmol) in 250 mL of toluene was refluxed under nitrogen overnight. The solvent was evaporated and the residue was flash chromatographed on silica gel with heptane:CH₂Cl₂ (1:1) to pure CH₂Cl₂. A colourless oil was obtained that crystallised. Yield: 3.0 g (82%).

¹H NMR (300 MHz, CDCl₃) δ 7.56 (m, 1H), 7.3-7.2 (m, 2H), 6.69 (m, 1H), 5.89 (d, 1H), 5.51 (d, 1H)

(ii) 2-Fluoro-4-hydroxymethylbenzonitrile

Into a cooled solution (-78°C) of 2-fluoro-4-vinylbenzonitrile (1.3 g, 8.8 mmol; see step (i) above) in 40 mL of CH₂Cl₂ and 5 mL of methanol was bubbled ozone (50 L/h, 29 g/m³) for 30 min. Argon was subsequently bubbled through to remove excess ozone. Sodium borohydride (0.67 g, 0.018 mol) was added and the cooling bath was removed. The mixture was stirred and allowed to react for 1 h. The mixture was evaporated and 2M HCl was added. The mixture was extracted twice with diethyl ether and the combined ether fraction was dried (Na₂SO₄) and evaporated. The crude product crystallised. Yield: 1.1 g (81%).

¹H NMR (300 MHz, CDCl₃) δ 7.59 (m, 1H), 7.3-7.2 (m, 2H), 4.79 (d, 2H), 2.26 (t, 1H)

(iii) 4-Cyano-3-fluorobenzyl methanesulfonate

2-Fluoro-4-hydroxymethylbenzonitrile (1.3 g, 8.6 mmol; see step (ii) above) was dissolved in 50 mL of CH_2Cl_2 and cooled on an ice bath. Triethylamine (0.87 g, 8.6 mmol) and methanesulfonyl chloride (0.99 g, 8.7 mmol) were added. After stirring for 1.5 h the reaction mixture was washed with 1M HCl. The organic phase was dried (Na_2SO_4) and evaporated. The product could be used without purification. Yield of a colourless oil: 1.8 g (92%).

^1H NMR (400 MHz, CDCl_3) δ 7.66 (m, 1H), 7.35-7.3 (m, 2H), 5.26 (s, 2H), 3.07 (s, 3H)

(iv) 4-Azidomethyl-2-fluorobenzonitrile

To an ice cooled solution of 4-cyano-3-fluorobenzyl methanesulfonate (1.8 g, 7.9 mmol; see step (iii) above) was added sodium azide (0.80 g, 0.012 mol). The mixture was stirred overnight and then poured into 200 mL of water and extracted three times with diethyl ether. The combined ethereal phase was washed five times with water, dried (Na_2SO_4) and evaporated. The crude colourless oil could be used without further purification. Yield: 1.2 g (87%).

^1H NMR (300 MHz, CDCl_3) δ 7.64 (m, 1H), 7.25-7.18 (m, 2H), 4.47 (s, 2H)

(v) 4-Aminomethyl-2-fluorobenzonitrile

To a suspension of stannous chloride dihydrate (0.45 g, 2.4 mmol) in 20 mL of acetonitrile under stirring was added thiophenol (1.07 g, 9.7 mmol) and triethylamine (0.726 g, 7.17 mmol). Thereafter was added a solution of 4-azidomethyl-2-fluorobenzonitrile (0.279 g, 1.58 mmol; see step (iv) above) in a few mLs of acetonitrile. After 1.5 h, the azide was consumed and the solvent was evaporated. The residue was dissolved in methylene chloride and washed three times with 2M NaOH. The organic phase was extracted twice with 1M HCl. The combined acidic aqueous phase was washed with methylene chloride and then made alkaline with 2M NaOH and extracted three times with methylene chloride. The organic phase was dried (Na_2SO_4) and evaporated to give 0.172 g (72%) of the desired sub-title compound which could be used without purification.

¹H NMR (400 MHz, CDCl₃) δ 7.58 (m, 1H), 7.3-7.2 (m, 2H), 3.98 (s, 2H), 1.55-1.35 (broad, 2H)

5 (vi) Boc-Aze-NHCH₂-Ph(3-F, 4-CN)

To an ice cooled solution of Boc-Aze-OH (0.194 g, 0.96 mmol) in 5 mL of DMF was added TBTU (0.50 g, 9.6 mmol). After 30 min another solution, comprising 4-aminomethyl-2-fluorobenzonitrile (0.17 g, 0.81 mmol; see step (v) above) and diisopropylethyl amine (0.326 g, 2.53 mmol) in 7 mL of DMF was added. The resulting
10 solution was stirred overnight at room temperature. The solvent was evaporated and the product was purified by preparative RPLC using CH₃CN:0.1M NH₄OAc (50:50). Freeze-drying gave 0.237 g (74%) of the desired sub-title compound.

¹H NMR (300 MHz, CD₃OD) δ 7.70 (m, 1H), 7.35-7.25 (m, 2H), 4.65-4.35 (m, 3H), 4.0-
15 3.85 (m, 2H), 2.51 (m, 1H), 2.19 (m, 1H), 1.40 (s, 9H)

(vii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NHCH₂-Ph(3-F, 4-CN)

Boc-Aze-NHCH₂-Ph(3-F, 4-CN) (0.118 g, 0.354 mmol; from step (vi) above) was dissolved in 30 mL of EtOAc saturated with HCl(g). The reaction was stirred for 20 min
20 and evaporated. The resulting dihydrochloride and HATU (0.152 g, 0.400 mmol) were dissolved in 5 mL of DMF. That solution was added to an ice cooled solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (0.101 g, 0.400 mmol; see Example 1(viii) above) in 5 mL of DMF. The reaction was stirred overnight at ambient temperature. The solvent was evaporated and the product was purified by preparative RPLC with CH₃CN:0.1M NH₄OAc
25 (50:50). Freeze-drying gave 0.130 g (77%) of the desired sub-title compound.

¹H NMR (500 MHz, CD₃OD mixture of rotamers) δ 7.7-7.6 (m, 1H), 7.35-7.1 (m, 5H), 6.88 (t, 1H, rotamer), 6.86 (t, 1H, rotamer), 5.25-5.1 (m, 1H plus minor rotamer from the following proton), 4.80 (m, 1H, major rotamer), 4.6-4.4 (m, 2H), 4.36 (m, 1H, major
30 rotamer), 4.18 (m, 1H, major rotamer), 4.07 (m, 1H, minor rotamer), 3.98 (m, 1H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.53 (m, 1H, major rotamer), 2.29 (m, 1H, major rotamer), 2.16 (m, 1H, minor rotamer)

(viii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(3-F)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-NHCH₂-Ph(3-F, 4-CN) (0.130 g, 0.278 mmol; see step (vii) above) was dissolved in 80 mL of ethanol saturated with HCl(g). The mixture was allowed to react at room temperature overnight. The solvent was evaporated and the residue was re-dissolved in 100 mL of ethanol saturated with NH₃(g). The reaction was allowed to proceed slowly at room temperature for two days. The temperature was raised to 50°C and the reaction continued for another 3 days. The starting material was consumed and the solvent was evaporated. The product was purified by preparative RPLC and freeze-dried to give 17 mg (13%) of the title compound as its HOAc salt.


¹H NMR (600 MHz, CD₃OD mixture of rotamers) δ 7.65-7.6 (m, 1H), 7.4-7.3 (m, 3H), 7.25-7.1 (m, 2H), 7.15-6.7 (m, 1H), 5.25-5.1 (m, 1H plus minor rotamer of the following proton), 4.8 (m, 1H, major rotamer partially hidden by CD₃OH), 4.6-3.95 (m, 4H), 2.69 (m, 1H, minor rotamer), 2.56 (m, 1H, major rotamer), 2.28 (m, 1H, major rotamer), 2.14 (m, 1H, minor rotamer), 1.90 (s, 3H)

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 180.6, 173.4, 173.1, 172.9, 164.5, 162.3, 159.8

APCI-MS: (M + 1) = 485/487 m/z

Example 39

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)



(i) 2,6-Difluoro-4[(methylsulfinyl)(methylthio)methyl]benzonitrile

(Methylsulfinyl)(methylthio)methane (7.26g, 0.0584 mol) was dissolved in 100 mL of dry THF under argon and was cooled to -78°C . Butyllithium in hexane (16 mL 1.6M, 0.0256 mol) was added dropwise with stirring. The mixture was stirred for 15 min. Meanwhile, a solution of 3,4,5-trifluorobenzonitrile (4.0 g, 0.025 mmol) in 100 mL of dry THF was cooled to -78°C under argon and the former solution was added through a cannula to the latter solution over a period of 35 min. After 30 min, the cooling bath was removed and when the reaction had reached room temperature it was poured into 400 mL of water. The THF was evaporated and the remaining aqueous layer was extracted three times with diethyl ether. The combined ether phase was washed with water, dried (Na_2SO_4) and evaporated. Yield: 2.0 g (30%).

^1H NMR (500 MHz, CDCl_3) δ 7.4-7.25 (m, 2H), 5.01 (s, 1H, diastereomer), 4.91 (s, 1H, diastereomer), 2.88 (s, 3H, diastereomer), 2.52 (s, 3H, diastereomer), 2.49 (s, 3H, diastereomer), 2.34 (s, 3H, diastereomer), 1.72 (broad, 1H)

(ii) 2,6-Difluoro-4-formylbenzonitrile

2,6-Difluoro-4[(methylsulfinyl)(methylthio)methyl]benzonitrile (2.17 g, 8.32 mmol; see step (i) above) was dissolved in 90 mL of THF and 3.5 mL of concentrated sulfuric acid was added. The mixture was left at room temperature for 3 days and subsequently poured into 450 mL of water. Extraction three times with EtOAc followed and the combined ethereal phase was washed twice with aqueous sodium bicarbonate and with brine, dried (Na_2SO_4) and evaporated. Yield: 1.36 g (98%). The position of the formyl group was established by ^{13}C NMR. The signal from the fluorinated carbons at 162.7 ppm exhibited the expected coupling pattern with two coupling constants in the order of 260 Hz and 6.3 Hz respectively corresponding to an *ipso* and a *meta* coupling from the fluorine atoms.

^1H NMR (400 MHz, CDCl_3) δ 10.35 (s, 1H), 7.33 (m, 2H)

(iii) 2,6-Difluoro-4-hydroxymethylbenzonitrile

2,6-Difluoro-4-formylbenzonitrile (1.36 g, 8.13 mmol; see step (ii) above) was dissolved in 25 mL of methanol and cooled on an ice bath. Sodium borohydride (0.307 g, 8.12 mmol) was added in portions with stirring and the reaction was left for 65 min. The solvent was evaporated and the residue was partitioned between diethyl ether and aqueous sodium bicarbonate. The ethereal layer was washed with more aqueous sodium bicarbonate and brine, dried (Na_2SO_4) and evaporated. The crude product crystallised soon and could be used without further purification. Yield: 1.24 g (90%).

^1H NMR (400 MHz, CDCl_3) δ 7.24 (m, 2H), 4.81 (s, 2H), 2.10 (broad, 1H)

(iv) 4-Cyano-2,6-difluorobenzyl methanesulfonate

To an ice cooled solution of 2,6-difluoro-4-hydroxymethylbenzonitrile (1.24 g, 7.32 mmol; see step (iii) above) and methanesulfonyl chloride (0.93 g, 8.1 mmol) in 60 mL of methylene chloride was added triethylamine (0.81 g, 8.1 mmol) with stirring. After 3 h at 0°C , the mixture was washed twice with 1M HCl and once with water, dried (Na_2SO_4) and evaporated. The product could be used without further purification. Yield: 1.61 g (89%).

^1H NMR (300 MHz, CDCl_3) δ 7.29 (m, 2H), 5.33 (s, 2H), 3.07 (s, 3H)

(v) 4-Azidomethyl-2,6-difluorobenzonitrile

A mixture of 4-cyano-2,6-difluorobenzyl methanesulfonate (1.61 g, 6.51 mmol; see step (iv) above) and sodium azide (0.72 g, 0.0111 mol) in 10 mL of water and 20 mL of DMF was stirred at room temperature overnight. The resultant was subsequently poured into 200 mL of water and extracted three times with diethyl ether. The combined ethereal phase was washed five times with water, dried (Na_2SO_4) and evaporated. A small sample was evaporated for NMR purposes and the product crystallised. The rest was evaporated cautiously but not until complete dryness. Yield (theoretically 1.26 g) was assumed to be almost quantitative based on NMR and analytical HPLC.

^1H NMR (400 MHz, CDCl_3) δ 7.29 (m, 2H), 4.46 (s, 2H)

(vi) 4-Aminomethyl-2,6-difluorobenzonitrile

This reaction was carried out according to the procedure described in *J. Chem. Res. (M)* (1992) 3128. To a suspension of 520 mg of 10% Pd/C (50% moisture) in 20 mL of water was added a solution of sodium borohydride (0.834 g, 0.0221 mol) in 20 mL of water. Some gas evolution resulted. 4-Azidomethyl-2,6-difluorobenzonitrile (1.26 g, 6.49 mmol; see step (v) above) was dissolved in 50 mL of THF and added to the aqueous mixture on an ice bath over 15 min. The mixture was stirred for 4 h, whereafter 20 mL of 2M HCl was added and the mixture was filtered through Celite. The Celite was rinsed with more water and the combined aqueous phase was washed with EtOAc and subsequently made alkaline with 2M NaOH. Extraction three times with methylene chloride followed and the combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. Yield: 0.87 g (80%).

¹H NMR (400 MHz, CDCl₃) δ 7.20 (m, 2H), 3.96 (s, 2H), 1.51 (broad, 2H)

(vii) 2,6-Difluoro-4-tert-butoxycarbonylaminomethylbenzonitrile

A solution of 4-aminomethyl-2,6-difluorobenzonitrile (0.876 g, 5.21 mmol; see step (vi) above) was dissolved in 50 mL of THF and di-tert-butyl dicarbonate (1.14 g, 5.22 mmol) in 10 mL of THF was added. The mixture was stirred for 3.5 h. The THF was evaporated and the residue was partitioned between water and EtOAc. The organic layer was washed three times with 0.5 M HCl and water, dried (Na₂SO₄) and evaporated. The product could be used without further purification. Yield: 1.38 g (99%).

¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 2H), 4.95 (broad, 1H), 4.43 (broad, 2H), 1.52 (s, 9H)

(viii) Boc-Pab(2,6-diF)(OH)

A mixture of 2,6-difluoro-4-tert-butoxycarbonylaminomethylbenzonitrile (1.38 g, 5.16 mmol; see step (vii) above), hydroxylamine hydrochloride (1.08 g, 0.0155 mol) and triethylamine (1.57 g, 0.0155 mol) in 20 mL of ethanol was stirred at room temperature for 36 h. The solvent was evaporated and the residue was partitioned between water and

methylene chloride. The organic layer was washed with water, dried (Na_2SO_4) and evaporated. The product could be used without further purification. Yield: 1.43 g (92%).

^1H NMR (500 MHz, CD_3OD) δ 7.14 (m, 2H), 4.97 (broad, 1H), 4.84 (broad, 2H), 4.40 (broad, 2H), 1.43 (s, 9H)

(ix) Boc-Pab(2,6-diF) x HOAc

This reaction was carried out according to the procedure described by Judkins *et al*, *Synth. Comm.* (1998) 4351. Boc-Pab(2,6-diF)(OH) (1.32 g, 4.37 mmol; see step (viii) above), acetic anhydride (0.477 g, 4.68 mmol) and 442 mg of 10% Pd/C (50% moisture) in 100 mL of acetic acid was hydrogenated at 5 atm pressure for 3.5 h. The mixture was filtered through Celite, rinsed with ethanol and evaporated. The residue was freeze-dried from acetonitrile and water and a few drops of ethanol. The sub-title product could be used without further purification. Yield: 0.149 g (99%).

15

^1H NMR (400 MHz, CD_3OD) δ 7.45 (m, 2H), 4.34 (s, 2H), 1.90 (s, 3H), 1.40 (s, 9H)

(x) Boc-Pab(2,6-diF)(Teoc)

To a solution of Boc-Pab(2,6-diF) x HOAc (1.56 g, 5.49 mmol; see step (ix) above) in 100 mL of THF and 1 mL of water was added 2-(trimethylsilyl)ethyl p-nitrophenyl carbonate (1.67 g, 5.89 mmol). A solution of potassium carbonate (1.57 g, 0.0114 mol) in 20 mL of water was added dropwise over 5 min. The mixture was stirred overnight. The THF was evaporated and the residue was partitioned between water and methylene chloride. The aqueous layer was extracted with methylene chloride and the combined organic phase was washed twice with aqueous sodium bicarbonate, dried (Na_2SO_4) and evaporated. Flash chromatography on silica gel with heptane/EtOAc = 2/1 gave 1.71 g (73%) of pure compound.

25

^1H NMR (400 MHz, CDCl_3) δ 7.43 (m, 2H), 4.97 (broad, 1H), 4.41 (broad, 2H), 4.24 (m, 2H), 1.41 (s, 9H), 1.11 (m, 2H), 0.06 (s, 9H)

30

(xi) Boc-Aze-Pab(2,6-diF)(Teoc)

Boc-Pab(2,6-diF)(Teoc) (1.009 g, 2.35 mmol; see step (x) above) was dissolved in 50 mL of EtOAc saturated with HCl(g). The mixture was left for 10 min., evaporated and dissolved in 18 mL of DMF, and then cooled on an ice bath. Boc-Aze-OH (0.450 g, 2.24 mmol), PyBOP (1.24 g, 2.35 mmol) and lastly diisopropylethyl amine (1.158 g, 8.96 mmol) were added. The reaction mixture was stirred for 2 h and then poured into 350 mL of water and extracted three times with EtOAc. The combined organic phase was washed with brine, dried (Na₂SO₄) and evaporated. Flash chromatography on silica gel with heptane:EtOAc (1:3) gave 1.097 g (96%) of the desired compound.

¹H NMR (500 MHz, CDCl₃) δ 7.46 (m, 2H), 4.65-4.5 (m, 3H), 4.23 (m, 2H), 3.87 (m, 1H), 3.74 (m, 1H), 2.45-2.3 (m, 2H), 1.40 (s, 9H), 1.10 (m, 2H), 0.05 (s, 9H)

(xii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(Teoc)

Boc-Aze-Pab(2,6-diF)(Teoc) (0.256 g, 0.500 mmol; see step (xi) above) was dissolved in 20 mL of EtOAc saturated with HCl(g). The mixture was left for 10 min. and evaporated and dissolved in 5 mL of DMF. Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (0.120 g, 0.475 mmol; see Example 1(viii) above), PyBOP (0.263 g, 0.498 mmol) and lastly diisopropylethyl amine (0.245 g, 1.89 mmol) were added. The reaction mixture was stirred for 2 h and then poured into 350 mL of water and extracted three times with EtOAc. The combined organic phase was washed with brine, dried (Na₂SO₄) and evaporated. Flash chromatography on silica gel with EtOAc gave 0.184 g (60%) of the desired sub-title compound.

¹H NMR (400 MHz, CD₃OD, mixture of rotamers) δ 7.55-7.45 (m, 2H), 7.32 (m, 1H, major rotamer), 7.27 (m, 1H, minor rotamer), 7.2-7.1 (m, 2H), 6.90 (t, 1H, major rotamer), 6.86 (t, 1H, minor rotamer), 5.15 (s, 1H, major rotamer), 5.12 (m, 1H, minor rotamer), 5.06 (s, 1H, minor rotamer), 4.72 (m, 1H, major rotamer), 4.6-4.45 (m, 2H), 4.30 (m, 1H, major rotamer), 4.24 (m, 2H), 4.13 (m, 1H, major rotamer), 4.04 (m, 1H, minor rotamer), 3.95 (m, 1H, minor rotamer), 2.62 (m, 1H, minor rotamer), 2.48 (m, 1H, major rotamer), 2.22 (m, 1H, major rotamer), 2.10 (m, 1H, minor rotamer), 1.07 (m, 2H), 0.07 (m, 9H)

(xiii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(Teoc) (81 mg, 0.127 mmol; see step (xii) above) was dissolved in 0.5 mL of methylene chloride and cooled on an ice bath. TFA (3 mL) was added and the reaction was left for 75 min. The TFA was evaporated and the residue was freeze dried from water and acetonitrile. The crude product was purified
5 by preparative RPLC with CH₃CN:0.1M NH₄OAc (35:65) to produce 39 mg (55%) of the title compound as its HOAc salt, purity: 99%.


¹H NMR (400 MHz, CD₃OD mixture of rotamers) δ 7.5-7.4 (m, 2H), 7.32 (m, 1H, major rotamer), 7.28 (m, 1H, minor rotamer), 7.2-7.1 (m, 3H) 6.90 (t, 1H, major rotamer), 6.86 (t,
10 minor rotamer), 5.15 (s, 1H, major rotamer), 5.14 (m, 1H, minor rotamer), 5.07 (s, 1H, minor rotamer), 4.72 (m, 1H, major rotamer), 4.65-4.45 (m, 2H), 4.30 (m, 1H, major rotamer), 4.16 (m, 1H, major rotamer), 4.03 (m, 1H, minor rotamer), 3.95 (m, 1H, minor rotamer), 2.63 (m, 1H, minor rotamer), 2.48 (m, 1H, major rotamer), 2.21 (m, 1H, major rotamer), 2.07 (m, 1H, minor rotamer), 1.89 (s, 3H)

15 ¹³C-NMR (75 MHz; CD₃OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 171.9, 171.2, 165.0, 162.8, 160.4

APCI-MS: (M + 1) = 503/505 m/z.

Example 40

20 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(OMe)



(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(OMe,Teoc)

25 A mixture of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(Teoc) (64 mg, 0.099 mmol; see Example 39(xii) above) and O-methyl hydroxylamine hydrochloride (50 mg,

0.60 mmol) in 4 mL of acetonitrile was heated at 70°C for 3 h. The solvent was evaporated and the residue was partitioned between water and EtOAc. The aqueous layer was extracted twice with EtOAc and the combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. The product could be used without further purification.

5 Yield: 58 mg (87%).

¹H NMR (400 MHz, CDCl₃) δ 7.90 (bt, 1H), 7.46 (m, 1H), 7.25-6.95 (m, 5H), 6.51, t, 1H), 4.88 (s, 1H), 4.83 (m, 1H), 4.6-4.5 (m, 2H), 4.4-3.9 (m, 4H), 3.95 (s, 3H), 3.63 (m, 1H), 2.67 (m, 1H), 2.38 (m, 1H), 1.87 (broad, 1H), 0.98 (m, 2H), 0.01, s, 9H)

10

(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(OMe)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(OMe, Teoc) (58 mg, 0.086 mmol; see step (i) above) was dissolved in 3 mL of TFA, cooled on an ice bath and allowed to react for 2 h. The TFA was evaporated and the residue dissolved in EtOAc. The organic
15 layer was washed twice with aqueous sodium carbonate and water, dried (Na₂SO₄) and evaporated. The residue was freeze-dried from water and acetonitrile to give 42 mg (92%) of the title compound. Purity: 94%.


¹H NMR (300 MHz, CDCl₃) δ 7.95 (bt, 1H), 7.2-7.1 (m, 4H), 6.99 (m, 1H), 6.52 (t, 1H),
20 4.88 (s, 1H), 4.85-4.75 (m, 3H), 4.6-4.45 (m, 2H), 4.29 (broad, 1H), 4.09 (m, 1H), 3.89 (s, 3H), 3.69 (m, 1H), 2.64 (m, 1H), 2.38 (m, 1H), 1.85 (broad, 1H)

¹³C-NMR (100 MHz; CDCl₃): (carbonyl and/or amidine carbons) δ 172.1, 169.8, 151.9

APCI-MS: (M + 1) = 533/535 m/z

25 *Example 41*

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)



(i) 2,5-Difluoro-4-[(methylsulfinyl)(methylthio)methyl]benzonitrile

(Methylsulfinyl)(methylthio)methane (3.16 g, 0.0255 mol) was dissolved in 50 mL of dry THF under argon and then cooled to -78°C . Butyllithium in hexane (16 mL 1.6M, 0.0256 mol) was added dropwise with stirring. The mixture was stirred for 15 min. Meanwhile a solution of 2,4,5-trifluorobenzonitrile (2.0 g; 0.013 mol) in 50 mL of dry THF was cooled to -78°C under argon and the former solution was added through a cannula to the latter solution over a period of 3-5 min. After 30 min, the cooling bath was removed and when the reaction had reached room temperature it was poured into 200 mL of water. The THF was evaporated and the remaining aqueous layer was extracted three times with diethyl ether. The combined ether phase was washed with water, dried (Na_2SO_4) and evaporated. The crude product started to crystallise and could be used as such in the next step. Yield: 2.8 g (84%).

^1H NMR (500 MHz, CDCl_3) δ 7.51-7.44 (m, 2H, major diastereomer), 7.39 (dd, 1H, minor diastereomer), 5.00 (s, 1H, minor diastereomer), 4.92 (s, 1H, major diastereomer), 2.59 (s, 3H, minor diastereomer), 2.56 (s, 1H, major diastereomer), 2.46 (s, 1H, minor diastereomer), 2.40 (s, 1H, major diastereomer)

(ii) 2,5-Difluoro-4-formylbenzonitrile

2,5-Difluoro-4-[(methylsulfinyl)(methylthio)methyl]benzonitrile (2.8 g, 0.0107 mol; see step (i) above) was dissolved in 100 mL of THF and 6.5 g of concentrated sulfuric acid was added. The mixture was left at room temperature for 6 days and subsequently poured into 500 mL of water. Extraction three times with diethyl ether followed and the combined ethereal phase was washed several times with water, dried (Na_2SO_4) and evaporated. The crude product was flash chromatographed on silica gel using heptane:EtOAc (8:2). Yield:

1.2 g (67%). The position of the formyl group was established by use of ^{13}C NMR. The carbon signals from the fluorinated carbons at 160.1 and 158.4 respectively were doublets and not quartets, which they would have been if the formyl group had been in the 2-position.

^1H NMR (300 MHz, CDCl_3) δ 10.36 (d, 1H), 7.72 (dd, 1H), 7.54 (dd, 1H)

(iii) 2,5-Difluoro-4-hydroxymethylbenzonitrile

2,5-Difluoro-4-formylbenzonitrile (3.60 g, 0.0215 mol; see step (ii) above) was dissolved in 50 mL of methanol and cooled on an ice bath. Sodium borohydride (0.815 g, 0.0215 mol) was added in portions with stirring and the reaction was left for 45 min. Water (300 mL) was added and thereafter carefully 2M HCl was added until an acidic pH was attained. The mixture was extracted three times with diethyl ether, and the combined ethereal phase was washed with water, dried (Na_2SO_4) and evaporated. The crude product crystallised soon and could be used without further purification. Yield: 3.1 g (85%).

^1H NMR (300 MHz, CDCl_3) δ 7.45 (dd, 1H), 7.30 (dd, 1H), 4.85 (s, 2H), 2.10 (broad, 1H)

(iv) 4-Cyano-2,5-difluorobenzyl methanesulfonate

To an ice cooled solution of 2,5-difluoro-4-hydroxymethylbenzonitrile (3.10 g, 0.0183 mol; see step (iii) above) and methanesulfonyl chloride (2.21 g, 0.0192 mol) in 60 mL of methylene chloride was added triethyl amine (1.95 g, 0.0192 mol) with stirring. After 1.5 h at 0°C the mixture was washed with water, dried (Na_2SO_4) and evaporated. The product could be used without further purification. Yield: 4.5 g (99%).

^1H NMR (300 MHz, CDCl_3) δ 7.45-7.35 (m, 2H), 5.32 (s, 2H), 3.13 (s, 3H)

(v) 4-Azidomethyl-2,5-difluorobenzonitrile

A mixture of 4-cyano-2,5-difluorobenzyl methanesulfonate (4.5 g, 0.0182 mol; see step (iv) above) and sodium azide (2.0 g, 0.031 mol) in 20 mL of water and 40 mL of DMF was stirred at room temperature for 2 h. It was subsequently poured into 300 mL of water and

extracted three times with diethyl ether. The combined ethereal phase was washed several times with water, dried (Na_2SO_4) and evaporated. A small sample was evaporated for NMR purposes and the product crystallised. The rest was evaporated cautiously but not until complete dryness. Yield (theoretically 3.5 g) is assumed to be almost quantitative based on NMR and analytical HPLC.

^1H NMR (500 MHz, CDCl_3) δ .38 (dd, 1H), 7.32 (dd, 1H), 4.54 (s, 2H)

(vi) 4-Aminomethyl-2,5-difluorobenzonitrile

This reaction was carried out according to the procedure described in *J. Chem. Res. (M)* (1992) 3128. To a suspension of 300 mg of 10% Pd/C (50% moisture) in 20 mL of water was added a solution of sodium borohydride (0.779 g, 0.0206 mol) in 20 mL of water. Some gas evolution resulted. 4-Azidomethyl-2,5-difluorobenzonitrile (1.00 g, 5.15 mmol; from step (v) above) was dissolved in 60 mL of THF and added to the aqueous mixture on an ice bath. The mixture was stirred for 1.5 h whereafter 10 mL of 2M HCl was added and the mixture was filtered through Celite. The Celite was rinsed with more water and the combined aqueous phase was washed with EtOAc and subsequently made alkaline with 2M NaOH. Extraction three times with methylene chloride followed and the combined organic phase was washed with water, dried (Na_2SO_4) and evaporated. Yield: 0.47 g (54%).

^1H NMR (300 MHz, CDCl_3) δ 7.39 (dd, 1H), 7.29 (dd, 1H), 3.99 (s, 2H), 1.45 (broad, 2H)

(vii) 2,5-Difluoro-4-*tert*-butoxycarbonylaminomethylbenzonitrile

A solution of 4-aminomethyl-2,5-difluorobenzonitrile (0.46 g, 2.7 mmol; see step (vi) above) and di-*tert*-butyl dicarbonate (0.60 g, 2.7 mmol) in 10 mL of THF was stirred overnight. The THF was evaporated and the residue was partitioned between water and EtOAc. The organic layer was washed with water, dried (Na_2SO_4) and evaporated. The product could be used without further purification. Yield: 0.71 g (97%).

^1H NMR (300 MHz, CDCl_3) δ 7.35-7.2 (m, 2H), 5.11 (broad triplet, 1H), 4.38 (d, 2H), 1.45 (s, 9H)

(viii) Boc-Pab(2,5-diF)(OH)

A mixture of 2,5-difluoro-4-*tert*-butoxycarbonylaminomethylbenzonitrile (0.70 g, 2.6 mmol; see step (vii) above), hydroxylamine hydrochloride (0.54 g, 7.8 mmol) and triethylamine (0.79 g, 7.8 mmol) in 10 mL of ethanol was stirred at room temperature for 6 days. It was then partitioned between water and methylene chloride. The aqueous layer was extracted with methylene chloride and the combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. The product could be used without further purification. Yield: 0.72 g (92%).

¹H NMR (500 MHz, CD₃OD) δ 7.27 (dd, 1H), 7.12 (dd, 1H), 4.29 (s, 2H), 1.47 (s, 9H)

(ix) Boc-Pab(2,5-diF) x HOAc

This reaction was carried out according to the procedure described by Judkins *et al*, *Synth. Comm.* (1998) 4351. Boc-Pab(2,5-diF)(OH) (0.70 g, 2.3 mmol; see step (viii) above), acetic anhydride (0.25 g, 2.4 mmol) and 230 mg of 10% Pd/C (50% moisture) in 70 mL of acetic acid was hydrogenated at 5 atm pressure for 2.5 h. The mixture was filtered through Celite and evaporated. The residue was freeze dried from acetonitrile and water. The product could be used without further purification in the next step. Yield: 0.80 g (100%).

¹H NMR (500 MHz, CD₃OD) δ 7.49 (dd, 1H), 7.31 (dd, 1H), 4.33 (s, 2H), 1.91 (s, 3H), 1.46 (s, 9H)

(x) Boc-Pab(2,5-diF)(Teoc)

To a suspension of Boc-Pab(2,5-diF) x HOAc (0.80 g, 2.3 mmol; see step (ix) above) in 50 mL of THF was added 2-(trimethylsilyl)ethyl p-nitrophenyl carbonate (0.85 g, 3.0 mmol). A solution of potassium carbonate (0.80 g, 5.8 mmol) in 10 mL of water was added dropwise. The mixture was stirred overnight. The excess Teoc reagent was destroyed by addition of glycine (0.100 g) and potassium carbonate (0.75 g) to the solution, letting it react for an additional 2 h. The THF was evaporated and the residue was partitioned between water and methylene chloride. The aqueous layer was extracted with methylene chloride and the combined organic phase was washed with water, dried (Na₂SO₄) and

evaporated. Flash chromatography on silica gel with heptane:EtOAc (2:1) gave 0.72 g (72%) of pure compound.

¹H NMR (400 MHz, CDCl₃) δ 8.00 (dd, 1H), 7.15 (dd, 1H), 4.98 (broad, 1H), 4.36 (bd, 2H), 4.24 (m, 2H), 1.45 (s, 9H), 1.12 (m, 2H), 0.07 (s, 9H)

(xi) H-Pab(2,5-diF)(Teoc) x 2 HCl

Boc-Pab(2,5-diF)(Teoc) (0.38 g, 0.88 mmol; see step (x) above) was dissolved in 50 mL of EtOAc saturated with HCl(g). The mixture was left for 30 min and evaporated.

¹H NMR (500 MHz, CD₃OD) δ 7.75-7.6 (m, 2H), 4.46 (m, 2H), 4.3 (s, 2H), 1.15 (m, 2H), 0.07 (s, 9H)

(xii) Boc-Aze-Pab(2,5-diF)(Teoc)

To a stirred solution of Boc-Aze-OH (0.189 g, 0.94 mmol), H-Pab(2,5-diF)(Teoc) x 2 HCl (0.36 g, 0.89 mmol; see step (xi) above) and PyBOP (0.54 g, 1.03 mmol) in 5 mL of DMF was added diisopropylethyl amine (0.49 g, 3.8 mmol) and the mixture was allowed to react overnight. The resultant was then poured into aqueous sodium bicarbonate and extracted three times with EtOAc. The combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. Flash chromatography on silica gel with heptane:EtOAc (3:7) gave a sufficiently pure compound. Yield: 0.25 g (48%).

¹H NMR (500 MHz, CDCl₃) δ 7.98 (dd, 1H), 7.13 (dd, 1H), 4.69 (m, 1H), 4.53 (m, 2H), 4.22 (m, 2H), 3.92 (m, 1H), 3.79 (m, 1H), 2.55-2.35 (m, 2H), 1.44 (s, 9H), 1.11 (m, 2H), 0.06 (s, 9H)

(xiii) H-Aze-Pab(2,5-diF)(Teoc) x 2 HCl

Boc-Aze-Pab(2,5-diF)(Teoc) (0.25 g, 0.49 mmol; see step (xii) above) was dissolved in 50 mL of EtOAc saturated with HCl(g). The mixture was left for 30 min. and evaporated.

The product was used in the next step without further purification. Yield: 0.23 g (97%).

¹H NMR (400 MHz, CD₃OD) δ 7.59 (dd, 1H), 7.47 (dd, 1H), 5.14 (m, 1H), 4.54 (m, 2H), 4.48 (m, 2H), 4.15 (m, 1H), 3.96 (m, 1H), 2.87 (m, 1H), 2.56 (m, 1H), 1.17 (m, 2H), 0.05 (s, 9H)

(xiv) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(Teoc)

To a solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)OH (0.12 g, 0.47 mmol; see Example 1(viii) above), H-Aze-Pab(2,5-diF)(Teoc) x 2 HCl (0.23 g, 0.47 mmol; see step (xiii) above) and PyBOP (0.27 g, 0.52 mmol) in 10 mL of DMF was added diisopropylethyl amine (0.245 g, 1.90 mmol), and the mixture was stirred overnight. The resultant was poured into water and extracted three times with EtOAc. The combined organic phase was washed with water, dried (Na₂SO₄) and evaporated. Flash chromatography on silica gel with EtOAc gave 100 mg of a pure fraction and 30 mg of a 90% pure fraction. Total yield: 0.13 g (41%).

¹H NMR (400 MHz, CDCl₃) δ 9.80 (broad, 1H), 8.05 (bt, 1H), 7.94 (dd, 1H), 7.20 (m, 1H), 7.2-7.1 (m, 2H), 7.02 (m, 1H), 6.54 (t, 1H), 4.93 (s, 1H), 4.91 (m, 1H), 4.51 (m, 2H), 4.28 (broad, 1H), 4.23 (m, 2H), 4.13 (m, 1H), 3.74 (m, 1H), 2.69 (m, 1H), 2.43 (m, 1H), 1.73 (broad, 1H), 1.11 (m, 2H), 1.11 (s, 9H)

(xv) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(Teoc) (60 mg (0.093 mmol) of the pure fraction from step (xiv) above) was dissolved in 3 mL of TFA and left at room temperature for 1 h. The TFA was evaporated and the residue was freeze-dried from water and acetonitrile to produce 55 mg (96%) of the title compound as its TFA salt, purity: >99%.

¹H NMR (500 MHz, CD₃OD mixture of rotamers) δ 7.55-7.3 (m, 3H), 7.2-7.1 (m, 2H), 6.88 (t, 1H, major rotamer), 6.86 (t, 1H, minor rotamer), 5.22 (m, 1H, minor rotamer), 5.20 (s, 1H, major rotamer), 5.13 (s, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.6-4.45 (m, 2H), 4.36 (m, 1H, major rotamer), 4.19 (m, 1H, major rotamer), 4.07 (m, 1H, minor


rotamer), 3.98 (m, 1H, minor rotamer), 2.70 (m, 1H, minor rotamer), 2.54 (m, 1H, major rotamer), 2.28 (m, 1H, major rotamer), 2.14 (m, 1H, minor rotamer)

^{13}C -NMR (75 MHz; CD_3OD): (carbonyl and/or amidine carbons, mixture of rotamers) δ 173.0, 172.6, 172.1, 172.0, 162.4

5 APCI-MS: $(M + 1) = 503/505 \text{ m/z}$.

Example 42

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(OMe)



(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze Pab(2,5-diF)(OMe,Teoc)

A mixture of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(Teoc) (40 mg, 0.062 mmol; see Example 41(xiv) above) and O-methyl hydroxylamine hydrochloride (58 mg, 0.70 mmol) in 5 mL of acetonitrile was heated at 70°C for 2 h. The solvent was evaporated and the residue was partitioned between water and EtOAc. The aqueous layer was extracted with EtOAc and the combined organic phase was washed with water, dried (Na_2SO_4) and evaporated. The product could be used without further purification. Yield: 35 mg (84%).

20 ^1H NMR (600 MHz, CDCl_3) δ 7.99 (bt, 1H), 7.72 (s, 1H), 7.20 (m, 1H) 7.15-7.1 (m, 1H), 7.07 (dd, 1H), 7.01 (m, 1H), 6.53 (t, 1H), 4.90 (s, 1H), 4.88 m, 1H), 4.48 (m, 2H), 4.2-4.1 (m, 3H), 3.95 (s, 3H), 3.67 (m, 1H), 2.68 (m, 1H), 2.41 (m, 1H), 0.97 (m, 2H), 0.07 (s, 9H)

25 (ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(OMe)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(OMe,Teoc) (35 mg, 0.052 mmol; see step (i) above) was dissolved in 3 mL of TFA and allowed to react for 30 min. The TFA was evaporated and the residue freeze-dried from water and acetonitrile to give 29 mg (99%) of the title compound. Purity: 97%.

¹H NMR (300 MHz, CDCl₃) δ 8.01 (bt, 1H), 7.45 (dd, 1H), 7.20 (m, 1H), 7.15 (m, 1H), 7.09 (dd, 1H), 7.02 (m, 1H), 6.54 (t, 1H), 5.2-5.0 (m, 2H), 4.95-4.85 (m, 2H), 4.6-4.4 (m, 2H), 4.25 (broad, 1H), 4.13 (m, 1H), 3.90 (s, 3H), 3.71 (m, 1H), 2.69 (m, 1H), 2.43 (m, 1H)

¹³C-NMR (75 MHz; CDCl₃): (carbonyl and/or amidine carbons) δ 173.0, 170.9, 152.6
APCI-MS: (M + 1) = 533/535 m/z.

Example 43

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OEt)

(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OEt, Teoc)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (55 mg, 0.090 mmol; see Example 1(ix) above) and *O*-ethylhydroxyl amine hydrochloride (53 mg, 0.54 mmol) were dissolved in 4 mL of THF. The mixture was stirred at 60°C for 5 h. The solvent was evaporated. The residue was chromatographed on silica gel, eluting with methylene chloride:methanol (95:5) to afford 55 mg (93%) of the sub-title compound.

¹H-NMR (400 MHz; CDCl₃) : δ 7.84 (bt, 1H), 7.59 (bs, 1H), 7.47 (bd, 1H), 7.29 (bd, 1H), 7.21 (m, 1H), 7.14 (m, 1H), 7.02 (m, 1H), 6.53 (t, 1H), 4.90 (s, 1H), 4.86 (m, 1H), 4.55-4.4 (m, 2H), 4.25-4.1 (m, 5H), 3.69 (m, 1H), 2.66 (m, 1H), 2.41 (m, 1H), 1.33 (t, 3H), 0.98 (m, 2H), 0.02 (s, 9H)

(ii) Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OEt)

To an ice-cold solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OEt, Teoc) (55 mg, 0.084 mmol; see step (i) above) in 0.5 mL of methylene chloride was added 3 mL of TFA. The mixture was stirred (ice-bath) for 160 minutes. The material was purified using

preparative HPLC. The fractions of interest were pooled and freeze-dried (2x), yielding 20 mg (47%) of the title compound.

¹H-NMR (400 MHz; CD₃OD) rotamers: δ 7.59 (bd, 2H), 7.35 (m, 1H), 7.32 (bd, 2H),
5 7.25-7.1 (m, 2H), 6.89 (t, 1H, major rotamer), 6.86 (t, 1H, minor rotamer), 5.18 (s, 1H, major rotamer), 5.18 (m, 1H, minor rotamer), 5.11 (s, 1H, minor rotamer), 4.77 (m, 1H), 4.5-4.3 (m, 3H), 4.2-3.9 (m, 3H), 2.67 (m, 1H, minor rotamer), 2.52 (m, 1H, major rotamer), 2.28 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer), 1.28 (t, 3H)
¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 172.4,
10 171.9, 171.4, 153.8, 152.3
MS (m/z) 509 (M - 1)⁻, 511 (M + 1)⁺

Example 44

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OnPr)

15

(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OnPr, Teoc)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (53 mg, 0.087 mmol; see Example 1(ix) above) and *O*-*n*-propylhydroxyl amine hydrochloride, 58 mg (0.52 mmol) were dissolved in 4 mL of THF. The mixture was stirred at 60°C for 5
20 h. The solvent was evaporated. The residue was chromatographed on silica gel, eluting with methylene chloride:methanol (95:5) to afford 51 mg (88%) of the sub-title compound.

¹H-NMR (400 MHz; CDCl₃): δ 7.84 (m, 1H), 7.59 (bs, 1H), 7.47 (bd, 2H), 7.28 (bd, 2H),
25 7.21 (m, 1H), 7.14 (m, 1H), 7.02 (m, 1H), 6.53 (t, 1H), 4.90 (s, 1H), 4.85 (m, 1H), 4.55-4.4 (m, 2H), 4.2-4.05 (m, 5H), 3.69 (m, 1H), 2.65 (m, 1H), 2.41 (m, 1H), 1.74 (m, 2H), 1.05-0.95 (m, 5H), 0.03 (s, 9H)

(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OnPr)

To an ice-cold solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OnPr, Teoc) (51 mg, 0.078 mmol; see step (i) above) in 0.5 mL of methylene chloride was added 3 mL of TFA. The mixture was stirred (ice-bath) for 110 minutes. The material was purified using
30

preparative HPLC. The fraction of interest was evaporated and freeze-dried, yielding 20 mg (47%) of the title compound.

¹H-NMR (500 MHz; CD₃OD) rotamers: δ 7.61 (bd, 2H), 7.38 (m, 1H), 7.35 (bd, 2H), 7.22 (m, 1H, major rotamer), 7.18 (m, 1H), 7.15 (m, 1H, minor rotamer), 6.92 (t, 1H, major rotamer), 6.89 (t, 1H, minor rotamer), 5.20 (s, 1H, major rotamer), 5.20 (m, 1H, minor rotamer), 4.80 (m, 1H, major rotamer), 4.5-4.4 (m, 2H, including minor rotamer corresponding to major at 4.37), 4.37 (m, 1H, major rotamer), 4.18 (m, 1H, major rotamer), 4.09 (m, 1H, minor rotamer), 3.99 (m, 2H), 2.70 (m, 1H, minor rotamer), 2.54 (m, 1H, major rotamer), 2.30 (m, 1H, major rotamer), 2.18 (m, 1H, minor rotamer), 1.73 (m, 2H), 1.01 (t, 3H)

¹³C-NMR (125 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 171.4, 153.8, 152.3

MS (m/z) 523 (M - 1)⁻, 525 (M + 1)⁺

Example 45

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OiPr)

(i) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OiPr, Teoc)

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Teoc) (50 mg, 0.082 mmol; see Example 1(ix) above) and *O*-*i*-propylhydroxyl amine hydrochloride, 55 mg (0.49 mmol) were dissolved in 4 mL of THF. The mixture was stirred at 60°C for 5 h. The solvent was evaporated. The residue was chromatographed on silica gel, eluting with methylene chloride:methanol (95:5) to afford 46 mg (84%) of the sub-title compound.

¹H-NMR (400 MHz; CDCl₃) : δ 7.84 (m, 1H), 7.57 (bs, 1H), 7.48 (bd, 2H), 7.29 (bd, 2H), 7.21 (m, 1H), 7.14 (m, 1H), 7.02 (m, 1H), 6.53 (t, 1H), 4.91 (s, 1H), 4.87 (m, 1H), 4.55-4.45 (m, 2H), 4.42 (m, 1H), 4.2-4.1 (m, 3H), 3.69 (m, 1H), 2.66 (m, 1H), 2.42 (m, 1H), 1.30 (d, 6H), 0.98 (m, 2H), 0.02 (s, 9H)

(ii) Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OiPr)

To an ice-cold solution of Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OiPr, Teoc) (46 mg, 0.069 mmol; see step (i) above) in 0.5 mL of methylene chloride was added 3 mL of TFA. The mixture was stirred (ice-bath) for 150 minutes. The material was purified using preparative HPLC. The fraction of interest was evaporated and freeze-dried (2x), yielding
5 22 mg (58%) of the title compound.

¹H-NMR (400 MHz; CD₃OD) rotamers: δ 7.59 (d, 2H), 7.35 (m, 1H), 7.32 (d, 2H), 7.19 (m, 1H, major rotamer), 7.15 (m, 1H), 7.12 (m, 1H, minor rotamer), 6.89 (t, 1H, major rotamer), 6.86 (t, 1H, minor rotamer), 5.18 (s, 1H, major rotamer), 5.18 (m, 1H, minor
10 rotamer), 5.12 (s, 1H, minor rotamer), 4.78 (m, 1H, major rotamer), 4.5-3.9 (m, 5H), 2.67 (m, 1H, minor rotamer), 2.52 (m, 1H, major rotamer), 2.28 (m, 1H, major rotamer), 2.15 (m, 1H, minor rotamer), 1.26 (d, 6H).

¹³C-NMR (100 MHz; CD₃OD): (carbonyl and/or amidine carbons, rotamers) δ 171.9, 171.4, 153.6.

15 MS (m/z) 523 (M - 1)⁻, 525 (M + 1)⁺

Example 46

The title compounds of Examples 3, 6, 9, 10, 13 to 15, 17, 19, 21, 23, 25, 27, 28, 32, 34, 36, 38, 39 and 41 were tested in Test A above and were found to exhibit IC₅₀TT values of
20 less than 3.5 μM. Those of Examples 3, 6, 9, 10, 13, 15, 17, 19, 21, 23, 27, 32, 34 and 39 were found to exhibit values of less than 0.02 μM; those of Examples 25 and 28 less than 0.03 μM, that of Example 14 less than 0.04 μM; and those of Examples 38 and 41 less than 0.15 μM.

25 Example 47

The title compounds of Examples 3, 6, 13, 15, 17, 19, 21, 23, 25, 27, 28, 32 and 34 were tested in Test D above and were found to exhibit an IC₅₀ APTT value of less than 1 TM.

Example 48

30 The title compounds of Examples 1, 2, 4, 5, 7, 12, 16, 18, 20, 22, 24, 26, 29, 30, 33 and 43 to 45 were tested in Test E above and were found to exhibit oral and/or parenteral bioavailability in the rat as the corresponding active inhibitor (free amidine).

Example 49

- 5 Title compounds of Examples 1, 2, 7, 8, 11, 12, 16, 18, 20, 22, 24, 26, 29, 33, 37, 40, 43 and 45 were tested in Test G above and were found to be converted to the corresponding active inhibitor (free amidine) in liver microsomes from humans and from rats.

Abbreviations

10

Ac = acetyl

AcOH = acetic acid

APCI = atmospheric pressure chemical ionisation (in relation to MS)

API = atmospheric pressure ionisation (in relation to MS)

15

aq. = aqueous

AUC = area under the curve

Aze = (*S*)-azetidine-2-carboxylate (unless otherwise specified)

AzeOH = azetidine-2-carboxylic acid

Bn = benzyl

20

Boc = *tert*-butoxycarbonyl

BSA = bovine serum albumin

Bu = butyl

Bzl = benzyl

CI = chemical ionisation (in relation to MS)

25

d = day(s)

DCC = dicyclohexyl carbodiimide

DIBAL-H = di-isobutylaluminium hydride

DIPEA = diisopropylethylamine

DMAP = 4-(*N,N*-dimethyl amino) pyridine

30

DMF = dimethylformamide

DMSO = dimethylsulfoxide

DVT = deep vein thrombosis

- EDC = 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
e.e. = enantiomeric excess
Et = ethyl
ether = diethyl ether
5 EtOAc = ethyl acetate
EtOH = ethanol
Et₂O = diethyl ether
h = hour(s)
HATU = *O*-(azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate
10 HBTU = [*N,N,N',N'*-tetramethyl-*O*-(benzotriazol-1-yl)uronium hexafluorophosphate]
HCl = hydrochloric acid, hydrogen chloride gas or hydrochloride salt (depending on context)
Hex = hexanes
HOAc = acetic acid
15 HPLC = high performance liquid chromatography
LC = liquid chromatography
Me = methyl
MEM = methoxyethoxymethyl
MeOH = methanol
20 min = minute(s)
MS = mass spectroscopy
MTBE = methyl *tert*-butyl ether
NADH = nicotinamide adenine dinucleotide, reduced form
NADPH = nicotinamide adenine dinucleotide phosphate, reduced form
25 NIH = National Institute of Health (US)
NIHU = National Institute of Health units
NMR = nuclear magnetic resonance
OAc = acetate
Pab = *para*-amidinobenzylamino
30 H-Pab = *para*-amidinobenzylamine
Ph = phenyl
Pr = propyl

- Pro = (S)-prolinyl
PyBOP = (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate
QF = tetrabutylammonium fluoride
RedAl = sodium bis(2-methoxyethoxy)aluminium hydride
5 RPLC = reverse phase high performance liquid chromatography
rt/RT = room temperature
SOPs = standard operating procedures
TBTU = [N,N,N',N'-tetramethyl-O-(benzotriazol-1-yl)uronium tetrafluoroborate]
TEA = triethylamine
10 Teoc = 2-(trimethylsilyl)ethoxycarbonyl
TEMPO = 2,2,6,6-tetramethyl-1-piperidinyloxy free radical
TFA = trifluoroacetic acid
THF = tetrahydrofuran
THP = tetrahydropyranyl
15 TLC = thin layer chromatography
TMSCl = trimethylsilyl chloride
TMSCN = trimethylsilyl cyanide
UV = ultraviolet
Z = benzyloxycarbonyl

20

Prefixes *n*, *s*, *i* and *t* have their usual meanings: normal, secondary, iso and tertiary. The prefix *c* means cyclo.

25

Claims

1. A combination product comprising:

- 5 (a) a compound of claim 1 in PCT/SE01/02657 or a pharmaceutically-acceptable derivative thereof; and
- (b) (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof)
- 10 wherein each of components (a) and (b) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

2. A combination product as claimed in Claim 1 which comprises a pharmaceutical formulation including a compound of claim 1 in PCT/SE01/02657 or a compound of claim
- 15 20 in PCT/SE01/02657 or sub-set 1, 2 or 3 of the compounds of claim 20 or a pharmaceutically-acceptable derivative thereof, and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof) or a pharmaceutically-acceptable derivative thereof, in admixture with a pharmaceutically-acceptable adjuvant,
- 20 diluent or carrier.

3. A combination product as claimed in Claim 1 which comprises a kit of parts comprising components:

- (a) a pharmaceutical formulation including a compound of claim 1 in PCT/SE01/02657 or
- 25 a compound of claim 20 in PCT/SE01/02657 or sub-set 1,2 or 3 of the compounds of claim 20 or a pharmaceutically-acceptable derivative thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; and
- (b) a pharmaceutical formulation including (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or

(3) Compound A or B or C or D (or pharmaceutically-acceptable salts thereof), in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier, which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

5

4. A kit of parts as claimed in Claim 3, wherein components (a) and (b) are suitable for sequential, separate and/or simultaneous use in the treatment of a condition where anticoagulant therapy is indicated.

10

5. A combination product as claimed in any one of Claims 1 to 4, which comprises Compound A or B or C or D (or pharmaceutically-acceptable salts thereof).

6. A combination product as claimed in any one of Claims 1 to 5 wherein the compound of PCT/SE01/02657 is selected from:

15

Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Cl)(5-OCHF₂)-(S)CH(CH₂OH)C(O)-Aze-Pab;

Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab;

20

Ph(3-OCHF₂)-(R)CH(OH)-CO-Aze-Pab;

Ph(3-OCF₃)-(R)CH(OH)-CO-Aze-Pab;

Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab;

25

Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Br)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab;

Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab;

30

Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Pro-Pab;

Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((2-amidino)-5-pyridinyl);

- Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-amidino)-2-pyrimidinyl);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(3-F);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF);
 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF).
 5 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OEt);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OnPr);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OiPr);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OcBu);
 10 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(COOCpentyl);
 Ph(3-Cl)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(Z);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂-3-(5-Me-isoxazole));
 15 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂-3-pyridine);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OiBu);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OEt);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Obn);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OcHexyl);
 20 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OcBu);
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(OCH₂CH₂OPh(3-CF₃));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Obn(4-Cl));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Obn(3-MeO));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Obn(2-Br));
 25 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(Obn(4-Me));
 Ph(3-Cl)(5-OCF₃)-(R)CH(OH)C(O)-Aze-Pab(O-4-heptyl);
 Ph(3-Cl)(5-OCF₃)-(S)CH(CH₂OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH₂CF₃)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 30 Ph(3-Cl)(5-OCH₂F)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OMe);
 Ph(3-Cl)(5-OCH(CH₂F)₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);

Ph(3-F)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
Ph(3-Br)(5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(OMe);
Ph(3-Cl, 5-OCH₂CHF₂)-(R)CH(OH)C(O)-Aze-Pab(OH);
Ph(3-Cl, 5-OCH₂CH₂F)-(R)CH(OH)C(O)-Aze-Pab(OH);
5 Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Pro-Pab(OMe);
Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((2-methoxy-amidino)-5-pyridinyl);
Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-NH-CH₂-((5-methoxy-amidino)-2-pyrimidinyl);
Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,6-diF)(OMe); or
Ph(3-Cl, 5-OCHF₂)-(R)CH(OH)C(O)-Aze-Pab(2,5-diF)(OMe) and pharmaceutically-
10 acceptable derivatives thereof.

7. A method of making a kit of parts as defined in any one of Claims 3 to 6, which method comprises bringing a component (a), as defined in any one of Claims 3 to 6, into association with a component (b), as defined in any one of Claims 3 to 6, thus rendering
15 the two components suitable for administration in conjunction with each other.

8. A kit of parts comprising:
(I) one of components (a) and (b) as defined in any one of Claims 3 to 6; together with
(II) instructions to use that component in conjunction with the other of the two
20 components.

9. A method of treatment arrhythmia, which comprises administration of a combination product as defined in any one of Claims 1 to 8 to a patient suffering from, or susceptible to, such a condition.
25

10. The use of a combination product as defined in any one of Claims 1 to 8 for the manufacture of a medicament for the treatment or prophylaxis of a condition where anticoagulant therapy is indicated.

11. The use of a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in PCT/SE01/02657 (or derivative thereof) or a pharmaceutically-acceptable derivative thereof and (1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-
s acceptable salts thereof) for the manufacture of a medicament for the treatment or prophylaxis of a condition where anticoagulant therapy is indicated.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
21

ABSTRACT

There is provided a combination product comprising:

- (1) a compound of claim 1 in PCT/SE01/02657 or a compound of claim 20 in
5 PCT/SE01/02657 (or derivative thereof) or a pharmaceutically-acceptable derivative
thereof; and
(1) a compound as defined in claim 1 of WO 01/28992 or (2) a compound of Claim 34 of
WO 01/28992 or (3) Compound A or B or C or D (or pharmaceutically-acceptable salts
thereof)
10 for use in treating arrhythmia or a coagulation controlled complication thereof.

PCT/SE01/02657